

## Solute – Solvent and Solute – Solute Interactions of Tetrapentylammonium Iodide in Dimethylsulphoxide – Acetone Systems at Different Temperatures

Baljeet Singh Patial

Department of Chemistry, BTC DAV College, Banikhet (Dalhousie), Distt. Chamba, H.P.(India)

---

**Abstract:** Ultrasonic velocity, viscosity and density studies on solution of tetrapentylammonium iodide ( $\text{Pen}_4\text{NI}$ ) in Dimethylsulphoxide (DMSO) and Dimethylsulphoxide + Acetone solvent mixtures containing 50,60,70,80,90 and 100 mol % of DMSO at 298 and 308K have been reported. From the velocity, viscosity and density data values, various parameters namely, the adiabatic compressibility ( $\beta$ ), apparent molar compressibility ( $\phi_k$ ), apparent molar volume ( $\phi_v$ ), limiting apparent molar compressibility ( $\phi_k^\circ$ ), limiting apparent molar volume ( $\phi_v^\circ$ ), free volume ( $V_f$ ), internal pressure ( $\pi_i$ ), relaxation time ( $\tau$ ) and viscosity B-coefficient have been calculated. All these parameters have been discussed separately to throw light on the solute-solvent and solvent-solvent interactions.

**Keywords:** Adiabatic compressibility, apparent molar compressibility, apparent molar volume, free volume, B-coefficient.

---

### I. Introduction

The principles related to acoustic behavior, viscosity etc., can be used to study the physicochemical behavior and molecular interactions such as ion – solvent interaction and solvent-solvent interactions involved in any electrolyte system<sup>1-6</sup>. The knowledge of solvation behaviour of a species is very essential to understand the solution chemistry<sup>1</sup>. Studies of densities, viscosities and ultrasonic speeds of electrolytic solutions are of great use in characterizing the structure and properties of solutions. Recently, solute-solute and solute-solvent interactions for the tetraalkylammonium salts have been reported<sup>7-9</sup>.

Various types of interactions exist between the solutes in the solutions, and these solute-solute and solute-solvent interactions are of current interest in all branches of chemistry. These interactions provide a better understanding of the nature of the solute and solvent i.e., whether the solute modifies or distorts the structure of the solvent.

The present investigation reports the ultrasonic velocity, density and viscosity studies of tetrapentylammonium iodide ( $\text{Pen}_4\text{NI}$ ) in Dimethylsulphoxide (DMSO) and Dimethylsulphoxide (DMSO) – Acetone (Ac) mixtures at 298 and 308K. From these experimental data, the number of thermodynamic parameters viz. the adiabatic compressibility ( $\beta$ ), apparent molar compressibility ( $\phi_k$ ), apparent molar volume ( $\phi_v$ ), limiting apparent molar compressibility ( $\phi_k^\circ$ ), limiting apparent molar volume ( $\phi_v^\circ$ ), free volume ( $V_f$ ), internal pressure ( $\pi_i$ ), relaxation time ( $\tau$ ) and viscosity B-coefficient have been calculated. These parameters have been used to interpret various molecular interactions occurring in solutions at different temperatures.

### II. Experimental

Dimethyl sulphoxide (extra pure AR grade; S D Fine Chemicals Ltd) was kept over CaO and distilled<sup>10</sup>. Acetone (Ac) of 99.5% purity (extra pure AR grade; S D Fine Chemicals Ltd) was dried over 4Å molecular sieves and distilled<sup>11,12</sup>. Tetrapentylammonium iodide ( $\text{Pen}_4\text{NI}$ ) from Fluka, was dried and used as described earlier<sup>13-16</sup>. Viscosity measurements were carried out as described elsewhere<sup>15-18</sup>. Viscosity values were found to be in good agreement with those reported in literature<sup>13-18</sup>. The densities of pure solvent, solvent system and various electrolytic solutions were measured with the help of a sealable type of pycnometer (supplied by M/s. Harsh & Co., Ambala Cantt.) of 20 cm<sup>3</sup> capacity).

The value of ultrasonic velocity for the conductivity water was found to be 1490 m/s at 298.15 K at 1 MHz, which is in agreement with literature value<sup>19</sup>.

Ultrasonic velocity was measured using interferometer (Model-81, supplied by Mittal Enterprises, New-Delhi) operating at a frequency of 1 MHz, which is a direct and simple device for measuring ultrasonic velocity in liquids.

The viscosities and densities of the above electrolyte in DMSO and DMSO + Ac solvent systems were measured at 298 and 308 K. The overall accuracy of the viscosity and density measurements in this study was estimated to be  $\pm 0.2\%$  and  $\pm 0.1\%$  respectively.

Different parameters such as the adiabatic compressibility ( $\beta$ ), apparent molar compressibility ( $\phi_k$ ), apparent molar volume ( $\phi_v$ ), limiting apparent molar compressibility ( $\phi_k^\circ$ ), limiting apparent molar volume ( $\phi_v^\circ$ ), free volume ( $V_f$ ), internal pressure ( $\pi_i$ ), relaxation time ( $\tau$ ) and viscosity B-coefficient have been calculated at

different temperatures, with the help of ultrasonic velocity ( $u$ ), density ( $\rho$ ) and viscosity( $\eta$ ) values using the following relations<sup>7,20-22</sup>:

**Adiabatic compressibility ( $\beta$ )**

The adiabatic compressibility values for various compositions of the binary solvent mixtures have been calculated from the measured ultrasonic velocities ( $u$ ) and densities ( $\rho$ ):

$$\beta = \frac{1}{u^2\rho}$$

**Apparent molar compressibility ( $\phi_k$ )**

The apparent molar compressibility ( $\phi_k$ ) have been calculated using the relation:

$$\phi_k = \frac{1000}{c\rho_o}(\rho_o\beta - \beta_o\rho) + \frac{\beta_o M}{\rho_o}$$

where  $\beta, \rho$  and  $\beta_o, \rho_o$  are the adiabatic compressibility and density of the solution and solvent, respectively.  $M$  is the molar mass of the solute, and  $c$  is the molar concentration.

**Limiting apparent molar compressibility ( $\phi_k^o$ ):**

The values of the limiting apparent molar compressibility  $\phi_k^o$  of solute solution are obtained by the use of least square treatment to the plots of the use of  $\phi_k$ , apparent molar compressibility of solution versus  $C^{1/2}$  in accordance with Masson's empirical<sup>23</sup>:

$$\phi_k = \phi_k^o + S_k^* + C^{1/2}$$

where  $\phi_k^o$  is the limiting apparent molar compressibility at infinite dilution and  $S_k$  is a constant.

**The apparent molar volume ( $\phi_v$ ):**

The apparent molar volume,  $\phi_v$  is calculated from the density data by using following expression:

$$\phi_v = 1000 \frac{(\rho_o - \rho)}{c\rho_o} + \frac{M}{\rho_o}$$

where  $\rho_o$  and  $\rho$  are densities of solvent and solution respectively;  $M$  is the molar mass of the solute, and  $c$  is the molar concentration.

**Limiting apparent molar volume ( $\phi_v^o$ ):**

The values of the limiting apparent molar volume  $\phi_v^o$  of solute solution are obtained by the use of least square treatment to the plots of the use of  $\phi_v$ , apparent molar volume of solution versus  $C^{1/2}$  in accordance with Masson's empirical<sup>23</sup>:

$$\phi_k = \phi_k^o + S_k^* + C^{1/2}$$

where  $\phi_v^o$  is the limiting apparent molar compressibility at infinite dilution and  $S_v$  is a constant.

**Relaxation Time ( $\tau$ )**

Relaxation can be calculated from viscosity coefficient ( $\eta$ ), density and ultrasonic velocity of binary mixtures and given by

$$\tau = \frac{4\eta}{3\rho u^2}$$

**Free Volume ( $v_f$ )**

The free volume of binary mixture is given by

$$V_f = \left[ \frac{M_{eff} u}{K\eta} \right]^{3/2}$$

where  $K$  is time independent constant whose value is  $4.28 \times 10^9$  in MKS system and  $M_{eff}$  effective molecular weight of the liquid is given by

$$M_{eff} = X_1 M_1 + X_2 M_2$$

where  $X_1$  &  $X_2$  are the mole fraction of first and second components and  $M_1$  &  $M_2$  are the molecular weights of first and second components respectively.

### Internal Pressure ( $\pi_i$ )

Internal pressure is given by

$$\pi_i = \frac{bRT[K'\eta]}{M_{eff}^{\frac{7}{6}} \rho^{\frac{2}{3}}}$$

where,  $b$  is the cubic packing factor which is assumed to be 2 in liquid systems.

$K = 4.28 \times 10^9$  and is independent to the nature of liquid.

$R$  is gas constant.

$\eta$  is the viscosity and  $\rho$  is the density of solution.

### III. Result And Discussion

The experimental values of density, viscosity and ultrasonic velocity of  $Pen_4NI$ , have been measured in DMSO and DMSO + Ac mixtures containing 100,90,80,70,60 and 50 mol% of DMSO in the concentration range (0.05-0.30) mol  $dm^{-3}$  at 298 and 308 K. The calculated values of adiabatic compressibility, apparent molar volume and apparent molar adiabatic compressibility are reported in Table-1. The value of limiting apparent molar compressibility, limiting molar volume, and the constants  $S_k$  and  $S_v$  for tetrapentylammonium iodide ( $Pen_4NI$ ) are presented in Table-2. The values of A- and B- coefficients are given in Table-3 and the calculated values of free volume, internal pressure and relaxation time in Table-4.

It is clear from the Table-1 that the adiabatic compressibility ( $\beta$ ) values of solution decrease with increase in the concentration of electrolyte at all compositions.  $\beta$  – values for a particular solute concentration increase with the increase of a particular solute concentration increase with the increase of acetone content to DMSO + Ac mixture. With increase of temperature,  $\beta$  – values of solution increase, indicates temperature dependence of  $\beta$  and increase of solute-solvent interactions.

The decrease of  $\beta$  with increase of solute concentration can be attributed due to the electrostatic effect of solute on the surrounding solvent molecules, which results in increase of internal pressure and thus solution becomes harder to compress. The decrease of  $\beta$  with increase of concentration of solute in a particular system is indicative of presence of solute-solvent interactions. Similar observations were made by Syalet al<sup>24</sup> and Kumar et al<sup>25</sup>. The increase in  $\beta$  with increase of Ac content in DMSO in DMSO + Ac mixture may be interpreted to the presence of dipole-dipole interaction/association between DMSO and Ac molecules and presence of solute-solvent interaction between electrolyte and solvent system.

**Table-1: Adiabatic compressibility ( $\beta$ ), Apparent molar compressibility ( $\phi_k$ ) and apparent molar volume ( $\phi_v$ ) of  $Pen_4NI$  in DMSO- Ac mixtures.**

C mol $dm^{-3}$	$\beta \times 10^{-10}$ ( $Kg^{-1}m s^{-1}$ )		$\phi_k \times 10^4$ ( $atm^{-1}m^3mol^{-1}$ )		$\phi_v$ ( $m^3mol^{-1}$ )	
	298K	308K	298K	308K	298K	308K
Pure DMSO						
0.00	4.20	4.40	--	--	--	--
0.05	1.20	4.39	141.94	149.01	368.00	372.96
0.10	4.19	4.39	139.01	147.21	365.87	371.70
0.15	4.18	4.38	137.71	145.89	364.85	371.01
0.20	4.17	4.36	136.72	143.39	363.66	370.00
0.25	4.16	4.35	134.74	141.25	362.39	369.24
0.30	4.15	4.34	133.53	139.89	361.06	368.56
90%DMSO						
0.00	4.48	4.75	--	--	--	--
0.05	4.47	4.73	138.20	141.76	362.05	368.76
0.10	4.45	4.71	133.91	140.02	360.98	367.16
0.15	4.43	4.69	131.53	138.86	359.78	366.17
0.20	4.42	4.67	130.63	136.84	358.98	365.33
0.25	4.40	4.65	129.53	133.35	357.69	364.17
0.30	4.39	4.63	127.36	133.23	356.90	362.82
80%DMSO						
0.00	4.81	5.12	--	--	--	--
0.05	4.79	5.10	130.90	136.34	358.15	365.09
0.10	4.77	5.07	128.48	134.38	356.7819	363.78
0.15	4.75	5.04	126.78	131.75	355.57	362.58
0.20	4.73	5.01	125.38	129.06	354.28	361.32
0.25	4.71	4.99	123.92	126.95	353.34	360.19
0.30	4.68	4.96	122.29	125.30	352.31	359.08
70%DMSO						

0.00	5.20	5.53	--	--	--	--
0.05	5.17	5.50	126.57	128.56	355.28	360.36
0.10	5.14	5.46	123.65	126.14	353.74	359.17
0.15	5.11	5.43	121.55	123.72	352.20	357.99
0.20	5.09	5.39	120.59	122.17	350.66	356.81
0.25	5.06	5.36	117.91	119.26	349.27	355.65
0.30	5.03	5.32	116.53	117.20	348.34	354.85
60%DMSO						
0.00	5.63	6.00	--	--	--	---
0.05	5.59	5.96	118.55	120.89	351.21	356.43
0.10	5.55	5.92	115.95	118.96	349.78	354.99
0.15	5.51	5.87	114.53	116.64	348.36	353.66
0.20	5.48	5.83	112.99	113.60	346.96	352.88
0.25	5.44	5.78	111.07	1114.42	345.56	351.74
0.30	5.40	5.74	108.77	109.05	344.55	350.92
50%DMSO						
0.00	6.09	6.54	--	--	--	--
0.05	6.05	6.48	113.71	115.79	346.56	353.69
0.10	6.00	6.43	111.09	113.53	345.19	352.04
0.15	5.96	6.38	110.22	110.84	344.09	350.76
0.20	5.91	6.32	107.52	108.42	343.19	349.98
0.25	5.87	6.27	105.55	106.30	342.42	349.11
0.30	5.82	6.22	103.17	104.20	341.56	348.93

The apparent molar adiabatic compressibility ( $\phi_k$ ) values for Pen<sub>4</sub>NI are positive at all the compositions and decrease with increase of Ac content to DMSO + Ac mixtures. These values show a general decrease with the increase of concentration for a particular composition showing the presence of interactions. From Table-1, it is evident that  $\phi_k$  values increase with increase of temperature for Pen<sub>4</sub>NI in all the composition. A similar trend is shown by Syalet al<sup>24</sup> in EMK+DMF solvent systems.

The limiting apparent molar compressibility  $\phi_k^\circ$  and  $S_k$  for each of electrolytic solutions have computed by least-squares method. From Table-2, the observed  $\phi_k^\circ$  values are large and positive for Pen<sub>4</sub>NI, which are found to be similar as reported earlier<sup>26</sup>.

$\phi_k^\circ$  values decrease with decrease in DMSO content in DMSO + Ac mixtures. Large positive values of  $\phi_k^\circ$  are indicative of solute-solvent and intermolecular interactions. These  $\phi_k^\circ$  values show little temperature dependence showing slight increase in the values with increased in temperature. The corresponding  $S_k$  values which indicative the solute-solute interactions<sup>27</sup> decrease with an increase in temperature.

**Table-2: Limiting apparent molar compressibility ( $\phi_k^\circ$ ), Limiting apparent molar volume ( $\phi_v^\circ$ ) and constant  $S_k$  and  $S_v$  of Pen<sub>4</sub>NI in DMSO- Ac mixtures.**

$X_{DMSO}$	$\phi_k^\circ 10^4$ ( $m^3 atm^{-1} mol^{-1}$ )		$S_k 10^4$		$\phi_v^\circ$ ( $m^3 mol^{-1}$ )		$S_v$	
	298K	308K	298K	308K	298K	308K	298K	308K
1.00	159.78	156.14	-25.12	-29.12	372.63	376.65	-22.63	-19.68
.90	144.11	150.22	-30.86	-31.85	365.90	372.80	-20.12	-18.50
.80	136.86	145.24	-25.76	-36.23	362.59	369.97	-18.14	-18.68
.70	133.20	136.96	-31.14	-34.76	360.49	364.50	-17.56	-17.42
.60	124.86	130.24	-29.04	-37.24	356.21	360.30	-16.00	-17.19
.50	121.12	124.15	-31.22	-36.11	350.01	357.45	-15.30	-16.96

The apparent molar volume ( $\phi_v$ ) behaves in a similar fashion to that of the apparent molar adiabatic compressibility in the salt solution.  $\phi_v^\circ$  is regarded as a measure of solute –solvent interaction. It is evident from the Table-2 that  $\phi_v^\circ$  values are positive for Pen<sub>4</sub>NI in DMSO + AC mixtures, suggesting the presence of strong solute-solvent interactions. Small decrease in  $\phi_v^\circ$  as the amount of Ac in the solution increases indicating the decreasing trend of solute-solvent interactions. The values of  $S_v^\circ$  are large and indicative the presence of weak solute-solute interaction in the solution. Moreover, the values of  $S_v^\circ$  become less negative with increase in Ac content in the system, suggesting increased solute-solute interaction in the present solvent system. In fact, negative  $S_v^\circ$  values are often obtain in solvents high dielectric constant<sup>28</sup>. This is attributed to fact that in solvents of high dielectric constant, like DMSO + Ac system, the salts remain completely ionized, even at fairly high concentrations. As a result, appreciable interionic penetration is likely to occur, giving rise to negative slope for  $\phi_v$  versus  $C^{1/2}$  plots<sup>29</sup>.

**Table-3: Free volume ( $V_f$ ), internal pressure ( $\pi_i$ ) and relaxation time ( $\tau$ ) of Pen<sub>4</sub>NI in DMSO-Ac mixtures.**

c mol dm <sup>-3</sup>	$V_f \times 10^{-8}$ (m <sup>3</sup> mol <sup>-1</sup> )		$\pi_i \times 10^9$ (Pa)		$\tau \times 10^{12}$ (s)	
	298K	308K	298K	308K	298K	308K
Pure DMSO						
0.00	5.02	6.43	7.80	6.57	11.01	9.51
0.05	4.53	5.93	8.04	6.72	11.96	10.21
0.10	4.22	5.62	8.19	6.81	12.52	10.64
0.15	3.96	5.31	8.33	6.90	13.26	11.12
0.20	3.72	5.05	8.47	6.99	13.84	11.66
0.25	3.48	4.77	8.62	7.09	14.50	12.18
0.30	3.32	4.59	8.72	7.15	15.16	12.57
90% DMSO						
0.00	6.64	8.30	7.10	6.02	9.35	8.30
0.05	5.96	7.62	7.32	6.17	10.18	8.84
0.10	5.54	7.18	7.47	6.26	10.64	9.25
0.15	5.18	6.81	7.61	6.35	11.29	9.63
0.20	4.87	6.48	7.74	6.43	11.72	10.02
0.25	4.57	6.31	7.87	6.53	12.39	10.54
0.30	4.36	5.89	7.97	6.58	12.85	10.88
80% DMSO						
0.00	8.26	10.16	6.59	5.63	8.29	7.48
0.05	7.36	9.23	6.96	5.78	8.99	7.98
0.10	6.83	8.68	7.10	5.87	9.47	8.35
0.15	6.37	8.19	7.21	5.96	9.93	8.72
0.20	6.02	7.82	7.35	6.04	10.49	9.03
0.25	5.33	7.40	7.45	6.12	10.92	9.41
0.30		7.11		6.18	11.46	9.70
70% DMSO						
0.00	10.53	12.57	9.06	5.24	7.20	6.67
0.05	9.37	11.46	6.28	5.38	7.82	7.12
0.10	8.68	10.82	6.42	5.46	8.26	7.42
0.15	8.14	10.25	6.54	5.54	8.06	7.73
0.20	7.63	9.67	6.66	5.62	9.08	8.02
0.25	7.20	9.15	6.77	5.70	9.47	8.35
0.30	6.79	8.80	6.88	5.76	9.89	8.60
60% DMSO						
0.00	13.36	15.43	5.60	4.87	6.30	5.95
0.05	11.86	14.02	5.80	5.80	6.85	6.37
0.10	11.02	13.25	5.93	5.93	7.22	6.63
0.15	10.28	12.56	6.05	6.05	7.59	6.58
0.20	9.69	11.90	6.15	6.15	7.92	6.84
0.25	9.12	11.30	6.25	6.25	8.27	7.11
0.30	8.63	10.82	6.35	6.35	7.61	7.38
50% DMSO						
0.00	16.54	18.13	5.20	4.60	5.62	5.50
0.05	14.64	16.39	5.40	4.75	6.11	5.90
0.10	13.54	15.44	5.53	4.83	6.16	6.16
0.15	12.63	14.63	5.64	4.90	6.79	6.41
0.20	11.87	13.85	5.74	4.97	7.10	6.66
0.25	11.18	13.16	5.84	5.04	7.41	6.91
0.30	10.66	12.63	5.91	5.10	7.67	7.12

Free volume can be defined as the average volume in which the central molecule can move inside the hypothetical cell due to repulsion of surrounding molecules. Free volume can also be referred as the void space between the molecules i.e. volume present as holes of monomeric size, due to irregular packing of molecules.

The calculated values of free volume ( $V_f$ ) are given in Table-3 for studied electrolyte in all the compositions at both temperatures.  $V_f$  values in general decrease in magnitude with the increase of concentration of salt. However, with the increase of Ac content in DMSO+ Ac mixture,  $V_f$  values increase. Increase of temperature also increases the magnitude of  $V_f$ .

Internal pressure ( $\pi_i$ ) is the resultant of forces of attraction and repulsion between solute and solvent molecules of the solution. It is evident from the Table-3 that  $\pi_i$  values increase with the increase of solute concentration and decrease with increase of temperature in all composition for Pen<sub>4</sub>NI. Increase of  $\pi_i$  with concentration indicates increase in intermolecular interactions due to the forming of aggregates of solvent molecules around the solute, which affects the structural arrangement of solution system. This may also be attributed to the presence of solute-solvent interactions. Similar behavior has been reported for PVP polymer in DMSO+H<sub>2</sub>O system<sup>30</sup>.

From Table-3, it has been found that the viscous relaxation time ( $\tau$ ) values increase with increase in concentration of solute in the studied solvent systems at both temperatures. Acoustic relaxation time decreases

with the rise in temperature, in accordance with the decrease of density, ultrasonic velocity and viscosity of solution system with the increase of temperature.

Relaxation time values decrease with the increase of acetone contents in DMSO + Ac mixture. This may be accounted for the decrease of dielectric constant of the medium, and change of intermolecular interactions between DMSO and Ac molecules. Increase of  $\tau$  with increase of solute concentration may be attributed to the presence of solute-solvent interactions. Similar results have been reported for tetraalkylammonium salts by Syalet al<sup>17</sup>.

**Table-4: Values of A (dm<sup>2/3</sup> mol<sup>-1/2</sup>) and B (dm<sup>3</sup> mol<sup>-1</sup>) parameters of Jones-Dole equation for Pen<sub>4</sub>NI in DMSO- Ac mixtures.**

X <sub>DMSO</sub>	A x10 <sup>2</sup>		B	
	298K	308K	298K	308K
1.00	10.64	7.42	1.09	0.93
0.90	12.90	9.72	1.09	0.92
0.80	15.39	10.97	1.10	0.93
0.70	16.37	11.60	1.09	0.92
0.60	17.21	12.96	1.09	0.92
0.50	18.34	15.22	1.10	0.91

The viscosity data of the present solutions were analysed by using Jones-Dole equation<sup>31</sup>,

$$\eta_r = \frac{\eta}{\eta_0} = 1 + AC^{1/2} + BC$$

where  $\eta$  and  $\eta_0$ , respectively are the dynamical viscosities of solution and solvent,  $\eta_r$  is the relative viscosity of the solution. A is Falkenhagen coefficient<sup>32</sup> and is a measure of solute-solute interactions theoretically. On the other hand, B, the Jones -Dole coefficient, is empirical and is a function of the solute-solvent interactions. The coefficients A – and B- were calculated by least square fitting of experimental  $\eta$  values in the Jones-Dole equation as given above. The values of A and B thus obtained are listed in Table-4. The values of the A-coefficient are found to be positive in all the cases, indicating strong solute-solute interactions in the concentration range investigated. . Most of the studies in pure and mixed solvents have been positive<sup>16</sup>. The values of B-coefficients for tetraalkylammonium Iodide are positive. This is identical with the general observation<sup>16,33</sup> that B-coefficients are commonly large and almost always positive for salts in non-aqueous solvents. The positive B-coefficients value attributed to strong solute-solvent interaction in the system. From Table-4, it is clear that viscosity B-coefficients for Pen<sub>4</sub>NI is decrease with rise in temperature suggesting the structure-making tendency of Pen<sub>4</sub>NI in the studied solvent system. This found to be consistent with the work reported in literature<sup>14-16</sup>.

## References

- [1]. Marcus Y, Ion solvation, John Wiley, New York(1985).
- [2]. Chauhan, Kumar K, &Patil B.S., Indian J Pure ApplPhys, 51(2013)531.
- [3]. Syal V K, Bhall V &Chauhan S, Indian J Pure ApplPhys, 81(1995)276.
- [4]. Syal V K, Gautam R &Chauhan S, Indian J Pure &ApplPhys, 36(1998)108.
- [5]. Chauhan S, Syal V K, Chauahn M S & Sharma P, ActaAcoustica United with Acoustica, 93(2007)566.
- [6]. Mehra R & Gaur A K, J Chem, Eng Data, 53(2008)863.
- [7]. Hemalatha B., Vasantharani P &Senthilkumar; International J AdvEng& Tech,6(2013)795.
- [8]. Nikam P S, Hasan Mehdi, Pawar T B, Sawant A B; Indian J Pure Appl Phys,42(2004)172.
- [9]. Nikam P S, Nikhumbh A B, J ChemEng Data,47(2002)400.
- [10]. Syal V K,Chauhan S &Chauhan M S, Z. Phys. Chem.(NF),49(1988)159.
- [11]. Syal V K, Chauhan S &Kumari U, Ind. J. Pure & Appl. Phys.,43(2005)844.
- [12]. Reddick J A, Burger W B &Sankano T K,Technique in Chemistry Vol.II 4th edition(John Wiley, N.Y.,1986).
- [13]. Syal V K, Patial B S &Chauhan S, Acoust. Lett., 23(2000)137.
- [14]. Patial B S, IOSR-JAC, 8(2)(Feb.2015)01-04
- [15]. Patial B S, IOSR-JAC, 7(Dec.2014)01-05.
- [16]. Patial B S , Chauhan S, Chauhan M S &Syal V K, Ind. J. of Chem.,41A(2002)2039
- [17]. Syal V K, Patial B &Chauhan S, Ind. J. Pure and Appl. Phys.,37(1999)366.
- [18]. Syal V K, Bist P and Chauhan M S, Proc. Nat. Acad. Sci., India,67A(1977)1.
- [19]. Syal V.K. , LalG., Bist P. &Chauhan S., J. Mol. Liq., 63(1995)317.
- [20]. Syal V.K., Chauhan Anita &chauhanSuvarcha, J. Pure Appl. Ultrason,27(2005)61-69.
- [21]. Kumar G. Pavan, Babu Ch. Praveen, Samatha K., Jyosthna N., Showrilu K., International Letters of Chemistry, Physics and Astronomy 10(2014)25-37.
- [22]. Eyring H. and Kincaid J.F.,J.Chem.,6(1928)620.
- [23]. Masson D O; Philos. Mag. 8(1929)218.
- [24]. Syal V K, Chauhan S & Uma Kumari, Indian J. pure Appl. Phys., 43(2005)844.
- [25]. Kumar R, S. Jayakumar&Kannappan, Indian J. pure Appl. Phys., 46(2008)169-175.
- [26]. Syal V K, Patial B S, &Chauhan S; Acoustic Letter,29(2000)366.
- [27]. Muhuri, Parkash K, Bijan Das, Hazara, Dilip K; Indian J Chem,35A(1996)288.
- [28]. Subha M C S, Rao K C, &Rao S B, Indian J Chem ,25A(1986)424.
- [29]. Parmar M L, Khanna A & Gupta V K; Ind. J Chem,28A(1989)565.
- [30]. Syal V K,Chauahn A &Chauahn S; J Pure And Appl Ultrason,27(2005)61.
- [31]. Jone G & Dole M, J. Am. Chem. Soc.,51(1929)2950.
- [32]. Falkenhagen H & Dole M, Phys. Z.,30(1929)611; Falkenhagen H and Vernon E L, Phys. Z.33(1932)140.
- [33]. Jenkins H D B & Y Marcus; ChemRev.95(1995)2695.