

Weathering, Maturity And Tectonic Setting Of The Eocene Sediments Of Ogwashi-Asaba Formation

¹Ebiegberi Oborie ¹Felix D. Akori ²Archimedes U. Nwanne

¹Department Of Geology, Niger Delta University, Wilberforce Island, Bayelsa State.

²Nwanne Geoscan Ltd, Port Harcourt, Rivers State, Nigeria

Abstract:

Nine samples of fine grained clastic rocks which includes shales, fine sandstones, lignites siltstons and white clays were obtained from the outcropping unit of the Ogwashi-Asaba Formation in order to determine weathering, maturity and tectonic setting of the sediments. The results of the X-Ray Fluorescence analysis (XRF) showed enrichment of major oxides like SiO₂, Al₂O₃ and TiO₂, while Na₂O were depleted. The chemical index of Alteration (CIA), chemical index of weathering (CIW) and the plagioclase Index of Alteration (PA) showed averages of 91.84 (with a range of 77.8 – 98.46), 97.73 (94.40-99.56) and 96.83 (91.21-99.8) respectively. These values suggests a high intensity of source rock weathering. SiO₂/Al₂O₃, K₂O/Na₂O and the Index of composition variability (ICV) has averages of 3.8 (with a range of 1.8-8.42), 21-6 (with a range of 53.52-21.66) and 0.472 (1.04-0.863) respectively, indicating maturity variations from moderate to dominantly low and interpreted as the deposition of the sediments from multiple sources. A plot of SiO₂ vs Al₂O₃ + Na₂O + K₂O indicates a semi-humid to humid paleoclimatic conditions of deposition, while the tectonic discrimination plot suggests a passive margin setting.

Keywords: Outcropping, Formation, Weathering, Maturity, tectonic setting, source rock

Date of Submission: 06-04-2024

Date of Acceptance: 16-04-2024

I. Introduction

Numerous sedimentary basins in Nigeria were formed as a result of the South Atlantic Ocean opening up during the Cretaceous period, 65 million years ago, when Gondwanaland broke apart, causing the North American and African plates to separate. About 75,000 km² make up the Niger Delta Basin, and 12,000 km is made up of clastic fill [1]. The marine shales of the Akata Formation, the paralic deposits of the Agbada Formation, and the continental facies of the Benin Formation are the three diachronous units that make up the clastic fill of the basin. These units were all deposited during the Tertiary geological era. The same formations have been dubbed Imo-Shale (Akata), Ameki-Bende (Agbada), and Ogwashi-Asaba (upper Agbada facies) along the Northern axis of the Niger Delta where proximal portions of these lithostratigraphic units are visible. The Ogwashi-Asaba Formation's well-exposed rock facies are located in Onitsha, the Ogwashi-Uku region of Asaba, and Umuahia. According to [2] and [3], the formation is characterized by medium coarse planar cross-bedded ferruginized sandstones, white clays and claystones, silty muds, seams of black lignites, grey shales, and laterites.

II. Location And Accessibility

The outcropping Ogwashi-Asaba formation under investigation is located at the Ogbunike area of Onitsha toll gate, along the Enugu-Awka Express way and bounded by the coordinates 6°10'55.8"N and 06°51'55"E. It is a laterally and vertically extensive quarry which exposed a wide variation of interbedded rock facies. The facies comprise lignities, siltstones, shales, white clay (kaolin clays), coarse, medium and fine planar cross bedded sandstones, silts and siltstones, claystones and laterites.




Figure 1: The study area

III. Materials And Methods

Both field and laboratory methods were employed in the study. A one day field work was undertaken to the outcrop in order to determine bed thicknesses, grain sizes, strike-and dip and recorded in the course of the study. Both lateral and vertical sampling was done. Three different sections of the quarry was measured, sampled and labelled 1, 2 and 3. With the aid of the geological hammer and chisel, three samples were obtained from each of the measure sections of the quarry, making a total of nine samples of different lithologies including lignite, fine sands, siltstones, white clays (Kaolin) and shales. The nine samples were stored and labelled in polythene bags and subsequently sent to the research laboratory of the Geological Survey Agency, Kaduna for X-Ray Fluorescence Analysis (XRF).

Using an arget pulverizing machine, the samples were ground up. It was made sure that the ground samples could pass through 150 micromesh sieves. 5g of the pulverized sample and 1g of binding aid (starch soluble) were weighed into a beaker to check homogeneity. After the mixture was well combined to ensure homogeneity, it was pressed under high pressure to create pellets, which were then packed and labeled for examination. Model "minipal 4" Energy Dispersive x-ray fluorescence (EDXRF) spectrometry was used for the study. After the pellets were carefully positioned in their designated measurement positions on the machine's sample changer, the following conditions were created as the machine was turned on: the nature of the samples to be analyzed as pressed power (pellets), and the elemental composition of the samples were determined. For main oxides, 14 KV of current was employed, and for trace elements and rare earth metals, 20 KV. Ag/Al-thin for trace elements and rare earth metals, and "Kapton" for significant oxides were the chosen filters. A specific periodic table that was employed for the elemental analysis served as a reference for the filter selection.

Every sample was measured for 100 seconds, with air serving as the medium the whole time. After calibrating the machine, % results were obtained. The depletion or enrichment of elemental oxides in the samples is discussed in terms of the Element Enrichment factor (EF), defined as:

$$EF = \frac{\left\{ \frac{X}{AL} \right\}_{sample}}{\left\{ \frac{X}{AL} \right\}_{UCC}} \dots\dots\dots (1)$$

Where X = element; AL = Aluminum in sample; XUCC = element concentration in the upper continental crust; ALUCC = Aluminium concentration in the upper continental crust.

IV. Results And Discussion

The enrichment or depletion of element is interpreted as follows: EF>1 is enriched; EF<1 is depleted, while EF= 1 implies no change in concentration. The enrichment or depletion of elemental concentration in clastic rocks is due to source rock composition, resistance to chemical weathering or diagenesis and distance of recycling during transportation [4]. The calculated enrichment factor in table 2 showed that the samples are enriched in SiO₂ which ranges from 2.16 – 6.6 with an average of 5.02, TiO₂ Ranges from 0.16 – 4.09, and averaged 2.57, and Al₂O₃ with a range of 0.83 – 1.86, averaging 1.2. Their enrichment may be due to their resistance to chemical destruction during weathering, grain size effect and slow decomposition [5]. The depleted element includes CaO which has an average EF of 0.09 (with range of 0.003 – 0.37), MgO averaged 0.03, K₂O averaged 0.59 Na₂O average 0.17, MgO averaged 0.11 and Fe₂O₃ is 0.7. Their depletion may also be due largely due to their source of composition and the ease of chemical destruction during weathering.

Table 1: Oxides’ enrichment factor in samples

Oxide	Lithology	1A	1B	1C	2A	2B	2C	3A	3B	3C	Average
SiO ₂	White clay	64.51	68.248	51.960	57.80	74.127	51.435	65.302	44.549	75.867	61.52
CaO	Fine sandstone	0.017	0.124	0.120	0.220	Nd	Nd	Nd	0.368	Nd	0.569
MgO	Lignite	0.050	0.021	0.020	0.032	Nd	0.129	Nd	0.176	Nd	0.146
K ₂ O	Siltstone	0.225	0.131	0.315	1.780	2.855	2.854	2.435	1.820	3.015	1.9
Na ₂ O	Siltstone	0.010	0.007	0.008	0.044	0.650	0.540	0.120	0.034	0.760	0.227
TiO ₂	Lignite	1.369	0.892	2.944	0.879	1.197	2.209	1.670	2.28	1.703	1.694
MnO	Shale	0.002	0.012	0.006	0.008	0.003	0.015	0.005	0.055	0.008	0.0123
Fe ₂ O ₃	Fine sandstone	1.519	11.617	2.166	5.545	1.310	6.866	3.119	12.177	2.278	4.002
Al ₂ O ₃		24.28	16.770	28.789	23.000	18.458	20.052	19.107	19.436	12.802	20.11
SiO ₂ /Al ₂ O ₃		2.660	4.070	1.80	2.51	4.016	2.57	8.42	2.29	5.926	3.82
K ₂ O/Na ₂ O		22.50	18.714	39.375	26.81	4.392	5.285	20.435	53.529	3.963	18.88
CIA		98.35	98.460	98.48	91.849	84.04	85.52	88.2	89.74	77.48	98.81
CIW		99.26	99.22	99.56	98.87	96.60	97.38	99.37	94.98	94.40	98.19
PIA		99.25	99.21	99.8	91.21	96.00	96.96	96.96	99.29	92.79	104.96
ICV		0.104	0.757	0.187	0.369	0.326	0.650	0.386	0.868	0.606	0.464

Table 2: Elemental enrichment factor in samples

Element	1A	1B	1C	2A	2B	2C	3A	3B	3C	AV EF	UCC
SiO ₂	2.65	0.94	6.6	3.34	4.26	5.217	1.79	4.49	3.17	5.02	66.60
MgO	0.12	0.08	0.004	0.001	Nd	0.04	Nd	0.06	Nd	0.03	2.480
K ₂₀	0.06	0.05	0.07	0.50	0.9	0.91	0.81	0.59	1.50	0.59	2.800
Na ₂₀	0.002	0.002	0.001	0.009	0.17	0.12	0.29	0.08	0.28	0.17	3.270
TiO ₂	1.7	0.16	3.20	1.14	3.3	3.3	2.7	3.6	4.09	257	0.500
Mno	0.012	0.11	0.32	0.05	0.025	0.15	0.20	0.04	0.09	0.11	0.100
Fe ₂₀₃	0.17	1.9	0.21	0.65	0.08	0.96	0.50	1.80	0.60	0.76	5.400
Al ₂₀₃	1.6	1.08	1.86	1.49	1.19	1.30	1.24	1.26	0.83	1.2	15.40

Weathering

The degree or intensity of chemical weathering at a source area is measured by application of various weathering indices. The chemical composition of clastic rocks is a function of their sources area composition and weathering [6]. Thus [6] proposed the Chemical Index of Alteration (CIA), defined as:

$$CIA = \left\{ \frac{Al_2O_3}{Al_2O_3 + CaO^* + Na_2O + K_2O} \right\} \times 100 \dots\dots\dots (2)$$

CaO* is the amount of CaO incorporated in the silica fraction of the samples [7]. The CIA measures the ratio of primary minerals and secondary products such as clay minerals, which result from the destruction of plagioclase, hence Na, Ca, and K are progressively depleted from source rocks during chemical weathering [8]. The chemical index of weathering (CIW) proposed by [9] is also used for the evaluation of intensity of weathering. It is similar to the CIA, except the removal of K₂O in the equation.

The CIW proposed by Harnois is expressed as

$$CIW = \left\{ \frac{Al_2O_3}{Al_2O_3 + CaO^* + Na_2O} \right\} \times 100 \dots\dots\dots (3)$$

According to [10] and [11], the CIA and CIW are interpreted similarly, with values of 50 for unweathered top continental crust and around 100 for extensively weathered material with full elimination of alkali and alkaline earth element. The CIA values of the sample ranges from 77.8 – 98.46 with an average of 91.84. while the CIW ranges from 94.40 – 99.56 with an average of 97.73. These values are indication of a high intensity of weathering at the source area. This is also demonstrated in the bivariate plot of Al₂O₃ versus CIA [12] showing a clustering of the data points in the segments of high weathering intensity (figure 2). The plagioclase Index of Alteration (PIA) by [7] is another weathering indices defined as:

$$PIA = \left\{ \frac{Al_2O_3 + K_2O}{Al_2O_3 + CaO^* + Na_2O + K_2O} \right\} \times 100 \dots\dots\dots (4)$$

Where CaO* is the CaO content in the silicate fraction. The CIA determines the progressive breakdown of feldspars to clay minerals. It is also interpreted like the CIA and CIW where 0 – 50 indicates unweathered plagioclase, while 50 – 100 is an indication of moderate to high intensity of the chemical destruction of plagioclase. The samples of The Ogwashi Asaba Formation displayed PIA values that are quite similar to those of CIA and CIW with value ranges 91.21 – 99.8, averaging 96.83. This also suggest an intense weathering of source rocks.




Figure 2: Al₂O₃ vs CIA Plot showing intensity of weathering in the samples [After 12]

Maturity

The SiO₂/Al₂O ratio is very sensitive to weathering, recycling and diagenetic alteration processes, hence can be used as an index of sediment maturity. It is a measure of the quartz content of siliciclastic rocks. Consequently, quartz outlives feldspar, mafic minerals, and lithics as sediment maturity increases [13]. Those that are chemically immature are represented by low ratios, while those that are mineralogically mature (quartzose, rounded) exhibit high ratios.

Finer grain sizes may indicate maturity, however the true indicator of immaturity is more closely linked to the amount of clay present. In unmodified igneous rocks, the average SiO/Al₂O₃ ratios vary from 3.0 in basic rocks to 5.0 in siliceous rocks. Values of SiO₂/Al₂O₃ ratios of the sample under investigation ranged between 1.80 – 8.42 with an average of 3.81, which suggest immaturity (low maturity) of the sediment. Also the K₂O/Na₂O ratio is also used as a maturity index which measures the alkali concentration in sediments. Abundant feldspar represented by Na₂O and K₂O is a reflection of immaturity in sandstones and shales. The K₂O/Na₂O ratio in the sample has value range of 3.96 – 53.52 averaging 21.66, which suggest a variation between moderate and low maturity. This suggest that the sediments were derived from different sources and depositional settings. A plot of SiO₂ vs Al₂O₃ + K₂O/Na₂O (figure 3) [14] showed that the data points are plotted in the Humid – semi humid climatic axis. Humid climate promote the intensity of chemical weathering, leading to the progressive maturity of clastic rocks. It could therefore be inferred that the humid conditions may have accelerated the intensity of weathering and maturity of the samples. The Index of Compositional variability (ICV) is another important geochemical proxy used for the maturity of clastic rocks. The ICV proposed by [15] is expressed as;

$$ICV = \left\{ \frac{Fe_2O_3 + CaO + K_2O + Na_2O + MgO + MnO + TiO_2}{Al_2O_3} \right\} \dots\dots\dots (5)$$

According to [15], ICV > 1 indicates compositionally immature, derived from a cratonic basin, while ICV < 1 are compositionally mature and deposited in a tectonically quiescent cratonic environment. The samples under study has ICV values that ranges from 0.104 – 0. 868 with an average of 0.472, which is an indication of matured sediments that were recycled by long distance transport and deposited in a tectonically stable basin.




Figure 3: Climatic condition and maturity of the samples, [After 14]

Tectonic Setting

The tectonic origin (setting) of siliclastic rocks may also be inferred from major elements concentrations in sediments. In order to ascertain the tectonic setting of clastic sedimentary rocks, [13] constructed a tectonic decimation diagram (figure 4) using the ratio K₂O/Na₂O versus SiO₂.

Sediments deposited in the passive margin (PM), active continental margin (ACM), and oceanic island arc (OIA) may be distinguished from one another using a bivariate graphic. The plot suggest that the samples of the Ogwashi Asaba Formation were deposited in a passive margin tectonic setting (figure 4).




Figure 4: Tectonic Discrimination Diagram of the samples [After [13]]

V. Conclusion

In order to study the intensity of weathering, maturity and tectonic setting of the outcropping Eocene Sediments of the Ogwashi Asaba Formation, nine samples of mixed lithologies including lignites, fine sandstones, shales and white clays were retrieved from three sections of the Ogbunike quarry (Onitsha) during a one day field work. The samples were analyzed using the x-ray fluorescence (XRF) technique to determine major elemental concentrations in the fine grained sediments. From the results obtained, major oxides like SiO₂, Al₂O₃ and TiO are enriched. This was attributed to their stable and immobile characteristics during weathering and recycling. Conversely, mobile alkali oxides like Na₂O, CaO, K₂O, Fe₂O₃ and MnO were depleted due to their susceptibility to chemical destruction during weathering and diagenesis, sources area composition and grain size effect. Average values of CIA, CIW and PIA includes 90.23, 97.73 and 96.83 respectively which indicates an intense weathering at the source areas. SiO₂/Al₂O₃, K₂O/Na₂O and ICV maturity indices has average values of 3.81, 21.66 and 0.472 respectively. These figures indicates a maturity variation from moderate to low maturity. These variations possibly suggests a deposition of the sediments from different sources. The bivariate tectonic discrimination plot placed the sediments in the passive margin field.

References

- [1] Nwajide, C.S. and Reijers, T.J.A. (1996).. geology of the Southern Anambra Basin. In Reijers, T.J.A. (Ed), Selected chapters in Geology; sedimentary geology and sequence stratigraphy in Nigeria and three case studies and a field guide. SPDC corporate reprographic services, warri, Nigeria. Pp. 131-148.
- [2] Whiteman, A.J. (1982). Nigeria: Its petroleum geology, reserves, and potential Graham and Trotman, London, Vol 2. 394pp.
- [3] Akpofure, E. and Akori, F. (2022). Major element analysis of Ezeaku Sandstones outcropping at Abini, cross Rivers State, Nigeria. geosciences and remote sensing journal, clausius scientific press Canada. Vol. 5 No. ISSN 2523-2592.
- [4] Nesbih, H.W. and Young, G.M. (1982). Early proterozoic climates and plate motions inferred from major element chemistry of lutites. Nature, 299. 715-717.

- [5] Madukwe, H.Y., Ayodele, S.O., Akinyemi, S.A., and Adebayo, O.F. (2016). Classification, maturity, provenance, tectonic setting and source area weathering of Ipole and Erin Ijesha stream sediments, South West Nigeria. *put jurn. Of Adv. Sci and tech. res.* Vol. 2249-9954.
- [6] Nesbitt, H.W., Young, G.M. (1989). Determination of tectonic setting of sandstone-mudstone suites using SiO_2 content and $\text{K}_2\text{O}/\text{Nk}_2\text{O}$ ratio. *J. Geol.*, 94. P. 635-650.
- [7] Fedo, C.M., Nesbitt, H.W., and Young, G.M. (1995). Unraveling the effect of potassium metal somatism in sedimentary rocks sand paleosols, with implication for paleoweathering conditions and provenance *J. of Geology*, 23, 921-924.
- [8] Baivegunhi, C., Liu, K., Gwayaya, O. (2017). Geochemistry of sandstone sand shales from the ECCA group, Karoo super group, in Eastern Cape province of south Africa implications for provenance, weathering and tectonic setting. *Aegruyter open geosciences journal.* Vol. 9. P. 340-360.
- [9] Harnois, L., (1988). Geochemical classification of terrigenous and shales from core or log data. *Journal of sedimentary petrology.* 58(5); p/ 820-829.
- [10] Mchennan, S.M., Hemming, S., McDaniel, D.K., Hanson, G.N. (1993). Geochemical approaches to sedimentation, provenance and tectonics. *Geol. Soc. Am/ Spec. pap.* 284, 21-40.
- [11] Mongelli, G., Cullers, R.L and Muelheisen, S. (1996). Geochemistry of late cretaceous Oligocene shales from the varicowri formation, Southern apennines, Italy: implications for mineralogical, grain-size control and provenance. *Eur. J. Mineral.* 8: p. 773-754.
- [12] Madukwe, H.Y., Akinmosun, A, Akinyemi, S.A., Aturamu, A.O and Ojo, A. (2014). Provenance, Tectonic setting and maturity of the Ishara sandstone, South Western Nigeria. *insight into major elemental geochemistry put journal of current research.* Vol.6. pp. 11123-1113.
- [13] Roser, B.P. and Korsch, R.J. (1983). Provenance signature of sandstone-mudstone suite determined using discernment function analysis of major element data. *Chem.. geol.* 67: p. 119-139.
- [14] Suttner, L.J., Dutta, P.L., (1986). Alluvial Sandstone composition and paleo climate L. Framework mineralogy. *Journal of sedimentary petrology.* Vol. 56. P. 329-345.
- [15] Cox, R., Lowe, D.R. and Cullers. R.C. (1995). The influence of sediment recycling and basement composition of evolution of mudrocks chemistry in the South Western united state, geochemical cosmochimica Acta. Vol. 59; pp 2919-2940.