

Deriving the Useful Expression for Time Dilation in the Presence Of the Gravitation by means of a Light Clock

Dr. Hasabalrasoul Gesm Allah Ismail Hamza¹, Mr .Wasil Abdalgader Abdalla Alhassan²

¹ Department of Physics & Mathematics, faculty of Education Hasaheisa /Gezira University - Wad Madani , Sudan .

² Department of Physics ,Tahseen Aladaa Training Center /Salman bin Abdulaziz University ,Saudi Arabia.

Abstract: Generalized special relativity is one of the most promising models that is found to cure many defects of special relativity. These defects include un capability of explaining gravitational red shift and satisfaction of the Newtonian limit.

Despite these successes generalized special relativity suffers from noticeable setbacks. First of all its way of derivation make it restricted to weak fields only. Although recent derivation make it applicable to fields other than the weak and gravitational field, but this derivation needs to be strengthened.

In this study a light clock is used to derive a useful expression for time in General special relativity. These expressions are typical to that derived before by using curved space. Time, but they are not restricted gravitational and weak fields. The expressions for time hold for all fields.

Keywords: alight clock mirror, Generalized, gravitational, time Dilation.

I. Introduction

Einstein theory of special relativity (GR) is one of the biggest achievements in physics. It changes radically the classical concept of space and time. The GR theory is based on two postulates. One of them is concerned with the homogeneity of space and time the other is concerned with the constancy of speed of light[1]

The theory of GR succeeds in explaining a large number of experimental observations, like meson decay, pair production and nuclear mass defect [2].

Unfortunately GR suffers from noticeable setbacks. First of all, it cannot explain the change of neutrino mass, it does not satisfy correspondence principle, since its expression of relativistic energy does not satisfy Newtonian limit since it does not consist of a term representing the potential energy.

It cannot explain also the gravitational red shift which indicates that the photon mass as well as its periodic time change with the gravitational field[3]

II. Set backs of the Theory of Special Relativity

In GR theory the expression of time t , mass m ,space l and energy E .

$$t = \frac{t_0}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (1)$$

$$l = l_0 \sqrt{1 - \frac{v^2}{c^2}} \quad (2)$$

$$m = \frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (3)$$

$$E = mc^2 \quad (4)$$

Where t_0, l_0, m_0 stands for time, length, and mass in arrest frame While $t, l, ,$ and m represents time, length and mass in frame moving with velocity v with respect to the rest frame.

The expression of relativistic mass in the Newtonian limit , where the light speed is law is by[4,5,6,7].

$$E = m_0 c^2 \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}} \approx m_0 c^2 \left(1 + \frac{1}{2} \frac{v^2}{c^2}\right)$$

$$E \approx m_0 c^2 + \frac{1}{2} m_0 v^2 \quad (5)$$

This is not typical to the Newton expression of energy[8].

$$E = \frac{1}{2} m_0 v^2 + V = T + V \quad (6)$$

Since it does not consists of on expression representing the potential energy.

The gravitational red shift phenomenon is also related to GR. It is concerned with the change of photon frequency from f to f' when it enters a gravitational field of potential V . According to the principle of Newtonian energy conservation

$$hf' = hf + V \quad (7)$$

Since the photon energy is also given by (4) it follows that

$$hf' = m' c^2$$

$$hf = mc^2 \quad (8)$$

Thus

$$m' c^2 = mc^2 + V$$

$$\Delta m = m' - m = \frac{V}{c^2} \quad (9)$$

Equation (9) indicates that the photon changes with the potential which is in direct conflict with the equation (3) , which states that the mass does not change with the potential V [9].

III. General Expression for Kinetic and Potential Energy

$$m \frac{dv}{dt} = F = -\nabla V = -\frac{\partial V}{\partial x} \quad (10)$$

For m independent of x and ϕ dependent on x only [9].

When

$$V = m\phi$$

$$m \frac{dv}{dx} \frac{dx}{dt} = -m \frac{d\phi}{dx} \tag{11}$$

$$m \int_{v_0}^v v dv = -m \int_0^\phi d\phi \tag{12}$$

$$\frac{1}{2} m [v^2 - v_0^2] = -m\phi \tag{13}$$

$$v^2 = v_0^2 - 2\phi \tag{14}$$

This can be also be dived from

$$v^2 = v_0 - 2ay \tag{15}$$

$$V = \text{potential} = Fy \tag{16}$$


$$may = m\phi$$

$$\phi = ay$$

$$v^2 = v_0 - 2\phi \tag{17}$$

IV. Derivation of Time Dilation in the presence of Fields

Consider an attractive field like gravitational field. Let a light mirror of length l_0 move upward with initial speed v_0 .



at

Figure (1) a clock in free space

When clock moving up its speed be comes v at a time t .

For the clock in free space the time to is given by

$$\begin{aligned} t_0 &= l_0 / c \\ l_0 &= ct_0 \end{aligned} \tag{18}$$




Figure (2) For a clock moving under the action of the field

$$\begin{aligned} y &= vt \\ x &= l_0 \end{aligned} \quad (19)$$

The velocity v_f and the vertical displacement y can be expressed in terms of the initial velocity v_o and acceleration a as follows

$$\begin{aligned} v_f &= v_o - at \\ y &= v_o t - \frac{1}{2} at^2 = (v_o - \frac{1}{2} at)t \end{aligned}$$

Since the average velocity v is given by

$$\begin{aligned} v &= \frac{v_o + v_f}{2} \\ &= \frac{2v_o - at}{2} \\ &= v_o - \frac{1}{2} at \end{aligned}$$




Figure (3) light moving with speed c from p_1 to p_2

Assuming the light moving with speed c from p_1 to p_2 in a trajectory inclined to the vertical plane it follows that

$$c^2 t^2 = l_0^2 + y^2 = c^2 t_0^2 + y^2$$

$$c^2 t^2 - y^2 = c^2 t_0^2 \quad (20)$$

But

$$y = vt$$

$$y^2 = v^2 t^2$$

$$c^2 \left[1 - v^2/c^2 \right] t^2 = c^2 t_0^2 \quad (21)$$

Thus the time in the presence of the field is given by

$$t = \frac{t_0}{\sqrt{1 - v^2/c^2}} = \frac{t_0}{\sqrt{1 + \frac{2\phi}{c^2} - \frac{v^2}{c^2}}} \quad (22)$$

This expression holds for any field.

V. Discussion

A useful expression for time dilation were found in equations (22). These expressions were derived by considering a light mirror clock moving in a field. The motion of this clock is obeys the ordinary equations of motion in any arbitrary field. Fortunately the expressions is found to be typical to that obtained in GSR .

VI. Conclusion

This model which uses a light clock to derive GSR, shows that the GSR can hold for any field, gravitational or non-gravitational.

References

- [1]. O.D Jefimenko , "Electromagnetic Retardation , and theory and Relativity "2nd Edition , Electret Scientific, ISBN 0917406214 9780917406218, 1997.
- [2]. F.L. Derek "an Introduction to Tensor Calculus and Relativity "3rd Edition Jon Wily & Sons Ltd , New York 1982, ISBN 0.486-42540-1(plk).
- [3]. M . H . M .Hilo , et al , "Using of Generalized Special Relativity .in Estimating the Proton (Nuclear) Mass to Explain the Mass Defect ,"Natural science Vo l .3, No.2, 141-144 (2011).
- [4]. R.F. Evens and J. Dunning – Davies "The Gravitational Red-Shift "University of Hull , England , arXiv:gr-qc/0403082V12004.p.3.
- [5]. Röhrlich, F. An elementary derivation of $E=mc^2$.Am. J. Phys. 58: 348–349. 1990.
- [6]. Thomas, E.G. What’s so special about $E=mc^2$. Eur. J. Phys. 26: 125- 130 , 2005.
- [7]. Sharma, A. The Origin of the Generalized Mass-Energy Equation $E=mc^2$ and Its Applications in General Physics and Cosmology. Physics Essays . 17: 195-222 . 2004.
- [8]. R.C. Tolman , "Relativity, Thermodynamic- and Cosmology 1s Edition Oxford university press, oxford 1934. LCCN 340-32023. Reissued (1987) New York: Dover ISBN 0-486-65383-8..p.591.
- [9]. H. I. Hamza ,Mubarak .Dirar. Abdalla. Speed of Light and Momentum within the frame work of Generalized Special Relativity, University of Africa Journal of Sciences ,volume No 3 – 2013.