

Trend of Cervical Dysplasia And Cervical Cancer -in Papanicolaou Smears in Perimenopausal Females In Tertiary Care Centre, Gwalior Madhya Pradesh

¹Dr Shelly Gupta, ²Dr Sudha Iyengar, ³Dr Bharat Jain

¹(Resident, Department Of Pathology) ²(Associate Proff Department Of Pathology) ³(Hod Department Of Pathology)

¹²³Department Of Pathologygajra Raja Medical College Gwalior, Madhya PradeshAddress:Senior Girls Hostel Grmc ,Gwalior

*Corresponding author: 1Dr Shelly Gupta

Abstract:

Introduction Cervical Dysplasia & cervical Cancer remain important health problem. Cervical cytology by Papanicolaou (Pap) smear is an effective means of screening for cervical premalignant & malignant conditions.

Aim :The aim of this study was to assess the trend of cervical dysplasia & cervical cancers in perimenopausal women in tertiary care centre of Madhya Pradesh and to assess the association (if any) of Neoplastic/dysplastic changes.

Material & Methodsa total of 3,610 cases were enrolled in 4 year study cervical scrape smears were collected & stained using Rapid-Pap method. The emphasis was on epithelial abnormalities classified in accordance with the Bethesda System. The cases belonged to the perimenopausal age group (41-50 years).

results :Trend of Cervical Dysplasia - 20.5% (2013), 11% (2014), 10.5% (2015), 8% (2016)

Trend of SCC -2.5% (2013), 1.8% (2014), 1% (2015), 0.5% (2016).

Conclusion :Study has shown a relatively decreasing trend of cervical dysplasia and cervical cancer in cervical smears which are taken regularly as a part of ongoing family planning programme in our tertiary care set up.

Date of Submission: 01-12-2017

Date of acceptance: 09-12-2017

I. Introduction

Cervical Intraepithelial neoplasia, Dysplasia & cervical Cancer remain important health problem. Cervical cytology by Papanicolaou (Pap) smear is an effective means of screening for cervical premalignant & malignant conditions. Carcinoma cervix worldwide accounts for 15% of all cancers diagnosed in women.[1] Cervical cancer is one of the leading cancer in women with the estimated 5.0 lakhs new cases every year of which 80% occur in developing countries.[2] In India, it is estimated that the number of new cases are over 1,40,000 .[3] Cancer cervix occupies the top rank or second among cancers in women in developing countries ,where as in the affluent countries cancer cervix does not find a place even in top five leading cancers in women. 70% or more of these cancers are in stage 3 or higher at the time of diagnosis. The role of Papanicolaou (Pap) smear as a cancer screening tool for cervix has been substantiated by several studies in the last 50 years and the methods has resulted in falling incidence and mortality of cervical cancer in developed world. Therefore, the data on the prevalence of cervical dysplasia and cancer in various population in this country is not known. There is an urgent need for initiation of community screening and educational programs for the control and prevention of cervical cancer in India.[4]Therefore, objective of this study was to assess the trend of cervical dysplasia, & cervical cancers in perimenopausal women in tertiary care centre of Madhya Pradesh .

II. Material & Methods

A total of 3,610 cases were enrolled in 4 year study cervical scrape smears were collected & stained using Rapid-Pap method. And classified in accordance with the Bethesda System.[5] The cases belonged to the perimenopausal age group (41-50 years).[6]It was ensured that no local douche, antiseptic cream and no local internal examination was done on the day of rest. The patient was placed in dorsal lithotomy position and ausco's bivalve speculum was introduced through vagina and cervix was visualized. The longer projection of the Ayre's spatula was placed in the cervix near squamo-columnar junction and rotated through 360°. The cellular material thus obtained was quickly, but gently smeared on a clean glass slide. The glass slide was then immediately put into the coplin jar containing 95% ethyl alcohol which acted as a fixative.


The prepared smears were then stained according to Papanicolaou's technique. The cytological interpretation of the smears was made according to the 2001 Bethesda system.

III. Results


Total no of 3,610 cervical scrape smears taken & examined . results are divided in – Cervical dysplasia and Squamous cell carcinoma.

Table: Categorization Of Cytodiagnosis:


S.N.	Category	2013	2014	2015	2016
1	Total Number Of Smears Collected	681	854	919	1156
2	Cervical Dysplasia	9.0%	8.0%	6.0%	5.0%
3	ScC	1.8%	1.5%	0.8%	0.5%



LSIL: Smear showing superficial and intermediate cells with koilocytic changes as perinuclear halos indicating HPV infection with mild dysplasia



Smear showing cells with scant cytoplasm and marked hyperchromatic nucleus and irregular nuclear contour.



SCC:Smear showing macro-nucleolus with tumor diathesis Tadpole and fiber cells are present: feature of keratinizing type SCC

IV. Discussion

Uterine cervix is ideal for screening due to easy accessibility of cervix for inspection, palpation, exfoliative cytology and screening .cancer cervix is a preventable and curable disease due to effective screening methods available and a long preinvasive phase of the disease and various treatment modalities available .Cervical cancer screening represents one of the great success stories in cancer prevention. Thus screening of perimenopausal women should be made mandatory as it helps in picking of cases of cervical dysplasia which can be treated easily with different treatment modalities and progression of cervical precursors to invasive cancer can be reduced, thus reducing the incidence of cervical cancer cervix. It has been established that low grade dysplasia is usually asymptomatic. The common complaints in high grade dysplasia are postmenopausal bleeding and intermenstrual and postcoital bleeding per vaginum.[7] In our study also, the most common complaints in cervical dysplasia was postcoital bleeding & postmenopausal bleeding per vaginum and on per speculum examination, most common finding was cervical polyp. In the past also, visual inspection of the cervix had been stressed by Sankarnarayanan et al.[8] Thus screening of perimenopausal women should be mandatory as it helps in picking of cases of cervical dysplasia which can be treated easily with different treatment modalities and progression of cervical precursors to invasive cancer can be reduced , thus reducing the incidence of cancer cervix.

V. Conclusion

Study has shown a relatively decreasing trend of cervical dysplasia and cancer in cervical smears which are taken regularly as a part of ongoing family planning programme in our tertiary care set up. Pap smear is good tool to find abnormal cervical cytology-cervical dysplasia & cervical cancers in population. Periodic cytological screening would go a long way in the early detection of cervical dysplasia and help in reducing the incidence of cervical cancer.

Refrence

- [1]. Boyle P,Ferlay J. Cancer incidence and mortality in Europe,2004. *Ann Oncol* 2005;16:481-8
- [2]. Tristen C ,Bergstrom S. Cancer in developing countries. A threat to reproductive health. *Lakartidningen* 1996;93:3374-6
- [3]. Juneja A,Sehgal A, Sharma S, Pandey A. Cervical cancer screening in India: Strategies revisited . *Indian J Med Sci* 2007;61:34-47
- [4]. Murty NS, Chaudhary K ,Saxena S. Trends in cervical cancer incidence – Indian Scenario. *Eur J Cancer Prev* 2005;14:513-8
- [5]. Solomon D, Davey D,Kurman R, Moriarty A,O'Connor D, Prey M,et al. Forum Group Members ,Bethesda2001 Workshop. The 2001 Bethesda system: Terminology for reporting results of cervical cytology. *JAMA* 2002;287:2114-9
- [6]. William's Gynaecology ,2 edition, Chapter 21, Menopausal Transition
- [7]. Chiitale AR, Bhuvaneshwari, Krishna UR. Histological assessment of cytologically abnormal smears. *Indian J Cancer* 1976;13:324-9
- [8]. Sankarnarayan R, Budukh AM, Rajkumar R. Effective screening programs for cervical cancer in low and middle –income developing countries. *Bull World Health Organization* 2001;79:954-62

- [9]. Mulay K, Swain M, Patra S, Gowrishankar S. A comparative study of cervical smears in an urban hospital in India and a population based screening program in Mauritius. *Indian J Pathology Microbiology* 2009;52:34-7
- [10]. Patel TS, Bhullar C, Bansal R, Patel SM. Interpreting epithelial cell abnormalities detecting during cervical smear screening- A cytologic approach. *Eur J Gynaecology Oncology* 2004;25:725-8