

A Comparative Study of Treatment of Pterygium by Excision and Limbal Autograft or Augmented with Post-Op Mitomycin C

Dr. Anurag Majumdar, Dr. Abhijit Ray, Dr. Shivam Gupta
Dept. of Ophthalmology, Agartala Govt. Medical College, Agartala, Tripura (West)
Corresponding Author: Dr. Abhijit Ray

Abstract: Pterygium is a cause of diminution of vision when it encroaches on the pupillary area. It typically develops in patients who have been living in hot climates and may represent a response to chronic dryness and exposure to ultraviolet light, dust and hot winds. It is rarely seen in patients younger than the age of 20 years. The study was conducted to study the clinical profile of pterygium and to compare results of recurrence by excision with limbal conjunctival autografts or postoperative topical Mitomycin C drops.

The present study was conducted at the Department Of Ophthalmology at Agartala Government Medical college and GB Pant Hospital over a period of 18 months (October 2016 to May 2018).

One of the main aims of the study was to compare the effectiveness of conjunctival limbal autografting and mitomycin C in preventing the recurrence of pterygium. In the present study it was found that recurrence of pterygium occurred in 2 (5%) out of 40 eyes treated with limbal conjunctival autograft and 3 (7.5%) out of 40 eyes in whom pterygium excision was followed by postoperative mitomycin C.

Keywords: Pterygium, Mitomycin-C, Recurrence, Limbal autograft, Vascularisation

Date of Submission: 14-11-2018

Date of acceptance: 29-11-2018

I. Introduction

Pterygium is a cause of diminution of vision when it encroaches on the pupillary area. The patient also has cosmetic problems. Epidemiological surveys indicate that the prevalence rates of pterygium vary depending on the exact population [1-6]. Overall prevalence rates range from 0.7 to 31% in various populations of the world [1-6]. As a general rule, prevalence rates for a pterygium increase with age although a decline in prevalence rates has been reported for patients over 60 to 70 years old [1,3]. It typically develops in patients who have been living in hot climates and may represent a response to chronic dryness and exposure to ultraviolet light, dust and hot winds. It is rarely seen in patients younger than the age of 20 years. With the development of new methods of treatment, frequency and severity of pterygium have declined. The study was conducted to study the clinical profile of pterygium and to compare results of recurrence by excision with limbal conjunctival autografts or postoperative topical Mitomycin C drops.

II. Aim And Objectives

To prospectively analyze a clinical study of pterygium and results of treatment by excision and limbal autograft surgery or augmented with post operative topical mitomycin C.

III. Material And Methods

Present study was a prospective, comparative case study involving 80 eyes of 40 patients who attended the Department of Ophthalmology at Agartala Government Medical College and GB Pant Hospital. The duration of the study was 2 years. The study was reviewed by the institutional review board. Each eligible patient provided informed consent prior to the enrollment in the study.

Inclusion Criteria

All cases of pterygium attending the OPD including recurrent pterygium.

Exclusion Criteria

Patients coming from far away places who were not able to come for regular follow up were excluded from this study

A detailed history was taken and recorded regarding the disease which included age, occupation, residence, exposure to dust and hot wind. The extent of corneal involvement by the pterygium was also noted.

Patient were assigned to two groups (A and B) randomly. Group A underwent limbal conjunctival autograft. Group B underwent pterygium excision augmented with postoperative topical mitomycin C application (**Figure**

1 and 2). In patients belonging to Group A excision of pterygium was done followed by limbal conjunctival autograft taken from upper temporal quadrant and sutured on the bare sclera using 6-0 vicryl sutures (**Figure 3** and **4**). Patients were given gatifloxacin or moxifloxacin with dexamethasone eye drop 4 times a day for 2 weeks, then 3 times a day for 2 weeks, and then 2 times a day for 2 weeks. Gatifloxacin or Moxifloxacin with dexamethasone ointment was applied at night for 6 weeks. The patients were followed after one week and after that monthly for a year. Patients belonging to group B were treated with mitomycin C in a strength of 0.01 mg/ml in carboxy methylcellulose. On first and second postoperative day patients were told to use either gatifloxacin or moxifloxacin eye drops 6 times a day. From third to twenty first postoperative day patients were given mitomycin C drops 4 times a day and gatifloxacin (3 mg) with dexamethasone (1 mg) eyedrops 4 times a day.

Patients were followed weekly for three weeks and then monthly for a period of one year. Best corrected visual acuity were noted on every visit. Patients were asked for any pain, photophobia, redness, lacrimation.

Slit lamp examination was done and eye was looked for:

- 1) Any recurrence.
- 2) Scleral thinning.
- 3) Corneal vascularisation.

IV. Results

The present study was conducted at the Department Of Ophthalmology at Agartala Government Medical college and GB Pant Hospital over a period of 18 months (October 2016 to May 2018).

80 eyes of 40 patients were enrolled in this study. Patients were assigned to two groups (A and B) randomly. Group A underwent limbal conjunctival autograft, 40 patients. Group B underwent pterygium excision augmented with postoperative topical mitomycin C application included 40 patients.

Age Distribution

Out of the 80 patients maximum number of patients were in the age group from 41 to 50 years in which the youngest patient was 23 years of age and the oldest patient was 70 years of age.

Sex Distribution

Out of the 80 patients there was a male preponderance in this study with 47 males (58.75%) as compared to 33 females (41.25%). The higher incidence in males could be attributed to their greater exposure to hot, dry and dusty climate.

Eye Involved

Out of 91 eyes in 80 patients right eye was involved in 54 patients (67.5%), left eye was involved in 37 patients (46.25%).

Site of Pterygium

The site of pterygium is found to be nasal in 88 eyes (96.70%) of eyes, while temporal in 3 (3.30%) of eyes.

Size of Pterygium

The cornea was encroached by pterygium between 2 - 3 mm in 88 eyes (96.70%) while the cornea was involved greater than 3 mm by pterygium in 3 eyes (3.30%).

Recurrence Rates

One of the main aims of the study was to compare the effectiveness of conjunctival limbal autografting and mitomycin C in preventing the recurrence of pterygium. In the present study it was found that recurrence of pterygium occurred in 2 (5%) out of 40 eyes treated with limbal conjunctival autograft and 3 (7.5%) out of 40 eyes in whom pterygium excision was followed by postoperative mitomycin C.

Recurrence Time after Surgery

Chen *et al.* [17] reported the mean time of recurrence varied from 3.7 to 4.8 months with only 6% of recurrences occurring after the sixth postoperative month.

In the present study, cases where limbal conjunctival autografting was done there was recurrence occurring in the first and third month whereas in cases where pterygium excision followed by mitomycin was done recurrence occurred in third, fourth and sixth month of follow up (**Table 4**). So the present study agrees with previous studies.

Age of Patients with Recurrence

In the study conducted by Figueiredo *et al.* [18] patients with age less than 50 years had significantly more recurrences. It has been suggested that lipid degeneration in the cornea is an inhibiting factor to pterygium growth, based on observations that pterygium does not cross an arcus senilis to any great extent. The presence of increasing amounts of lipid degeneration with age might explain in part, the strong association between age and recurrence.

The present study (**Table 5**) shows that there were two recurrences in eyes with pterygium excision followed by conjunctival autograft with age less than 50 years of age whereas in eyes with pterygium excision followed by mitomycin C there was one recurrence in patients less than 50 years and two recurrences in patients greater than 50 years of age which agrees with previous studies.

Table 1. Shows the site of pterygium involved.

Site	No. of eyes	Percentage
Nasal	88	96.70%
Temporal	3	3.30%
Total	91	100%

Table 1. Shows the site of pterygium involved.

Size	No of eyes	Percentage
2 - 3 mm	88	96.70
Greater than 3 mm	3	3.30

Table 3. Shows the recurrence rates.

Pterygium autograft		Pterygium excision with mitomycinC
No of eyes treated	40	40
Recurrence no. %	2 (5%)	3 (7.5%)

Table 4. Shows the recurrence time after surgery

Time of recurrence	Group A (conjunctival autograft)	Group B (pterygium excision followed by mitomycin C)
1 month	1	0
2 months	0	0
3 months	1	1
4 months	0	1
5 months	0	0
6 months	0	1
7 months	0	0
8 months	0	0
9 months	0	0
10 months	0	0
11 months	0	0
12 months	0	0

Table 5. Shows the age of patients with recurrence.

Age of the patient	Group A (conjunctival autograft)	Group B (pterygium excision followed by mitomycin C)
Less than or equal to 50 years	2 (5%)	2 (5%)
Greater than 50 years	0	1 (2.5%)

Table 6. Shows the postoperative complications.

Type of complications	Group A (conjunctival autograft)	Group B (pterygium excision followed by mitomycin C)
Scleral thinning	0	2 (5%)

5.9. Postoperative Complications

Lam *et al.* [19] reported 2 cases (5.57%) of scleral thinning in the group in which they applied MMC.

The present study shows (Table 6) 2 cases (5%) of scleral thinning in eyes with pterygium excision followed by MMC which agrees with previous studies.

V. Conclusion

Conjunctival limbal autograft and postoperative Mitomycin C (0.02%) are both safe and effective adjuncts to primary pterygium surgery. The main prejudices against autografting are the expertise and time required for the procedure. The use of biologic adhesives to fixate the autograft in place may simplify the procedure. Age of the patients was strongly associated with recurrence regardless of which procedure was used.

More research needs to be done to delve into this seemingly innocuous pathology of conjunctiva to effectively manage the disease condition

References

- [1]. R. M. Youngson, "Pterygium in Israel," *American Journal of Ophthalmology*, Vol. 74, No. 5, 1972, pp.954-959.
- [2]. R. Detels and S. P. Dhir, "Pterygium: A Geographical Study," *Archives of Ophthalmology*, Vol. 78, No. 4, 1967, pp. 485-491.<http://dx.doi.org/10.1001/archoph.1967.00980030487014>
- [3]. P. Sivasubramaniam, "Pterygium in Ceylon," *British Journal of Ophthalmology*, Vol. 55, 1971, pp. 55-59.<http://dx.doi.org/10.1136/bjo.55.1.55>
- [4]. M. S. Norn, "Prevalence of Pinguecula in Greenland and in Copenhagen and Its Relation to Pterygium and Spheroid Degeneration," *Acta Ophthalmologica*, Vol. 57, No. 1, 1979, pp.96-105.
<http://dx.doi.org/10.1111/j.1755-3768.1979.tb06664.x>
- [5]. R. T. Rasanayagam, "The Incidence and Racial Distribution of Pterygium in West Malaysia," *Transactions of the Ophthalmological Society of New Zealand*, Vol. 25, 1973, pp.56-59.
- [6]. J. R. Rojas and H. Malaga, "Pterygium in Lima, Peru," *Annals of Ophthalmology*, Vol. 18, No. 4, 1986, pp. 147- 149.
- [7]. A. Meseret, A. Bejiga and M. Ayalew, "Prevalence of Pterygium in Rural Community," *Ethiopian Journal of Health Development*, Vol. 22, No. 2, 2008, pp.191-194.
- [8]. J. S. Kwon and O. Choi, "A Study of Pterygium in Cheju Island," *Yonsei Medical Journal*, Vol. 18, No. 2, 1977, pp. 151-156.
- [9]. S. Duke Elder, "System of Ophthalmology," Vol. VIII, Part I, Henry Kimpton, London, 1965, pp.573-582.
- [10]. S. K. Rao, T. Lekha, B. N. Mukesh, *et al.*, "Conjunctival-Limbal Autograft for Primary and Recurrent Pterygia," *Indian Journal of Ophthalmology*, Vol. 46, No. 4, 1998, pp.203-209.
- [11]. J. A. McCoombes, L. W. Hirst and G. P. Isbell, "Sliding Conjunctival Flap for the Treatment of Primary Pterygium," *Ophthalmology*, Vol. 101, No. 1, 1994, pp. 169- 173.
- [12]. R. Rachmiel, H. Leiba and S. Levartovsky, "Results of Treatment with Topical Mitomycin C 0.02% Following Excision of Primary Pterygium," *British Journal of Ophthalmology*, Vol. 79, No. 3, 1995, pp. 233-236.<http://dx.doi.org/10.1136/bjo.79.3.233>
- [13]. J. Shimazaki, H. Y. Yang, K. Tsubota, *et al.*, "Limbal Autograft Transplantation for Recurrent and Advanced Pterygia," *Ophthalmic Surgery Lasers*, Vol. 27, No. 11, 1996, pp.917-923.
- [14]. R. S. Rubinfeld, R. R. Pfister and R. M. Stein, *et al.*, "Serious Complications of Topical Mitomycin C after Pterygium Surgery," *Ophthalmology*, Vol. 99, No. 11, 1992, pp.1647-1654.
- [15]. B. D. S. Allan, P. Short, G. J. Crawford, *et al.*, "Pterygium Excision with Conjunctival Autografting: An Effective and Safe Technique," *British Journal of Ophthalmology*, Vol. 77, No. 11, 1993, pp. 698-701.<http://dx.doi.org/10.1136/bjo.77.11.698>
- [16]. J. A. Cardillo, M. R. Alves, L. E. Ambrosio, *et al.*, "Single Intraoperative Application versus Postoperative Mitomycin C Eye Drops in Pterygium Surgery," *Ophthalmology*, Vol. 102, No. 12, 1995, pp.1949-1952.
- [17]. P. P. Chen, R. G. Ariyasu, V. Kaza, *et al.*, "A Randomized Trial Comparing Mitomycin C and Conjunctival Autograft after Excision of Primary Pterygium," *American Journal of Ophthalmology*, Vol. 120, No. 2, 1995, pp. 151-160.
- [18]. R. S. Figueiredo, E. J. Cohen, J. A. P. Gomes, *et al.*, "Conjunctival Autograft for Pterygium Surgery: How Well Does It Prevent Recurrence," *Ophthalmic Surgery Lasers*, Vol. 28, No. 10, 1997, pp.99-104.
- [19]. D.S.C.Lam,A.K.K.WongandD.S.P.Fan,"Intraoperative Mitomycin C to Prevent Recurrence of Pterygium after Excision. A 30-Month Follow-Up Study," *Ophthalmology*, Vol. 105, No. 5, 1998, pp. 901-905. [http://dx.doi.org/10.1016/S0161-6420\(98\)95034-5](http://dx.doi.org/10.1016/S0161-6420(98)95034-5)

Dr. Anurag Majumdar. , "A Comparative Study of Treatment of Pterygium by Excision and Limbal Autograft or Augmented with Post-Op Mitomycin C." " IOSR Journal of Dental and Medical Sciences (IOSR-JDMS), vol. 17, no. 11, 2018, pp 45-48.