

Review of outcome of pregnancy with history of previous caesarean section- predictors of safe labour

Dr. Uma kamat, Dr.Taibai Gavde, Dr.K.Nithisha, Dr.AkshataAmonkar,
Dr.Prajaktakatti, Dr.NishaNaik, Dr.Anamika Mishra. , Dr.GuruprasadPednecar
Goa Medical College, Department Of Obstetrics & Gynaecology, Goa, India.
Corresponding author: Dr. Uma kamat

Abstract

Objectives: To study the factors which affect outcome of trial of labor

Materials and Methods: All pregnant women admitted to Goa Medical College, from October 2013 to June 2015, fulfilling the inclusion criteria were enrolled in this study. Data was collected in a proforma, meeting the objectives of the study. Antenatal/ past obstetric predictive factors with respect to successful TOLAC were analysed.

Results: The study included 1,302 pregnant women with history of one or more previous caesarean deliveries, who fulfilled the inclusion criteria. 918 cases (70 %), including 236 subjects with previous 2 LSCS and 682 subjects with previous 1 LSCS; were taken for repeat LSCS (ERCD). Of the 384 cases (30 %) who underwent TOLAC; success rate was 80.5% (309/384).

Younger women (<30 years) (OR=1.61,P= 0.0003), Women with BMI <25(P=0.03), IUGR babies(OR=1.38,P=0.63), Preterm gestation <37 weeks (P= 0.001), Low Birth weights (<3 Kg) (P=0.0045), a history of previous vaginal deliveries (P= 0.0014), Previous history of VBAC (P=0.018), Inter delivery interval >3 years of having a VBAC (OR=2,P=0.03), Bishops score at admission ≥ 10 (P=0.000), Spontaneous onset of labour (P=0.034) were associated with significantly higher rate VBAC. Maternal diabetes (ODM/GDM) (OR=0.6,P=0.59), PROM (OR=0.39,P=0.013) previous LSCS was done for Cephalo-pelvic disproportion (OR=0.5) & Failure to progress (OR=0.29) reduces the chance of VBAC. Place of stay (P=0.75), Educational status (P=0.76), Pre-eclamsia (P=0.22) were not statistically significant factors affecting TOLAC. **Conclusion:** Positive predictors for VBAC from our study were younger age group, lower maternal BMI, absence of maternal diabetes, lower gestational age and birth weight, higher Bishops score at admission, spontaneous onset of labour, vaginal delivery prior to or after caesarean section, malpresentation as indication for prior LSCS and longer inter-delivery interval (>3 years).

Key words : Trial of labor after caesarean (TOLAC), Vaginal Birth After Cesarean (VBAC), Elective repeat caesarean delivery (ERCD)

Date of Submission: 25-11-2018

Date of acceptance: 07-12-2018

I. Introduction

Cesarean section is the delivery of a fetus through an abdominal and uterine incision¹. Since 1985, the obstetricians world-over have considered an ideal rate for caesarean sections to be around 10-15%². Even so, since then, caesarean section rates have increased in both developing and developed countries². Two new Human Reproduction Programme (HRP) studies show that when caesarean section rates are around 10%, neonatal and maternal deaths decrease. When the rate is above 10%, there is no improvement in maternal mortality². In India, until 2005-2006, caesarean section rate was 10.5% of all deliveries, just below the recommended level of 15%; according to a NFHS-3 report³. But in the last decade, the numbers have escalated in many states—reaching as high as 41% of deliveries in Kerala, and 58% in Tamil Nadu; as reported by the ICMR School of Public Health⁴. Reports have shown to be amongst the states with high rates of caesareans, around 25% (NFHS-3 survey)³. Goa and Maharashtra have lower odds of having a private institutional caesarean delivery³.

There are implications of a previous caesarean with respect to maternal, fetal, and neonatal complications in subsequent pregnancies such as placenta accreta and uterine rupture⁵. The risk worsens in patients with multiple repeat caesareans⁵. And, although the absolute risk is small, caesarean section is associated with an increase in severe maternal morbidity; even mortality. Also, higher caesarean rates imply higher health care costs⁵.

Trial of labor after caesarean (TOLAC) and Vaginal Birth After Cesarean (VBAC) has had a measurable impact on decreasing total caesarean deliveries⁵. A trial of labor after caesarean (v/s. repeat caesarean) has been a

polarizing and unresolved issue in obstetrics for over 30 years⁶. There are no randomized controlled trials (RCT) comparing VBAC with repeat cesarean⁶. Hence, conflicting guidelines have surfaced, influenced by emotional, financial, and medico-legal considerations rather than high quality evidence⁶. Owing to safety concerns, VBAC rates have declined drastically leading to an increase in rates of cesarean deliveries⁵. Therefore, in an attempt to reduce cesarean section rates, efforts must be made to increase VBAC rate⁵. Therefore, present study was undertaken to review of pregnancy following previous Cesarean section to identify predictors of VBAC.

II. Materials and Methods

All pregnant women admitted to Goa Medical College, from October 2013 to June 2015, fulfilling the inclusion criteria were enrolled in this study. Data was collected in a proforma, meeting the objectives of the study.

Inclusion criteria: Women with pregnancy above 28 weeks of gestation, with a live fetus, with a history of one or more previous cesarean deliveries.

Amongst all study subjects, one group underwent Elective repeat cesarean, without a trial of labor due to various indications. Subjects in 2nd group were allowed a trial of labor. Amongst the subjects who were allowed trial of labor, a subgroup who failed trial of labor were taken for emergency cesarean for various indications.

Thus, study subjects were classified as: A: women chosen for elective repeat cesarean without a Trial of labor B: women who were given a trial of labor and delivered vaginally C: women who were given a trial of labor, but due to failed trial had to be taken for repeat cesarean

For each study subject, demographic details in the form of age, weight, BMI, residence (rural v/s urban) & socio-economic/ educational status were recorded.

Detailed history including menstrual history (to determine gestational age), past obstetric history was elicited from all subjects. Past obstetric particulars included gravidity, parity, and number of previous cesareans, number of previous vaginal deliveries (including prior VBACs) and timing & indications of last cesarean delivery. Past history of intra/ post-operative complications was recorded. Birth weight & gestation at the time of last cesarean delivery were also recorded.

At admission, per abdomen and per vaginal examination (when indicated) was done to find out malpresentation and to note dilatation and effacement of cervix, station of vertex and adequacy of pelvis for vaginal delivery.

Patients without any contraindication for vaginal delivery were counselled with respect to risks and benefits of trial of labor. After obtaining informed consent subjects willing for VBAC were given trial of labor. During trial of labor, patients underwent careful monitoring for any sign of impending rupture uterus such as maternal tachycardia, FHR variability, scar tenderness etc.

Augmentation of labor was done with artificial rupture of membranes/ oxytocin infusion when indicated. Ventouse/forceps were used in second stage of labour if delay was anticipated.

Antenatal/ past obstetric predictive factors with respect to successful TOLAC were analyzed.

III. Results

Younger age group women <30 years of age were found more likely to have successful TOLAC (OR=1.6) v/s women \geq 35 years of age (OR= 0.28). The difference was statistically significant (P=0.0003)

Table 1: Effect of Age on TOLAC subjects

Age group	TOLAC	VBAC	Failed TOLAC	OR
<30	190	160	30	1.61
31-34	145	120	25	1.40
\geq 35	49	29	20	0.28
Total	384	309	75	
P= 0.0003				

Table 2: Effect of Maternal BMI on TOLAC

BMI	TOLAC	VBAC	Failed TOLAC	OR
<25	176	150	26	1.77
25-29.9	124	99	25	0.94
\geq 30	84	60	24	0.51
TOTAL	384	309	75	
P= 0.0312				

Higher maternal BMI \geq 30 was associated with lower chances of VBAC (OR= 0.51). The difference was statistically significant (P= 0.03).

Table 3: Effect of Maternal Diabetes in TOLAC group

Diabetes	TOLAC	VBAC	Failed TOLAC	OR
Absent	359	293	66	2.5
Present	25	16	9	0.6
	384	309	75	

P = 0.059

Presence of Maternal diabetes (ODM/GDM) reduced the chances of VBAC, although difference was not observed to be statistically significant (OR= 0.6, P= 0.059).

Table 4: Relation of IUGR in TOLAC group

IUGR	TOLAC	VBAC	Failed TOLAC	OR
Absent	345	276	69	0.73
Present	39	33	6	1.38
Total	384	309	75	

P=0.63

IUGR babies were observed to have higher odds of VBAC (OR= 1.38) although difference was not statistically significant (P=0.63).

Subjects with PROM, were less likely to have VBAC; reaching statistical significance (OR=0.39, P=0.013).

Table No. 5 Relation of PROM in TOLAC group

PROM	TOLAC	VBAC	Failed TOLAC	OR
ABSENT	341	281	60	2.51
PRESENT	43	28	15	0.39

P= 0.0128

Table 6: Effect of Gestational age on TOLAC

Gestational age	TOLAC	VBAC	Failed TOLAC	OR
<34Wks	42	42	0	1.84
34.1-37Wks	71	62	9	0.40
37.1-40Wks	250	190	60	0.59
>40wks	21	15	6	

P=0.0000

Preterm gestation 34.1-37 weeks had significantly higher probability of a VBAC (OR= 1.84, P= 0.000). There was no failed TOL with gestation <34 weeks.

Table 7: Relationship of Birth weight on TOLAC

Birth weight	TOLAC	VBAC	Failed TOLAC	OR
<1500	22	22	0	-
1.5-2.499	85	70	15	1.17
2.5-2.99	211	172	39	1.15
3-3.49	54	39	15	0.57
>3.5	12	6	16	0.23

P=0.0045

Infants with higher Birth weights (>=3 Kg) had significantly lower odds of successful

TOLAC compared to lower Birth weights (<3 Kg) [P=0.0045].

Table 8: Relation of Bishops score at admission in TOLAC group

Bishop score	TOLAC	VBAC	Failed TOLAC	OR
6-7	54	24	30	0.12
8-9	186	144	42	0.68
>10	144	141	3	20.14

P=0.000

Higher Bishops score at admission >= 10 was associated with a higher rate of TOLAC success (OR=20.14, P= 0.000)

Table 9: Effect of type of labor on TOLAC

Type of labour	TOLAC	VBAC	Failed TOLAC	OR
Spontaneous	329	271	58	2.09
Induced	55	38	17	0.48

P= 0.034

Subjects who had spontaneous labor had significantly higher chances of a VBAC compared to subjects in whom labor was induced (P=0.034).

No statistically significant difference was found in success of TOLAC in urban v/s rural population (P=0.75). Educational status had no statistically significant correlation with success of TOLAC (P=0.76), although; higher odds (OR=1.2) of VBAC were found in less educated (<X, OR=1.2) subjects. No statistically significant relation was found between presence of preeclampsia & success of TOLAC though OR=2.15 showed increased chances of VBAC with absence of preeclampsia. Subjects with a history of previous vaginal deliveries had significantly higher odds of TOLAC being successful. Odds were higher for patients with multiple prior vaginal deliveries compared to one prior vaginal delivery (OR=3.76 v/s 2.57). Odds of having successful TOLAC in patients with no vaginal deliveries were low (OR=0.27, P= 0.0014).

IV. Discussion

In our study of 1,302 pregnant women with history of one or more previous cesarean deliveries, 918 cases (70 %) (including 236 subjects with previous 2 LSCS and 682 subjects with previous 1 LSCS) were taken for repeat LSCS (ERCD) without trial of labor.

The rate of TOLAC in our study was 30% (384 subjects), while Poddar⁷ and Najma KP⁸ found TOLAC rates of 22% and 20% respectively. Of the 384 cases underwent TOLAC; 309 achieved vaginal delivery. VBAC rate in our study was 80.5% (309/384). AHRQ review⁹ found a comparable VBAC rate of 74% and 79.6%, respectively.

Higher VBAC rate was found in subjects <35 years age (84.2% in <30 years & 82.8% in 31-34 years) v/s \geq 35 years age (59.2 %), reaching statistical significance (P=0.0003). The results were comparable with results shown by Cameron et al⁹ which showed decreasing odds of TOLAC success with increasing age (OR =0.4 for >40 yrs age). Bujold et al¹⁰, (P= 0.005), Doshi²² and Nighat Shaheen¹¹, (P<0.001) also found a statistically significant relation between increasing age and lower chances of VBAC. Educational status was found to have significant effect on outcome of TOLAC (P= 0.76); while King et al¹² in a study of 3,068 subjects found higher odds of VBAC with increase in years of education. Higher maternal BMI \geq 30 was associated with lower chances of VBAC (OR= 0.52, P= 0.03); our results correlated with findings of study by Landon¹³, (OR= 0.55) for BMI > 30 and Juhasz¹⁴, (OR= 0.53, P<0.001) for BMI >29 . Though in a study by Grobman¹⁵ odds ratio was 0.94 in patients with BMI >30 , there were less chances of VBAC with higher BMI. Presence of maternal diabetes (ODM/GDM) reduced the chances of VBAC (OR= 0.6). The results of the present study are in agreement with the findings of Srinivas¹⁶, 2007 (OR=0.68) showing lower odds for VBAC in patients with diabetes. Gyamfi¹⁷ presented still lower odds of VBAC (OR=0.42), in patients with diabetes. Conversely, Dharan et al¹⁸ presented higher odds of VBAC in non- diabetic mothers (OR=1.61, P<0.001). In our study, no statistically significant relation was found between presence of preeclampsia & success of TOLAC, though; OR=2.15 showed increased chances of VBAC with absence of preeclampsia. Relative risk (RR) for failure of TOLAC in subjects with preeclampsia in our study was 1.2. This finding is in accordance to finding by Srinivas et al¹⁶(2006) which stated that subjects with preeclampsia were more likely to fail VBAC (relative risk [RR], 1.56; 95% CI, 1.22 to 2.00). In our study, preterm gestation <37 weeks had significantly higher probability of a VBAC (P= 0.0000) whereas gestation beyond 37 weeks had lower odds of VBAC (OR=0.4 & 0.56 for gestation 37.1-40 weeks & >40 weeks respectively). Gestation <34 weeks had 100% success rate for TOLAC. This finding was in agreement with findings by Landon¹³, Smith¹⁹, Quiñones²⁰ and Srinivas¹⁶ which demonstrate lower odds ratio for VBAC with increasing gestational age. Dhillon²¹, also found lower percentage of VBAC with higher gestational age (P=0.0003). Infants with higher Birth weights (>3 Kg) had lower odds of VBAC (OR= 0.57 & 0.23 for birth weight 3-3.5 Kg & > 3.5 Kg, respectively) compared to lower Birth weights. This result was comparable to findings from study by Cameron⁹, Landon¹³, Doshi²², Smriti Gupta²³, Dhillon²¹ & Nighat Shaheen¹¹. In our study, higher Bishops score at admission was associated with a significantly higher rate of TOLAC success (P= 0.000). This is in accordance with findings by Bujolds¹⁰ 2004 (OR for Bishop's $>5= 2.07$) and Smriti Gupta²³ 2014 (OR for Bishop's $>5= 16$). Both studies found Bishops score to be a statistically significant predictor for VBAC. Subjects who had spontaneous labor had higher chances of a VBAC compared to subjects in whom labor was induced (P=0.034). This result was substantiated by results from Smriti²³ Gonen²⁴ and Landon¹³ published results with similar odds of VBAC in cases of induced labor. In our study, subjects with a history of previous vaginal deliveries had significantly higher odds of TOLAC being successful. The odds of VBAC increased with no. of prior vaginal deliveries (P=0.0014). This finding were comparable to Cameron⁹, Bujold¹⁰, Gyamfi¹⁷, Grobman¹⁵, Srinivas¹⁶, Mercer²⁵ and Bangal²⁶ which demonstrate a high chance of VBAC in subjects with a history of prior vaginal delivery. Patients with a history of vaginal delivery have less chances of inadequate pelvis. This explains the higher success rate of VBAC.

In our study, subjects with a history of previous VBAC had significantly higher odds of TOLAC being successful (OR= 3.8 for VBAC success with one prior VBAC v/s no prior VBAC). There was no failed TOLAC in subjects with ≥ 2 prior VBACs (P=0.0181).

Our results correlate with results by Flamm²⁷, Bujold¹⁰, Gyamfi¹⁷, Landon¹³, Grobman¹⁵ and Doshi.

Maconesz²⁸2001 also demonstrated lower TOLAC failure in subjects with history of prior VBAC (adjusted odds ratio, 0.13; 95% confidence interval, 0.05-0.31). The odds ratio for VBAC failure in our study was 0.21, for patients with history of prior VBAC. Our study found that, Inter delivery interval >3 years had higher odds of having a VBAC (OR=2.0). The relationship was statistically significant (P= 0.03). Doshi²²(2010), found VBAC was associated with significantly higher success rates in women whose inter-conceptual period exceeded two years (P<0.01).

Huang et al.²⁹2002, studied inter-delivery interval <19 month, and found no relationship in less spacing of deliveries compared to larger inter-delivery spacing for patients who had spontaneous onset of labor (P=0.98). However, for induced labor; <19 months interdelivery interval was associated with lower VBAC rate in these studies (P<0.01).

V. Conclusion:

Positive predictors for VBAC from our study were younger age group, lower maternal BMI, absence of maternal diabetes, lower gestational age and birth weight, higher Bishops score at admission, spontaneous onset of labour, vaginal delivery prior to or after caesarean section, malpresentation as indication for prior LSCS and longer inter-delivery interval (>3 years).

Ethics: Institutional ethical committee approval was taken prior to present study.

References:

- [1]. Gabert HA, Bey M: History and development of cesarean operation. *ObstetGynecolClin North Am* 15:591, 1988
- [2]. WHO statement on caesarean section rates: WHO/RHR/15.02; April 2015
- [3]. Sancheeta Ghosh and K.S James: Increasing caesarean section delivery: A threat to urban women's health? 2013 princeton.edu/papers/130873
- [4]. Vaishnavi S D :Analysis of the extent and cost of caesarean section deliveries in Kerala and Tamil Nadu, India: Evidence from DLHS-3 survey 2005-2006
- [5]. David Branch, Robert Silver: Managing the Primary Cesarean Delivery Rate: *Clinical Obstetrics and Gynecology* Volume 55, Number 4, 946-960 ; december 2012
- [6]. James Scott: Update on Vaginal Birth After Cesarean: *Clinical Obstetrics and Gynecology* Volume 55, Number 4, 945; december 2012
- [7]. SumanPoddar: A Study of the Outcome of Subsequent Pregnancy following Previous Cesarean Section; *Journal of Evolution of Medical and Dental Sciences* 2014; Vol. 3, Issue 06, February 10; Page: 1466-1473
- [8]. Najma KP, Smitha K Rao : Pregnancy Outcome in Women With Previous one Cesarean Section; *International Journal of scientific Research*, Volume : 4 | Issue : 1 | January 2015 • ISSN No 2277 - 8179
- [9]. Vaginal Birth After Cesarean: New Insights Prepared for: Agency for Healthcare Research and
- [10]. Quality ; number 191;2010
- [11]. Bujold E, Hammoud AO, Hendler I, et al. Trial of labor in patients with a previous cesarean section: does maternal age influence the outcome? *Am J Obstet Gynecol.* 2004;190(4):1113-1118
- [12]. NighatShaheen, Safia Khalil, Pulwashalfitkha, Prediction of successful trial of labour in patients with a previous caesarean section; *JPak Med Assoc Vol.64, No.5, May2014*
- [13]. King DE, Lahiri K. Socioeconomic factors and the odds of vaginal birth after cesarean delivery. *JAMA.* 1994;272(7):524-529
- [14]. Landon MB, Leindecker S, Spong CY, et al. The MFMU Cesarean Registry: factors affecting the success of trial of labor after previous cesarean delivery. *Am J Obstet Gynecol.* 2005;193(3 Pt 2):1016-1023
- [15]. Juhasz G, Gyamfi C, Gyamfi P, et al. Effect of body mass index and excessive weight gain on success of vaginal birth after cesarean delivery. *Obstet Gynecol.* 2005;106(4):741-746
- [16]. Grobman WA, Gilbert S, Landon MB, Spong CY, Leveno KJ, Rouse DJ. Outcomes of induction of labor after one prior cesarean. *Obstet Gynecol.* 2007 Feb. 109(2 Pt 1):262-9.
- [17]. Srinivas, S.K., et al., Predicting failure of a vaginal birth attempt after cesarean delivery. *ObstetGynecol*, 2007. 109(4): p. 800-5.
- [18]. Gyamfi C, Juhasz G, Gyamfi P, Stone JL. Increased success of trial of labor after previous vaginal birth after cesarean. *Obstet Gynecol.* 2004;104(4):715-719.
- [19]. Dharan VB , Srinivas SK, Parry S, Ratcliffe SJ, Macones G., Pregestational diabetes: a risk factor for vaginal birth after cesarean section failure? *Am J Perinatol.* 2010 Mar;27(3):26570.
- [20]. Smith GCS, White IR, Pell JP, Dobbie R. Predicting cesarean section and uterine rupture among women attempting vaginal birth after prior cesarean section. *PLoS Med.* 2005;2(9):e252
- [21]. Quiñones JN, Stamilio DM, Paré E, Peipert JF, Stevens E, Macones GA. The effect of prematurity on vaginal birth after cesarean delivery: success and maternal morbidity. *Obstet Gynecol.* 2005 Mar. 105(3):51924
- [22]. Dhillion et al, Vaginal birth after cesarean section (VBAC) versus emergency repeat cesarean section at teaching hospitals in India: an ICMR task force study; *Int J ReprodContraceptObstet Gynecol.* 2014; 3(3) 592-597
- [23]. HareshDoshi et al. Prognostic factors for successful vaginal birth after cesarean section — Analysis of 162 cases; *Journal of Obstetrics and Gynecology of India* 12/2010; 60(6):498-502
- [24]. Smriti Gupta, Shanti Jeeyaselan et al, An Observational Study of Various Predictors of Success of Vaginal Delivery Following a Previous Cesarean Section; *J ObstetGynaecol India.* 2014 Aug; 64(4): 260-264
- [25]. Gonen R , Tamir A, Degani S, Ohel G , Variables associated with successful vaginal birth after one cesarean section: a proposed vaginal birth after cesarean section score; *Am J Perinatol.* 2004 Nov;21(8):44753
- [26]. Mercer BM, Gilbert S, Landon MB, Spong CY, Leveno KJ, Rouse DJ, et al. Labor outcomes with increasing number of prior vaginal births after cesarean delivery. *Obstet Gynecol.* 2008 Feb. 111(2 Pt 1):28591
- [27]. Bangal et al, Vaginal Birth after Cesarean Section; *N Am J Med Sci.* 2013 Feb; 5(2): 140-144
- [28]. Flamm BL, Geiger AM. Vaginal birth after cesarean delivery: an admission scoring system. *Obstet Gynecol.* 1997;90(6):907-910

Review of outcome of pregnancy with history of previous caesarean section- predictors of safe labour

- [29]. Macones GA , Hausman N, Edelstein R, Stamilio DM, MarderSJ. Predicting outcomes of trials of labor in women attempting vaginal birth after cesarean delivery: a comparison of multivariate methods with neural networks; Am J Obstet Gynecol. 2001 Feb;184(3):40913
- [30]. Huang WH , Nakashima DK, Rumney PJ, Keegan KA Jr, Chan K , Interdelivery interval and the success of vaginal birth after cesarean delivery, Obstet Gynecol. 2002 Jan;99(1):414

Dr. Uma kamat. , ““Review of outcome of pregnancy with history of previous caesarean section- predictors of safe labour..” ” IOSR Journal of Dental and Medical Sciences (IOSR-JDMS), vol. 17, no. 12, 2018, pp 17-22.