

A Study on Teenage Pregnancy and its Maternal and Fetal Outcome

Dr. Nibha¹, Dr. Anubha Vidyarthi²

¹Junior Resident, Department of Obstetric and Gynaecology, RIMS, Ranchi

²Corresponding author – Professor & HOD, Department of Obstetric and Gynaecology, RIMS, Ranchi

Abstract: Background: Teenage pregnancy rates varies vastly between different countries and different regions within a country. Such pregnancies are seen mostly amongst the poorer and less educated sections of the society. In developed countries majority of teenage pregnancies occur to unmarried girls unlike developing countries including India where teenage pregnancies occur to married girls and are associated with early marriages. Aim & Objectives: To find out the incidence and to evaluate the effect of pregnancy in teenage girls and its maternal and fetal outcome. Material & Methods: It is a one year hospital based clinical prospective study of all teenage pregnant females admitted to a tertiary care centre. Required details were collected on a proforma by taking history and following up the patients from admission to delivery. MS Excel was used for statistical purpose. Results: Teenage pregnancy 9.76% of the total Obstetric admissions. In this study 57.7% teenage pregnancies were associated with complications. The major maternal complications were Preterm labour 16.6 %, Hypertensive Disorders of Pregnancy 11.66 %, Premature Rupture of Membranes 15 %, anemia (13.3 %). Low Birth Weight 25 % and stillbirths 1.9 % were major adverse fetal outcomes. Conclusion: Teenage pregnancy is still a common occurrence. It has adverse impact on the health of teenage mothers leading to various adverse maternal and fetal outcome.

Date of Submission: 15-10-2019

Date of Acceptance: 31-10-2019

I. Introduction

World Health Organization defines Teenage Pregnancy as “any pregnancy from a girl who is 10-19 years of age”, the age being defined as her age at the time the baby is born. Often the terms “Teenage pregnancy” and “Adolescent pregnancy” are used as synonyms.

Teenage pregnancy is coming up as one of the most important social and public health problems all over the world with varying prevalence rates.

About 16 million women 15–19 years old give birth each year, about 11% of all births worldwide. Ninety-five per cent of these births occur in low- and middle-income countries. Although adolescents aged 10–19 years account for 11% of all births worldwide, they account for 23% of the overall burden of disease. Due to pregnancy and childbirth. Fourteen percent of all unsafe abortions in low- and middle-income countries are among women aged 15–19 years.

Stillbirths and death in the first week of life are 50% higher among babies born to mothers younger than 20 years than among babies born to mothers 20–29 years old. The rates of preterm birth, low birth weight and asphyxia are higher among the children of adolescents, all of which increase the chance of death and of future health problems for the baby.

Although adolescent marriage is cognizable offence in India, it is still a common practice in many parts of the country. A high fertility rate, social customs, poverty and ignorance make early marriage a common feature in this part of the world. So this teenage group is most vulnerable and maximum attention should be diverted to them. With modernization, teenage pregnancy rate is rapidly declining in developed countries, but it is still high in developing countries like India. The scenario of teenage pregnancy in developed countries is quite different from that of the developing countries and have distinctly different rates of pregnancy as well.

In India, teenage pregnancy constitutes 8–14% of total pregnancies. Complications of pregnancy and childbirth in women between 15–19 years of age are the leading cause of mortality among women in India. Hence teenage pregnancy is a serious problem today all over the world and more so in developing countries like India. Objectives of this study is to find out the incidence and to evaluate the effect of pregnancy in teenage girls (13–19 years) and its maternal and fetal outcome.

II. Materials and Methods

This is a one year clinical prospective study carried out in the Department of Obstetrics and Gynecology, at a tertiary care Centre of Jharkhand, India. All pregnant females admitted to the Hospital in the age group of 13-19 years during the study period were included and all pregnant females equal to or more than 20 years admitted to the Hospital during the same period were excluded from the study. The required details were collected by history taking and following up the patients from admission to delivery. Data was compiled in MS Excel sheet. Frequencies and percentages were computed for the results.

III. Result and Observation

In the present study there were 104 teenage mothers admitted during the study period amongst the total obstetric admissions of 1967, giving an incidence of 9.76 % of teenage pregnancy .Out of the 104 teenage pregnancies, 60 (57.69%) were associated with complications and the remaining 44(42.30%) were without any complication

Table 1: Distribution according to presence of complication

Complication	No. of pregnancies	Percentage
With complication	60	57.70
Without complication	44	42.30
Total	104	

The most common complication associated with teenage pregnancy was Preterm Labour (10). Amongst 7 mothers who had Hypertensive Disorders, 3 had Preeclampsia and 4 had Gestational Hypertension. 9 of them had premature rupture of membranes, 7 of them had hypothyroidism and 6 pregnant teenagers were found to be anaemic(Table No.2).

Table 2: Complications in teenage pregnancy

Complications	No. of cases	Percentage
Preterm labour	10	16.6
PROM	9	15
Malpresentation	9	15
Medical disorder	8	13.3
Hypertensive disorders	7	11.66
IUGR	7	11.66
CPD	6	10
Oligohydramnios	2	3.33
Multiple pregnancies	1	1.6
IUFD	1	1.6

Out of the 104 teenage mothers who delivered, majority i.e 57 (54.8%) had normal vaginal delivery and 47 (45.19%) had cesarean section.

Out of the 57 teenage mothers delivered by Cesarean Section, 22 mothers had more than one indication for Cesarean Section. Majority of Cesarean Section were due to Fetal Distress. It was followed by Malpresentation (16) of which 14 was for Breech presentation and one each for face and brow presentation. The other indications for C- Section in teenage mothers were oligohydramnios, severe preeclampsia, CPD and Obstructed Labour (TableNo. 3).

Table 3: Indication for Cesarean Section Indication Number Percentage Fetal distress 22.1 % Malpresentation 14.4 % Oligohydramnios 6.7 % Severe preeclampsia 4.8 % CPD 4.8 % Obstructed labour 3.8 %

Indication	Number	Percentage
Fetal distress	23	22.1
Malpresentation	15	14.4
Oligohydramnios	7	6.7
Severe preeclampsia	5	4.8
CPD	5	4.8
Obstructed labour	4	3.8

Majority of the babies (104) born out of such pregnancies were healthy babies. The most common adverse fetal outcome noted in the study was Low Birth Weight babies (26 babies). Amongst 7 babies who needed NICU admissions, 4 were Low Birth Weight babies. There were 2 Still born babies.

Table .4 Fetal outcome of teenage pregnancy

Fetal outcome	Number	Percentage
Active and healthy	70	67.3
Low birth weight	26	25
NICU admissions	7	6.7
Still birth	2	1.9

IV. Discussion

In the present study, 9.76% of the study population were teenage pregnancies which correlated with incidences of other studies, Gazala Yasmin et al, Ambadekar NNet et al, Dubashi SS et al (3, 16, 17). Incidence of C-Section in the present study was 57%. The number of vaginal deliveries were more in the teenage group probably because of the smaller size of the babies. Similar results were found in other studies Trivedi S S et al, Mukhopadhyay et al, Mahavarkar S. H et al and Sagili Het al [5,6,7,8].

However two studies have shown higher rate of caesarian sections. This may be due to a difference in the setting, Nwobodo et al and Jimoh et al [9,10]. Cases of obstructed labour are higher in the teenage group and may be due to weaker physical strength of the teenage mothers.

The incidence of Preeclampsia was 6.25%. Although anemia, and preeclampsia were higher in the cases, the difference was not statistically significant. Similar results were found in Mahavarkar S. H et al [7]. A higher study population is needed to conclude this finding. As pregnant teenagers often receive inadequate antenatal care, their anemia during labour and the postpartum period usually get worse. In developing countries more than 25% of teenage mothers were found to be anemic as revealed in studies conducted by Saxena et al, (11) Bhalerao et al (12) And Rahman MM et al. (13) In contrast to it our study found a lower incidence (3.125%). Consistent with previous studies, cases of preterm, LBW and foetal distress were higher in teenage group. But unlike other studies, the difference was not statistically significant [14,5,6,7,15]. 5.

V. Conclusion

The present study concludes that Preterm labour, Hypertensive Disorders of Pregnancy, Premature Rupture of Membrane, anemia, malpresentations, IUGR, IUFD were major maternal complications; Low Birth Weight and NICU admissions were major adverse fetal outcomes. Teenage pregnancy today, still represent one of the most important public health problems. There is no doubt that the obstetrical problems can be managed by modern medicine and so the risk of Teenage pregnancy can be diminished. The health care provider should consider Teenage pregnancy as a „high risk“ pregnancy and should educate the pregnant teenagers to have more number of antenatal visits so that the signs and symptoms of various complications of teenage pregnancy could be recognized at the earliest. Attention should be given to the use of various screening and diagnostic tests and to the interventions needed if any complication does occur during the course of pregnancy or labour. Proper monitoring of the progress of labour is important to prevent prolonged labour. Education of the female child can play a significant role in delaying marriage and hence delaying childbearing, thus protecting the young girl from being exposed to the various complications of teenage pregnancy. There is a need to promote the use of Contraceptives amongst the married teenagers and ensuring the availability of contraceptives at a wider scale. Good antenatal and intranatal services, good neonatal services, contraceptive services and abortion services, all together can minimize the various risks associated with teenage pregnancies to a large extent. With all these measures, we can hope for a world-wide decline in the trend of teenage pregnancy rates and complications in the years to come.

References

- [1] World Health Organization, United Nations Population Fund: Married adolescents: no place of safety. Geneva: WHO-UNFPA; 2006.
- [2] Adolescent pregnancy- Issues in adolescent health and development, WHO discussion papers on adolescence. WHO, 2004 Pg 86.
- [3] Gazala Yasmin, Aruna Kumar, Bharti Parihar. "Teenage Pregnancy - Its Impact On Maternal And Fetal Outcome". International Journal of Scientific Study. 2014(6):9-13.
- [4] Snigdha Kamini, Dr. Krishna Veni Avvaru. Teenage Pregnancy: Maternal and Fetal Outcomes. IOSR Journal of Dental and Medical Sciences. 2014(4): 41-44.
- [5] Trivedi S S., and Shikha Pasrija. Teenage pregnancies and their obstetric outcomes, Tropical doctor, 37(2), 2007, 85-88.
- [6] Mukhopadhyay, Prianka, R. N. Chaudhuri, and Bhaskar P, Hospital-based perinatal outcomes and complications in teenage pregnancy in India, Journal of health, population, and nutrition, 28(5), 2010, 494.
- [7] Mahavarkar S. H, C. K. Madhu, and V. D. Mule., A comparative study of teenage pregnancy, Journal of Obstetrics & Gynecology, 28(6), 2008, 604-607.
- [8] Sagili H, N Pramy, Karthiga P, Mariano M, and P. Reddi Rani.. Are teenage pregnancies at high risk? A comparison study in a developing country, Archives of gynecology and obstetrics, 285(3), 2012, 573-577. [9] Nwobodo, Emmanuel I., and Kasimu
- [9] Adoke, Obstetric Outcome of teenage pregnancies at a tertiary care hospital in Sokoto, Nigeria, Tropical Journal of Obstetrics and Gynaecology, 22(2), 2006, 168-170.
- [10] Jimoh, Adiodun S., and Ishaq-F. Abdul. Outcome of teenage pregnancies in Ilorin, Nigeria, Tropical Journal of Obstetrics and Gynaecology, 21(1), 2005, 27-31.

- [11] Saxena P, Salhan S, Chatopadhyay B, Kohli MPS, Nandan D and Adhish SV. Obstetrics and perinatal outcome of teenage and older primigravidas – A retrospective analysis. *Health and Population: Perspectives and Issues*. 2001; 33(1):16-22.
- [12] Bhalerao AR, Desai SV, Dastur NA, Daftary SN. Outcome of teenage pregnancy. *J Postgrad Med*. 1990;36:136–9.
- [13] Rahman MM, Hasan M, Akhter S, Sultan P. Adolescent pregnancy complication and wastage in Bangladesh. *J. Nepal Paediatr*. 2010; 30(3):147-153.
- [14] Kumar A., Singh T, Basu S, Pandey S, & Bhargava V, Outcome of teenage pregnancy. *The Indian Journal of Pediatrics*, 74(10), 2007, 927-931.
- [15] Yadav, Siddhartha, Dilip C, K. C. Narayan, Rajesh K M, Achyut S et al, Adversereproductive outcomes associated with teenage pregnancy." *McGill Journal of Medicine: MJM*, 11(2), 2008, 141.
- [16] Ambadekar NN, Khandait DW, Zodpey SP, Kasturwar NB, Vasudeo ND. Teenage pregnancy outcome: *Indian J Med Sci*. 1999;53:14-7.

Dr. Nibha. "A Study on Teenage Pregnancy and its Maternal and Fetal Outcome." *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)*, vol. 18, no. 10, 2019, pp 42-45.