

Incidence of Eosinophilia Inrural Population of Northern Region of Kashmir in India (A Study at Tertiary Care Hospital)

Dr. Qazi Imtiaz Rasool¹, Dr. Javedhussain Bhat²Dr. Nadeemarafiq³,
Dr. Sehar Aslam⁴, Dr. Nasir Choudary⁵Abdul Aziz Parry⁶

¹Head of Department, Associate Professor, Department of Physiology Government; Medical College Baramulla J&K India

²Resident in Department of Physiology Government; Medical College Baramulla J&K India

³ Assistant Professor Department of Physiology Government; Medical College Baramulla J&K India

⁴ Resident in Department of Physiology Government; Medical College Baramulla J&K India

⁵ Ex. Assistant Professor GMC Srinagar and Consultant Department of Pathology Majmaah Hospital SA

⁶ Junior laboratory Technicians Immuno-pathology Central Laboratory Government; Medical College Baramulla J&K India

Corresponding author: Dr. Qazi Imtiaz Rasool

Abstract:

Background: Eosinophilic count varies and fluctuates demographically because of culture, exercise and environment i.e. seasonal allergen exposure, the presence of more than 500/ μ l, and classified as mild less than 1500/cu mm, moderate (Hyper-eosinophilia)1500- 5000 /cu mm, severe Hypereosinophilicsyndromeson 2 occasions \geq 1 month apartmore than 5000 /cu mm. which is usually seen in middle and elderly subjects and may be primary or secondary type^{1,2}. Significant tissue eosinophilia can occur without an elevated blood count and vice versa ³. Percentages generally above 5% are regarded as elevated, but is not reliable indicator as it depends on total leucocytes, hence absolute count (AEC) should be considered for diagnosis of eosinophilia. AEC also does not accurately predict organ damage, hence evaluation of unexplained eosinophilia in an asymptomatic subject, needs further exploration, due the wide variety of potential pathogens, in underdeveloped countries which is a challenging task.

Method; A prospective tertiary Hospital study of non-randomized cohort of 2138 subjects aged 10 yrs.to 75yrs attending OPD or admitted was conducted. The aim of the study was to determine the incidence of eosinophilia in the rural border areas in Kashmir, an area where almost negligible literature is available on this subject and comparison with other places.

Conclusion: In tropical countries like India, high percentage of eosinophilia is expected, due to unhygienic living condition and poor sanitation. Eosinophilia was found in only5.09%,of subjects in this study which is very less as compared to other parts of India i.e. UP rural population were common etiologies was parasitic, protozoal or fungal infestations and infections, though this province of Baramulla Kashmir near the border hence underdeveloped,being at an altitude of >5128ft, cold, lash green hilly area with allergy related problem is prone for chronic respiratory problems..

Key Words;-Eosinophils, Allergy, Helminthic, Immunology

Date of Submission: 26-04-2019

Date of acceptance: 11-05-2019

I. Introduction

Eosinophil's are granulocytic white blood cells, recognized by Paul Ehrlich about 125 years ago ⁴and is the second least in number from the same series in peripheral blood. They are also found sparsely in tissues like the medulla of thymus, lower gastrointestinal tract, ovary, uterus, spleen and lymph nodes, normally not seen in the lungs, skin, esophagus and other internal organs, there presence in them indicates disease ^{5,6}. Eosinophilia is count above 0.5 $10^9/l.$, higher in neonates than in adults and the values gradually fall in the elderly. In some institution higher values are recommended^{7,8}

There is no sex or ethnic variation, in subjects with primary eosinophilia without organ involvement, no treatment may be necessary. Diurnal variation may be as much as 100% ^{9,10}, lowest counts in the morning 10 A.M-12A.M a time at which endogenous steroids are the lowest and the highest in the midnight 00 A.M- 04 A.M. The circulating life span of eosinophil is 6-12 hrs. Before it migrates to tissue In contrast, blood hyper eosinophilia (HE) is defined as an AEC of $\geq 1.5 \times 10^9/$ which is relatively rare and should prompt a thorough evaluation for an underlying cause. The majority of cases of eosinophilia are Secondary (reactive) with allergic disorders , drugs ,infectious diseases, gastrointestinal disorders, vasculitis's, rheumatologically disease,

respiratory disease, neoplasms (non-hematological and hematological in which the eosinophil's are not part of the neoplastic clone, lymphocytic variant hyper eosinophilic syndrome ,miscellaneous causes graft-versus-host disease, Gleich syndrome ¹¹Primary (clonal) eosinophilia Meteorologically seasonal variations usually starts in late April, peaks in May and gradually subsides in late June. Blood picture depend upon the physiological adaptation of the body to the new set of environmental situation.^{12, 13,}but unlike the neutrophils it can recirculate and have a much longer life. ¹⁴. In tertiary care hospital situated against a backdrop of a rural population, it was observed that surgeries were delayed as a result of an increased blood eosinophilia, due to bronchoconstriction during anesthesia. The clinical manifestations becomes important to prevent organ damage by these inflammatory substances. Refined hematological computerized equipment, are spotting a lot of new, repeatedly unanticipated cases of eosinophilia ¹⁵

II. Aims and Objectives


1. A study was undertaken to find out the Hospital based prevalence of blood eosinophilia in a rural population
2. Compare these figures with blood eosinophilia in a hospital based urban and rural population of other places as very less literature is available in this tropical country

III. Materials And Methods

This was a prospective study which was conducted in the Pathology department of the newly created govt. medical college and tertiary hospital in rural area of northern province in Kashmir.. A total number of **2138** subjects who visited our hospital were tested. These included OPD, indoor and emergency ward subjects. Subjects with hematological malignancy were excluded from our study, a detailed medical history that including an assessment for allergic disorders, such as asthma, eczema, urticarial and hay fever. Skin rashes or lymphadenopathy, cardiorespiratory and gastrointestinal symptoms i.e, fever drenching night sweats, weight loss, pruritus. A detailed travel and drug history was also recorded. A thorough physical examination was done. Total complete blood count was done IRIF-I count 5 hematology analyzer. Having normal range of 0.04-0.50 foreosinophils, 5.09% were found to have eosinophilia. The arbitrary eosinophil count accepted for a diagnosis of CEL, NOS [16]. and idiopathic HES ¹⁷ (Chusid et al, 1975) was not accepted because the new version of equipment analyses at an molecular level hence permits certain entities to be diagnosed with a lower count (because hypo granular eosinophil's may not be counted accurately by automated counters).Also the cytological features of eosinophil's are not helpful in the differential diagnosis given that striking abnormalities can occur in reactive and clonal eosinophil's are cytological fairly normal¹⁸ Eosinophilia was based on counts and not defined by the percentage (typically <5 percent in healthy individuals), as it varies with the total WBC count and the proportion of other WBC lineages (eg, neutrophils, lymphocytes).

IV. Results

Total number of subjects registered was 2138 Graph -1 shows subjects were 764 (35.73%) males and 1374(64.26%) were females. Age group of subjects varied from 10 yrs. to 75 yrs. Graph -2 shows Eosinophilia distribution of subjects according to severity of eosinophilia. 109, 47in males whereas 62 in females



.Compared with other states of India, mean eosinophil counts were not highereven in peak pollen month i.e. April as compared to the counts takes month earlier but the results had no statistically significance. The chief complaints of our subjects were related to fever, cough, rashes, and breathlessness, wheezing etc. other major group of subjects had symptoms of abdominal pain, diarrhea, pallor, pica and loss of appetite.

V. Discussion

Despite being the second least represented granulocyte subpopulation in the circulating blood, eosinophils are receiving a growing interest from the scientific community, due to their complex pathophysiological role in a broad range of local and systemic inflammatory diseases as well as in cancer and thrombosis. Eosinophilia was detected in 5.09% of the population who visited the hospital. The disparity in the incidence in males and females is probably due to the fact that the overall number of male subjects visiting the hospital is low as compared to females during this period which is beginning of farming season. Eosinophilia was seen in subjects of all age groups. Many subjects came with chief complaints associated with eosinophilia i.e. allergy. In others it was an incidental finding.¹⁹ Eosinophilia in the bloodstream can be diagnosed by a simple blood test. If problem is in body tissues then diagnoses will involve examination of the relevant tissue normal bone marrow contains between 1% and 6% eosinophils and these produce an eosinophil count in the peripheral blood of 0.05–0.59 $10^9/l$.²⁰ In tropical countries like India, the most common cause of eosinophilia is parasitic infestation, unhygienic living condition and poor sanitation that accounts for high worm infestation in India. These findings differ from another study from India done a decade ago, which reported parasitism as a more common cause of eosinophilia than allergic rhinitis.²¹ As the subjects who are otherwise well with mild to moderate eosinophilia may not require further testing but persistent eosinophilia (0.5-1.53 $10^9/l$), irrespective of suspected organ damage, should be considered for additional testing for an underlying^{22,23,24} In another study done (10.7%) subjects visiting the hospital of northern India had (62%) mild eosinophilia, of which (71.2%) were males and (28.8%) females.²⁵ In another study, all, 47% of the study population had mild and 30% had moderate among rural population of barabanki and neighbor districts were no access is to hygienic mode of stool disposition as reactive type based on clinical suspicion. In one of studies carried SKIMS Kashmir for patients with RA shows almost more than 70% had normal count though suffering from autoimmune diseases. The eosinophil count was lower in RA patients with high disease activity.

Table No. 1 Comparison at national and international level

	No. of Females	No. of Males	High count	normal
OUR STUDY	1374	764	5.09%	94%
1. SKIMS Kashmir with RA	103(76.9%)	31(23.1%)	29(21.6%)	105 (78.4%)
2. Barabanki and neighboring districts of eastern Uttar Pradesh ²⁷	73 (64.03%)	41(35.97%)	69(60%)	45(40%)
3. Incidence of eosinophilia in rural population in North India: A Study at tertiary care hospital ²⁸	71.2%	28.8%	(10.7%)	89.3%
4. Blood and tissue eosinophilia ²⁹	Eosinophilia 14. %	Eosinophilia 11.17%	100 (25%)	400(75%)
5. Seasonal variation in eosinophil count in normal healthy adult females ⁽³⁰⁾	-----	45	40 (88.88%) SEASONAL INCREASE IN APRIL.	5(11.2%)

The present study was conducted with an aim to document the prevalence of eosinophilia. Identification of possible cause's may be a next step as a significant proportions of children are suffering from anemia, under nutrition, anorexia, pain abdomen, cough and hence long term morbidity. All these chronic problems are preventable and treatable with simple corrective steps of which the most important ones are access to safe drinking water and toilet.

Further significant advancement in recent times has been made in understanding the mechanisms of eosinophil production, apoptosis, and how eosinophil immunology contributes to both host defenses against infections, tissue damage and autoimmune diseases. New syndromes like hyper- eosinophilic syndromes (HESs) that effects almost any organ has been documented where there is moderate to severe eosinophilia which leads to extensive extracellular deposition of eosinophil-derived proteins in tissue as demonstrated by immunostaining.³¹ Eosinophilia- myalgia syndrome which is a multisystem disease, chronic and fatal characterized by counts $> 1000 /\mu l$ without any cause. Subjects with $AEC \geq 1500 / \mu l$ should have a CBC repeated in one to two weeks to determine if the eosinophilia is transient, stable, or rising; the CBC should be repeated even when eosinophilia is detected incidentally in an asymptomatic patient.

VI. Conclusion

Eosinophilia appears to be a common occurrence in rural population around the tropical countries across the globe due to poverty, hygiene, diet and in addition Kashmir being an border area where facilities are not available easily, besides high altitude > 5128 ft. effects, cold area and green hilly belt area is more prone to allergy of various pollen at the peak of allergic season i.e. march to June are more vulnerable to COPD where counts can be a biomarker indicator, our study showed lower incidence of eosinophilia. In a significant proportion of subjects no definite etiology of eosinophilia could be zeroed upon, all ages were effected, majority

showed mild $0.5 - 1.53 \times 10^9/l$ The results proved that good living habits, hygiene, diet, education contributed for the lower incidence of mild category only, that may not require further testing. For asymptomatic subjects with eosinophilia $<1500/\mu l$, it may be reasonable to postpone a repeat CBC and evaluation for a month or longer. However, it is important to first ensure that there are no clinical findings suggestive of eosinophilic end-organ damage, no history of travel or residence in helminthic-endemic areas, no history of drugs ingestion and no features suggestive of a malignancy (eg, significant anemia or thrombocytopenia, splenomegaly, lymphadenopathy before deferring the evaluation. concurrent neutrophilia may suggest an infection or inflammatory condition, basophilia may reflect a myeloid malignancy or allergic disorder, and lymphocytosis may be associated with a lymphoid malignancy and persistent eosinophilia at least $1-5.3 \times 10^9/l$), irrespective of suspected organ damage, should be considered for additional testing for an underlying cause, i.e., stool test, allergic tests, marrow smears, ^{32,33} fluorescence in situ hybridization (FISH) or PCR. Secondary (reactive) eosinophilia should be confirmed or excluded at an early stage.

Further long-term studies could answer the question of immunological response to changes that represent or identify individuals who could be potential asthmatics or at high risk of developing seasonal allergic rhinitis in future nevertheless, the prevention of morbidity by the diagnosis and prompt treatment of parasitic helminthic infection is also an important task in these subjects.

References

- [1]. Andersen CL, Siersma VD, Hasselbalch HC, et al. Eosinophilia in routine blood samples and the subsequent risk of hematological malignancies and death. *Am J Hematol* 2013;88(10):843-847. [OpenUrlPubMed](#)
- [2]. Schulte C, Krebs B, Jelinek T, Nothdurft HD, von Sonnenburg F, Löscher T. Diagnostic significance of blood eosinophilia in returning travelers. *Clin Infect Dis* 2002;34(3):407-411. [OpenUrlAbstract/FREE Full Text](#)
- [3]. Harrison's Principles of Internal Medicine 18th edition vol;1 Long;Fauci;Kasper;Hauser;Jameson;Loscalzo page 1332
- [4]. Juhan Yoon, Jens U. Ponikau, Christopher B. Lawrence, Hirohito Kita. Innate Antifungal Immunity of Human Eosinophils Mediated by a $\beta 2$ Integrin, CD11b. *Jour Immunol*, 2008, 181: 2907-2915.
- [5]. Spry CJF, ed. Eosinophils: A Comprehensive Review and Guide to the Scientific and Medical Literature. Oxford, UK: Oxford University Press; 1988.
- [6]. Gleich, G.J. (2000) Mechanisms of eosinophil-associated inflammation. *The Journal of Allergy and Clinical Immunology*, 105, 651-663.
- [7]. Brigden ML, Horak MG. Incidence and clinical significance of unsuspected eosinophilia discovered by automated WBC differential counting. *Lab Med* 1993; 24: 173-6
- [8]. Wardlaw AJ. Eosinophils in the 1990s: new perspectives on their role in health and disease. *Postgrad Med J* 1994; 70: 536-52.
- [9]. Rothenberg ME. Eosinophilia. *N Engl J Med* 1998;338:1592-1699
- [10]. Uhrbrand H. The number of circulating eosinophils Normal figures and spontaneous variations. *Acta Medica Scandinavica* 1958;160:99-104
- [11]. Gleich, G.J., Schroeter, A.L., Marcoux, J.P., Sachs, M.I., O'Connell, E.J. & Kohler, P.F. (1984) Episodic angioedema associated with eosinophilia. *New England Journal of Medicine*, 310, 1621-1626.
- [12]. Deng-Bang WEI, Zhang JM, Lian WEI, Hong- Yan YU, Wang XJ. Seasonal changes in blood physiological parameters in plateau zokors. *Acta Zoologica Sinica* 2006; 52: 871-7.
- [13]. Kristal EB, Froom P, Harari G, Shapiro Y, Green MS. Seasonal changes in red blood cell parameters. *Br J Haematol* 2006; 85: 603-7.
- [14]. Tefferi, A. Blood eosinophilia: A new paradigm in disease classification, diagnosis, and treatment. *Mayo Clin Proc* 2005;80:75.
- [15]. Zucker-Franklin D. Eosinophilia of unknown etiology: a diagnostic dilemma. *Hosp Pract* 1971;7: 119-27.
- [16]. Bain, B.J., Gilliland, D.G., Vardiman, J.W. & Horny, H.-P. (2008) Chronic eosinophilic leukaemia, not otherwise specified. In: *World Health Organization Classification of Tumours of Haematopoietic and Lymphoid Tissue* (eds. by S.H. Swerdlow, E. Campo, N.L. Harris, E.S. Jaffe, S.A. Pileri, H. Stein, J. Thiele, J.W. Vardiman), pp. 51-53, 4th edn. IARC Press, Lyon.
- [17]. Chusid, M.J., Dale, D.C., West, B.C. & Wolff, S.M. (1975) The hypereosinophilic syndrome: analysis of fourteen cases with review of the literature. *Medicine (Baltimore)*, 54, 1-27.
- [18]. Barbara J Bain. *Blood Cell Morphology in Health and Diseases in Dacie & Lewis Practical Hematology*, 9th Edition, Eds: S. Mitchell Lewis, Barbara J Bain & Imelda Bates, Pg: 93-94.
- [19]. Nutman TB. Asymptomatic peripheral blood eosinophilia redux: Common parasitic infections presenting frequently in refugees and immigrants. *Clin Infect Dis* 2006;42(3): 368-369.
- [20]. Valent, P., Klion, A.D., Horny, H.P., Roufosse, F., Gotlib, J., Weller, P.F., Hellmann, A., Metzgeroth, G., Leiferman, K.M., Arock, M., Butterfield, J.H., Sperr, W.R., Sotlar, K., Vandenberghe, P., Haferlach, T., Simon, H.U., Reiter, A. & Gleich, G.J. (2012) Contemporary consensus proposal on criteria and classification of eosinophilic disorders and related syndromes. *The Journal of Allergy and Clinical Immunology*, 130, e609
- [21]. Makkar A, Rohtagi A, Goel A, et al. A study of clinical profile and spontaneous course of eosinophilia. *JK Science* 2005 7:199-201.
- [22]. Udwardia FE. Tropical eosinophilia. In: Herzog H. *Pulmonary eosinophilia: Progress in respiration research*. 7th ed. Basel: Karger 1975:35-155.
- [23]. Dacie YV, Lewis SM. Basic hematologic techniques. *Practical Hematology*, 7th ed. Edinburg: Churchill Livingstone 1991:37-68.
- [24]. Weller PF. Eosinophilic syndromes. In: Goldman L, Bennett JC, eds. *Cecil Textbook of Medicine*, 25th Ed. Philadelphia: W.B. Saunders 2000:1151-1154
- [25]. Incidence of Eosinophilia in Rural Population in North India: A Study at Tertiary Care Hospital Rimpri Bansal, Anureet Kaur, Anil K Suri, Puneet Kaur, Monika Bansal and Rupinderjeet Kaur Dept. of Pathology, Gian Sagar Medical College and Hospital, Banur, Dist Patiala, Punjab, India DOI: 10.21276/APALM.2017.1044

- [26]. Eosinophilia in rheumatoid arthritis patients and its relation to disease activity: A single center experience from Kashmir, India The Egyptian Rheumatologist Volume 39, Issue 2, April 2017, Pages 65-67
- [27]. Alarming High Incidence of Eosinophilia in Barabanki and Neighboring Districts of Eastern Uttar Pradesh: A Prospective Hospital-Based Study Neha Thakur Narendra Rai Journal of Tropical Pediatrics, Volume 62, Issue 6, December 2016, Pages 500–502.
- [28]. Incidence of Eosinophilia in Rural Population in North India: A Study at Tertiary Care Hospital DOI:Rimpi Bansal, Anureet Kaur, Anil K Suri, Puneet Kaur, Monika Bansal and Rupinderjeet Kaur 10.21276/APALM.2017.1044
- [29]. Seasonal variation in eosinophil count in normal healthy adult females Jane Alam*, Fayaz Qureshi, Muhammad Imran Suliman, Zaheeruddin Qureshi ;Gomal Journal of Medical Sciences July-December 2008, Vol. 6, No. 2
- [30]. Blood and Tissue Eosinophilia: A Study at the MVJ Medical College & Research Hospital with Review of Literature Yogendra V1, Roopa AN2*, Shammem Shariff 3 GUHS J Med Sciences, October 2011 / Vol 1 / Issue 4
- [31]. Valent P, Klion AD, Horny HP, et al . Contemporary consensus proposal on criteria and classification of eosinophilic disorders and related syndromes. J Allergy Clin Immunol 2012;130(3):607-612, e9. OpenUrlCrossRefPubMedWeb of Science
- [32]. Webb JKG.. Etiology of tropical eosinophilia. Indian Journal of Tuberculosis 1961;8 (3):95-98.
- [33]. Udwardia FE. Tropical eosinophilia. In: Herzog H. Pulmonary eosinophilia: Progress in respiration research. 7th ed. Basel. Karger 1975:35-155.

Dr. Qazi Imtiaz Rasool¹ “Incidence of Eosinophilia Inrural Population of Northern Region of Kashmir in India (A Study at Tertiary Care Hospital).” IOSR Journal of Dental and Medical Sciences (IOSR-JDMS), vol. 18, no. 5, 2019, pp 08-12.