

Presentation and Diagnostic accuracy of FNAC and USG in detecting Breast Cancer in tribal females of rural West Bengal

DR. Sanhita Mukherjee (MBBS, MD)¹, DR. Amiya Kumar Sarkar (MBBS, MD)², Dr Ashis Kumar Saha (MBBS, MS)³

¹Professor, Department of Physiology, Calcutta National Medical College, 32 Gorachand Road, Kolkata-700014

²Associate Professor, Department of Physiology, Calcutta National Medical College, 32 Gorachand Road, Kolkata-700014

³Professor, GENERAL SURGERY Professor, Department of General Surgery, Calcutta National Medical College, 32 Gorachand Road, Kolkata-700014

Corresponding author: DR. Amiya Kumar Sarkar (MBBS, MD)

Abstract: The present study tried to have a comparative analysis among the sensitivity, specificity and positive predictive value of Ultrasonography and Fine needle Aspiration Cytology in the detection of breast cancer among tribal population.

92 tribal females attending surgical outdoor with breast lumps were included in the study. There demographic parameters, mode of presentation, time lapse since first noticed, presence and absence of well-known breast cancer risk factors were studied. All patients were subjected to FNAC & USG and the findings of these two investigations were compared with the Histopathology reports when the patients came for a follow-up. Data were analyzed using the Statistical Package for Social Sciences software version 17 for Windows. Percentage, sensitivity, specificity, and positive predictive values (PPV) were calculated.

Among 92 patients 68 (74%) were diagnosed to have benign breast disease and 24 (26%) had carcinoma breast. Although the age of presentation of these patients did not differ from that of the western population but time lapse since first noticed and different well-known risk factors of breast cancer e.g. parity, usage of oral contraceptive pills differ markedly in the study group. The Sensitivity, Specificity & PPV of Ultrasonography were 91.67 %, 91.18% & 78.58 % respectively and that of FNAC was found as 95.83 %, 97% & 92 %.

The study observed a late presentation of breast lump in tribal rural women of Bankura, West Bengal. Although the accuracy of the screening procedures like USG and FNAC are comparable with that of western world.

Key Words: Breast cancer, Tribal female, Presentation, Diagnostic accuracy

Date of Submission: 17-01-2020

Date of Acceptance: 04-02-2020

I. Introduction:

Breast cancer is the most prevailing cancer in women worldwide. It is the principal cause of death from cancer among women globally (1). Obesity, lack of physical exercise, consumption of alcohol, smoking and use of oral contraceptive pills are the most common risk factors of breast cancer (2). The uncontrollable risk factors are genetic mutations and family history of it. In general, women from higher socio-economical groups, regardless of their race/ethnicity, have higher rates of occurrence of breast cancer. Most of the breast cancers are asymptomatic. In some cases, however, the first sign may be a new lump or mass in the breast. According to the American Cancer Society, any of the following atypical changes in the breast can be an indication of breast cancer e.g. swelling of all or part of the breast, dimpling or skin scar, pain in the breast or nipple, nipple retraction, redness, or increased thickness of the nipple or breast skin, a nipple secretion other than breast milk, a lump in the axilla. (3) However most of the above observations are done on the basis of Western Population.

At present, screening for breast cancer in women is regarded as effective way to reduce incidence of breast cancer. A rapid and accurate diagnostic tool is crucial and fundamental in the implementation of breast cancer screening. The USG-FNAC (ultrasonography- fine needle aspiration cytology) techniques have acceptable diagnostic validity indices and can be used for early staging of breast cancer patients. (4)

Recent studies indicate that cancer incidences, mortality and survival rate vary according to geographical location and tribe. (5) Socioeconomic factors have been shown to strongly influence the stage of disease presentation (6-9). Less access to medical care, awareness, illiteracy of the population leads to late breast cancer screening (10, 11).

Bankura District of West Bengal has much lower Human Development Index or HDI (0.53), Income Index (0.26) and education index (0.62) as compared to other districts of West Bengal. (12) According to 2001

census 10.36% tribal population is living in this region. Little is known about the presentation of breast cancer among the tribal women of rural West Bengal, India, where the culture, life-style, socio-economic level and literacy rate in female differ greatly from Western countries.

Therefore the purpose of the present study was to investigate the pattern of presentation, and role of various other clinic-pathological parameter in tribal female patient with breast lump presenting to a rural medical college and hospital of Bankura. The present study is also an attempt to have a comparative analysis among the sensitivity, specificity and positive predictive value of Ultrasonography and Fine needle Aspiration Cytology in the detection of breast cancer among this tribal population.

II. Methods & Materials:

This descriptive study was carried out in B. S. Medical College, Bankura, West Bengal, in the period of January 2011 to June 2012. All the tribal female patients presenting at surgical outpatient department of BSMC&H with clinically palpable breast lump either solitary or multiple, unilateral or bilateral irrespective of age, socioeconomic status and duration of complaint within this time period were included in the study. The tribal status of the subjects were recorded from their 'scheduled tribe' certificates issued by Govt. of West Bengal. Institutional ethical committee clearance and informed consent of all patients were obtained. Male patient with breast related diseases, patients with acute and tender breast lump like breast abscess, recurrent breast lump after definitive surgery for confirmed malignancy were excluded from the study. Total 92 patients fulfilled the criteria. They were admitted and were evaluated by detailed history, thorough physical examination were done. The variables studied included demographics, modes of presentation, time since first noticed and reason for delayed presentation.

Appropriate investigations of the subjects like ultrasound, FNAC and incisional or Trucut biopsy were done in the department of Surgery, Radiology and Pathology of BSMC, Bankura. The FNAC was performed using a 23G needle and 10 ml syringe with an average of four to six passes with constant suction. The breast lumps of the patients were under local/general anesthesia, and were sent for histopathologic examination. Where the lump was very big, incision biopsy was done at proper site and with proper orientation of the incision (With due consideration of future operation and other modalities of treatment, if they become necessary). The post-intervention patients were kept for observation overnight and discharged the next morning with the advice to continue antibiotics for 48 hours and to attend the outpatient department or emergency room in case any problem. The follow up visits were scheduled at 3 weeks, 3 months. The histopathology reports obtained at follow-up visits were used to compare the result of USG and FNAC. High frequency ultra sound image was used for diagnosis. Solid lesions with irregular shape, irregular margin, non-uniform distribution of internal echotexture, shadowing and calcifications in hypo-echoic lesions are suggestive of cancer by USG.

Data were analyzed using the Statistical Package for Social Sciences software version 17 for Windows. Sensitivity, specificity, and positive predictive values (PPV) were calculated.

III. Results:

A total of 92 female patients presenting with breast lump were included in the study. Their mean age was 39 ± 6.019 years (range 22 years – 54 years). Among 92 patients, 68 (74%) were diagnosed to have benign breast disease and 24 (26%) had carcinoma breast.

26.26 % (n = 26) patients with Benign breast disease were within the age range of 21-30 years and 20.6% were below 19 years of age whereas 50% (n=12) patients with breast cancer were more than 50 years of age.

All the 92 patients were from remote rural areas. There were 73 (79.34%) illiterate patients whereas 19 (20.65%) had primary to graduate level education. In 90.22% (n=83) cases the lump was detected accidentally by the patient herself, 47 (51.08%) of them took clinical opinion after 6 months of self-detection and 36.95% (n=34) within 2 to 6 months. (Table 1) 81.5 % patient took no clinical advice in this interim period whereas 17 (18.5%) patients used alternative medicines (e.g. homeopathy) during this time. 13 out of these 17 patients (76.47%) presented with breast cancer.

The most common presentation in both benign and carcinoma breast were painful lump (52%) and in most of the cases the pain was non-cyclical (81%). Other than pain and lump in the breasts, the most common symptoms were fever (19.5%, n=18), nipple discharge (10.86%, n=10), wound discharge (9.98%, n=9), redness (6.52%, n=6) and opposite breast pain (4.34, n=4). (Table 2) Only 12 (13.04%) patients gave history of benign breast disease among mother and first degree relatives, none of them had history of malignant breast disease in the family. 3 patients (12.5%) with malignant breast disease and 6 patients (8.82%) with benign breast tumor had history of Oral Contraceptive Pill intake for more than 2 years. Out of 24 breast cancer patient 19 (79.66%) were multipara and only 5 (20.83%) were nullipara.

The Sensitivity, Specificity & PPV of Ultrasonography were 91.67 %, 91.18% &78.58 % respectively and that of FNAC was found as 95.83 %, 97% & 92 %. (Table 3)

IV. Discussion:

Carcinoma of breast continues to be a major exterminator of women all around the world. The incidence and sequence of this disease, however, differ significantly between developed and under-developed countries (13, 14). In developing countries many factors modulate the presentation of the breast diseases such as non-uniform distribution of health care facilities, deficiency of knowledge about the disease, poorness, and social habits of the families avoiding women exposure.

In this study 26.26 % (n = 26) of patients with Benign breast disease (n=68) were within the age range of 21-30 years and 20.6% were below 19 years of age whereas 50% (n=12) patients with breast CA (n=24) were more than 50 years of age. This observation is in accordance with a previous study done by Vissya Shanti et al and Reeni Malik et al in 2003 (15).

Majority of our study subjects belonged to remote, under-resourced areas presenting late in the course of the disease (>6 months after detection of lump). Painful breast lump were there most common presentation and most of the lumps were observed by accidental self-detection. 79.34% (n=73) of patients in this study were illiterate and only 19 (20.65%) had primary to graduate level education. This shows the unawareness of this rural tribal female community about breast cancer symptoms and the health care facility available to detect it in early stage. This is consistent with the findings of many previous similar studies (16, 17, 18)

Only 3 patients (12.5%) with malignant breast disease and 6 patients (8.82%) with benign breast tumor of the present study had history of OCP intake for more than 2 years. This finding differs from the previous studies that show women using oral contraceptives (birth control pills) have a slightly greater risk of breast cancer than women who have never used them. (19, 20, 21)

Out of 24 breast cancer patient 19 (79.66%) were multipara and only 5 (20.83%) were nullipara. This observation is also not in agree with the previous study reports that demonstrate that women who have had no children or who had their first child after age 30 have a slightly higher breast cancer risk. (22, 23)

The Sensitivity, Specificity & PPV of Ultrasonography were 91.67 %, 91.18% & 78.58 % respectively and that of FNAC was found as 95.83 %, 97% & 92 %. This result is almost comparable to the observations obtained by S. Usmani et al (24) and Yumjaobabu Singh TaKhellambam et al (25) among western population. Thus it shows that it is not inaccuracy of screening programme in underdeveloped countries but lack of either basic advances in health literacy or absence of tailored approaches to help women with low literacy to navigate local health care systems, are major factors in delaying the diagnosis of breast cancer (26).

Limitation: It is a single institutional study. Further studies on the current topic using multiple centers and with a large sample size are therefore recommended to have more clear idea about the pattern of the presentation of breast cancer among rural tribal women.

Conclusion: The study observed a late presentation of breast lump in tribal rural women of Bankura, West Bengal. Although the accuracy of the screening procedures like USG and FNAC are comparable with that of western world low literacy rate, poor breast cancer awareness and reluctance to obtain health care facility are the major barrier of the early detection of breast cancer in these regions.

References:

- [1]. Susan G. Komen Breast cancer foundation. Racial and ethnic differences in breast cancer. www.komen.org.
- [2]. Courtney M. Townsend Jr., R. Daniel Beauchamp, B. Mark Evers, Kenneth L. Mattox. Sabiston Textbook of Surgery: The Biological Basis of Modern Surgical Practice. 19th Edition. Elsevier Saunders; 2012.
- [3]. F. Brunnicardi, Dana Andersen, Timothy Billiar, David Dunn, John Hunter, Jeffrey Matthews, Raphael E. Pollock. Schwartz's Principles of Surgery. 9th Edition. McGrawHill; 2010.
- [4]. Xi-Wen Wang & Yun-Hui Xiong & Xiao-Qing Zen, Hai-Bo Lin, Qing-Yi Liu. Diagnostic Accuracy of Ultrasonograph Guided Fine-needle Aspiration Cytologic in Staging of Axillary Lymph Node Metastasis in Breast Cancer Patients: a Meta-analysis. *Asian Pacific J Cancer Prev*, 13 (11), 5517-5523.
- [5]. K. Glanz, R. T. Croyle, V. Y. Chollette, and V. W. Pinn, "Cancer-related Health Disparities in Women," *American Journal of Public Health* 93, no. 2 (February 2003): 292-98; Centers for Disease Control and Prevention (hereinafter CDC), Division of Cancer Prevention and Control 2008/2009, "HealthDisparitiesCancer, http://www.cdc.gov/cancer/HealthDisparities/pdf/0809_hf_fs.pdf (accessed 25 May 2007); J. Swan and B. K. Edwards, "Cancer Rates among American Indians and Alaska Natives," *Cancer* 98, no. 6 (2003): 1262-72.
- [6]. Gordon NH, Crowe JP, Brumberg DJ, Berger NA. Socioeconomic factors and race in breast cancer recurrence and survival. *Am J Epidemiol*. 1992;135:609-618.
- [7]. Berg JW, Ross R, Latourette HB. Economic status and survival of cancer patients. *Cancer*. 1977;39:467-477.
- [8]. Dayal HH, Power RN, Chiu C. Race and socio-economic status in survival from breast cancer. *J Chronic Dis*. 1982;35:675-683.
- [9]. Freeman HP, Wasfie TJ. Cancer of the breast in poor black women. *Cancer*. 1989;63:2562-2569.
- [10]. Fletcher SW, Harris RP, Gonzalez JJ. et al. Increasing mammography utilization: a controlled study. *J Natl Cancer Inst*. 1993;85:112-120.
- [11]. Blendon RJ, Aiken LH, Freeman HE, Corey CR. Access to medical care for black and white Americans. *JAMA*. 1989;261:278-281.
- [12]. Census 2001. censusindia.gov.in/2011-common/censusdataonline.html.
- [13]. Jakesz R. Breast Cancer in Developing Countries: Challenges for Multidisciplinary Care. *Breast Care* 2008; 3: 4-5.
- [14]. Agarwal G, Pardeep PV, Aggarwal V, Yip CH, Cheung PS. Spectrum of breast cancer in Asian women. *World J Surg* 2007; 31: 1031-40.

Presentation and Diagnostic accuracy of FNAC and USG in detecting Breast Cancer in tribal ..

[15]. Vissa Shanthi, Kashmir Ali, Nandam Mohan Rao ,Baddukonda Appala, Rama Krishna, Kuppali Venkata Murali mohan. CLINICOPATHOLOGICAL STUDY OF BREAST LESIONS IN FEMALES WITH ASSESSMENT OF CORRELATION BETWEEN TUMOR GRADE AND PROGNOSTIC FACTORS. J Biosci Tech, Vol 2 (5),2011,367-378.

[16]. Nair MK, Sankaranarayanan R, Nair KS, Amma NS, V arghese C, Padmakumari G, et al. Overall survival from breast cancer in Kerala,India,in relation to menstrual, reproductive and clinical factors. Cancer 2006; 71: 1791-6.

[17]. Okobia MN, Osime U. Clinico-pathological study of carcinoma breast in Benin City. Afr J Reprod Health 2001; 5: 56-62.

[18]. Arshad Mehmood Malik, Rafique Pathan, Noshad Ahmed Shaikh, Jawaid Naem Qureshi, Khamiso Altaf Hussain Talpur. Pattern of presentation and management of Ca breast in developing countries. There is a lot to do. J Pak Med Assoc. Vol. 60, No. 9, September 2010.

[19]. Chlebowski RT, Anderson GL, Gass M, et al. Estrogen plus progestin and breast cancer incidence and mortality in postmenopausal women. JAMA. Oct 20 2010;304(15):1684-1692.

[20]. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer. Lancet. Oct 11 1997;350(9084): 1047-1059.

[21]. Beral V, Reeves G, Bull D, Green J. Breast cancer risk in relation to the interval between menopause and starting hormone therapy. J Natl Cancer Inst. Feb 16 2011;103(4):296-305.

[22]. Kelsey JL, Gammon MD, John EM. Reproductive factors and breast cancer. Epidemiol Rev. 1993;15(1):36-47.

[23]. Lambe M, Hsieh C, Trichopoulos D, Ekbom A, Pavia M, Adami HO. Transient increase in the risk of breast cancer after giving birth. N Engl J Med. Jul 7 1994; 331(1):5-9.

[24]. S Usmani, , H Khan, N Ahmed, F Marafi, and N Garvie. Scintimammography in conjunction with ultrasonography for local breast cancer recurrence in post-mastectomy breast. The British Journal of Radiology, 83 (2010), 934–939.

[25]. Yumjaobabu Singh TaKhellambam, Sunil Singh lourembam, opendro Singh Sapam, raju Singh KSheTrimaYum, bhupon Singh ningThoujam, TouSif Khan. Comparison of Ultrasonography and Fine Needle Aspiration Cytology in the Diagnosis of Malignant Breast Lesions. Journal of Clinical and Diagnostic Research, 2013.

[26]. Gilani GM, Kamal S, Akhter AS. A different study of Breast Cancer Patients in Punjab, Pakistan. J Pak Med Assoc 2003; 53: 478-81.

Table 1: Demographic parameters of the patients presenting with breast lump

Address of the Patient(n=92)	Literacy level (n=92)	Mode of Detection(n=92)	Time lapse since lump first noticed(n=92)
Rural remote areas 100%	Illiterate: 73 (79.34%)	Self-accidental detection 90.22% (n=83)	2-6 months: 81.5%
	Primary to graduate level: 19 (20.65%)	Physical examination by Physician 7.6%(n=7)	≥ 6 months: 36.95%
		Other 2.18% (n=2)	

Table 2: Symptoms and Signs of presentation of Breast lump:

Symptoms (n=92)		Pattern of Complaint(n=92)		Gross Appearance (n=92)		
Unilateral Breast Lump	88%	Pain	52%	Benign (n=68)	Malignant(n=24)	
Bilateral breast lump	12%	Fever	19.5%	Localized growth	100%	
		Discharge from nipple	10.86%		Localized growth	50%
		Redness	6.52%		Ulcerated growth with destruction of nipple	33.33%
		Pain in the opposite breast	4.34%		Infected fungating growth	16.67%

Table 3: Results of Ultrasonography and FNAC in the diagnosis of Breast Cancer

Test	Biopsy +/- Breast Cancer (n=24)	Biopsy -/ Breast Cancer (n= 68)	Total (n=92)	Sensitivity, Specificity & PPV
Ultrasonography				
Benign	2	62	64	Sensitivity 91.67 %, Specificity 91.18% & PPV 78.58 %
Malignant	22	6	28	
FNAC				
Benign	1	66	67	Sensitivity 95.83 %, Specificity 97% & PPV 92 %
Malignant	23	2	25	

DR. Amiya Kumar Sarkar (MBBS, MD), etal. "Presentation and Diagnostic accuracy of FNAC and USG in detecting Breast Cancer in tribal females of rural West Bengal." *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)*, 19(1), 2020, pp. 01-04.