

Association Of Maternal Serum Pigf Levels As A Biomarker In Preeclampsia: A Case Control Tertiary Care Hospital Based Study

Krishnaveni.Changalvala, Kiranmayee.P, Raghuveer C. V., Kalyani R , Venkateshu K. V.

¹(Departments Of Anatomy/ Rims, /Knruhs, Adilabad, India. &Sdumc&Sduaher, Kolar,India)

²(Department Of Cell Biology And Molecular Genetics², Sdumc, Sduaher, Kolar, India)

³(Department Of Pathology, Sdumc, Sduaher, Kolar, India)

Abstract:

Background: Preeclampsia (PE) is a multi-systemic disorder with circulatory pressure of more than 140/90 mm Hg and proteinuria with 0.3 g in a 24 h on dipstick emerging after 20 weeks of pregnancy. Due to incomplete spiral arterial transformation and incomplete cytotrophoblastic cell division. Due to that there will be improper trophoblastic development and improper implantation. As there is imbalance in Placental Inhibiting Growth Factor (PlGF) levels in PE women, it constricts the blood vessels and leads to endothelial dysfunction. Low circulating PlGF precedes the manifestation of clinical disease in PE pregnancies and intrauterine growth restriction in fetus. So the serum levels of PlGF in PE can be a biochemical or diagnostic marker needs to be focussed more. The present study aim is to compare the maternal serum PlGF levels in normotensive pregnant women to PE women in early and late gestational weeks.

Materials and Methods: In this case-control study, 150 were preeclamptic women as cases, and 150 as normotensive pregnant women as controls, a total of 300 subjects were associated in the present study. A 5ml of maternal venous blood was collected, the serum was separated and stored at -80°C till the analysis. By using commercially available Enzyme-Linked Immuno Sorbent Assay kits from Chongqing Biospes Co., Ltd., (suppliers: Info bio-Company, New Delhi) was measured with microplate reader at 450nm (MerilyzerEiaquant Company).

Results: Out of 300 pregnant women, 46 pregnant women were early gestational weeks and 254 were late gestational weeks. The statistical analysis was performed by Statistical Packages for Social Sciences Software 22. The area under the receiver operating characteristic curve is 0.93, 92% sensitivity, and 77% specificity. The significance in the maternal serum PlGF levels was calculated by the Mann-Whitney U test. By comparing the cases and controls it was found that maternal serum PlGF was significantly higher in PE women with z value =2.65 and U= 9253 and with p-value 0.005 significance.

Conclusion: Maternal serum PlGF levels in late gestational weeks in PE and in controls were significant when compared to early gestational weeks. So PlGF can be used as a late preeclamptic diagnostic marker in southeastern Kolar population.

Keyword: Pro-angiogenic Pregnancy Marker, Placental Inhibiting Growth Factor

Date of Submission: 01-05-2024

Date of Acceptance: 10-05-2024

I. Introduction

Pre-eclampsia (PE) is a pregnancy-specific, vascular angiogenesis imbalance associated with endothelial dysfunction. The abnormalities in the development of the placental vasculature occur at early or late gestation but the development of clinical manifestations was evident at the 20th week of PE pregnancy. Placental growth factor (PlGF), a component of the vascular endothelial growth factor (VEGF) family, play a key role in the remodeling process of maternal arteries in normal pregnancy. This growth factor is released into the bloodstream by the migration of trophoblasts. The placental growth factor (PlGF/ PLGF/PGF) belongs to the cysteine-knot superfamily.[1] Placental protein binds to form a heterodimer with VEGFR-1/FLT-1/soluble variant sFLT-1 (soluble fms Feline McDonough Sarcoma -like tyrosine kinase-1). It also binds to neuropilin receptor 1 and 2, recognized in the placenta but its role is still to be highlighted in detail.[2] PlGF is secreted in placental tissues, neurons, and pathological conditions like colon and breast carcinomas. It signals through the vascular endothelial growth factor receptor-1 VEGFR1/FLT1(fms-related tyrosine kinase-1) which stimulates endothelial cell proliferation and migration.[1] PlGF concentrations in early pregnancy are lower in women who

later develop PE. However, as the pregnancy progresses, there is a reciprocal relationship between sFLT-1 and PIGF. Low PIGF levels are evident for abnormal placentation, small for gestational age babies which further leads to abnormal growth in the second half of pregnancy. So it supports the hypothesis that PIGF is an indicator of abnormal placentation.[2] Circulating PIGF is prominently elevated in pregnancy from the placenta. The function of PIGF in the placenta is development and maturation of the placental vascular system. Placental expression of PIGF dominates from the second trimester when the uteroplacental circulation is advancing, with myometrial spiral arteries remodeling beginning at 16–18 weeks of gestation.[2] Concentrations of PIGF are low in the first trimester of an uncomplicated pregnancy and increases from 11 to 12 weeks onwards to a peak at 30 weeks, after which it decreases. Standard levels of PIGF are dependent on gestational age, normal levels ranging from approximately 141 pg ml⁻¹ at around 30 weeks gestation to 23 pg ml⁻¹ at term.[3] The main role of PIGF in tissues is angiogenesis, imbalance in the concentration of PIGF levels leads to pathological ischemia. PIGF expression in trophoblast (transcriptional activity) is suppressed by hypoxia and further related to intrauterine events.[2] With a combination of other angiogenic factors such as soluble endoglin and biomarkers-Doppler sonography are more useful for predicting early-onset PE. The angiogenic factors correlate with disease severity, can be detected several weeks before clinical presentation, and have predictive value for the diagnosis of severe-early onset PE.[4] So PIGF is specific to pregnancy and vascular development.[3] There was a lot of discrepancy to deliberate about the PIGF levels in the diagnostic precision between early onset (EO) and late onset (LO) of PE.[5] In this regard, the study aims to compare the maternal serum concentration of PIGF levels in normotensive pregnant women to PE women in early and late gestational weeks.

II. Material And Methods

After approval from the central ethics committee, the information was collected subsequently written informed consent from each study subject involved in this investigation were collected.

Study Design: Prospective Case-Control study design was adopted.

Study Location: This was a tertiary care teaching hospital-based study done in Department of Obstetrics & Gynecology in R.L. Jalappa Research Institute & Teaching Hospital.

Study Duration: August 2017 to October 2018.

Sample size: 300 patients.

Sample size calculation:

The sample size was estimated on the basis of a single proportion design. We assumed that the confidence interval of 10%, confidence level of 95%, power 80% with mean difference of 35%. The sample size actually obtained for this study was 142 subjects for each group. We planned to include 300 subjects.

Subjects & selection method: Out of 300 Pregnant women, 150 pregnant women who were determined with PE dependent on the American College of Obstetrics and Gynecologists. The remaining 150 were controls considered as normotensive pregnant women under early and late PE.

Diagnostic Criteria of Preeclampsia: According to the American College of Obstetrics and Gynaecologist PE was characterized as new beginning of hypertension (Blood Pressure \geq 140/90 mm Hg) and proteinuria (urine dipstick of \geq 1+) following 20 weeks of gestation. Without proteinuria, additionally PE was characterized by extreme hypertension (\geq 160/110 mm Hg) and hemolysis, low platelets count, and increased liver enzymes (HELLP syndrome) or indications (i.e. headache/migraine, visual changes, and right upper quadrant of stomach discomfort). Renal deficiency in creatinine \geq 1.1 mg/dl.[6] PE further ordered into early and late-beginning, which is related with various pathogenesis, clinical outcome, and prognosis.[7] The PE women with the \geq 20th week of gestation were incorporated as cases. Another 150 normotensive pregnant women with the \geq 20th week of gestation without any pregnancy complexity till delivery were controls.

Inclusion criteria:

Pregnant women who were diagnosed with preeclampsia after the 20th week of gestation, both primigravida and multigravida with reproductive age group.

Healthy normotensive pregnant women after the 20th week of gestation without any complications till delivery both primigravida and multigravida

Exclusion criteria:

- 1.Pregnant women with gestational hypertension,
- 2.Persistent hypertension,
- 3.Gestational diabetes,
- 4.History of multiple premature births and abortions,
- 5.Past pregnancy with an abnormal fetus and with developmental anomalies,
- 6.Thrombophilia-like disorders.

Procedure methodology:

Under complete aseptic precautionary measures, 5ml of maternal venous blood was collected in red-capped vacutainers. The serum was isolated by centrifugation and stored at -80°C till further investigation. Before 2 days the samples were stored at -20°C at that point steadily to 4°C and afterward gradually brought to room temperature. By using commercially available Enzyme-Linked Immuno Sorbent Assay (ELISA) kits from Chongqing Biospes Co., Ltd., (suppliers: Info bio-Company, New Delhi) was measured by following the complete protocol carefully according to guidelines. The maternal serum analysis was completed by estimating with an ELISA microplate reader at 450 nm (MerilyzerEiaquant Company).

Statistical analysis

The outcomes were interpreted by verifying normal distribution further analyzed by the Mann-Whitney U test due to non-distribution invariance. The AUC (Area Under the receiver operating characteristics (ROC) curve) estimation value of PIGF is 0.93. The statistical analysis of mean, median, sensitivity, and specificity with cut-off variables was determined and decipher as an ROC curve by utilizing Statistical Packages for Social Sciences Software SPSS (version 22.0; SPSS Inc, Chicago, IL, USA)

III. Result

Out of 300 pregnant women based on the gestational weeks, the early gestational weeks were 46 and late gestation was 254. In that 24% were early gestational PE subjects and 7% were early gestational weeks in control subjects. The late gestational weeks in PE were 76% and in controls 93%. Based on their gravida status primigravida were 149 (PE 79 and controls 70) and multigravida were 151 (PE- 71 and controls -80) subjects. In 52.6% primigravida were PE patients.

Table 1: Angiogenic Marker-:The statistical analysis was analyzed by using SPSS software. After analysing the normal distribution curve, the statistical significance of maternal PIGF levels in cases and controls was calculated by the Mann-Whitney U test Table 2. By comparing cases and controls the maternal serum PIGF levels were significantly higher in PE cases (z value =2.65)U= 9253. The p-value is significant with (≤ 0.05) especially if we compare with early and late gestational weeks, in late gestational weeks the PIGF levels were extremely significant. The AUC curve value is 0.93, 92% sensitivity, and 77% specificity (**Figure1**)

Content	Median± SE	IQ Range	AUC value	Sensitivity	Specificity	P-value
In both cases & controls	90 ± 60.11	196-48.5= 147.5	0.93	92%	77%	U=9253 Z=2.65757 P value=0.00782
PE	82.5± 10.2	142-39.5 = 102.5	NA	NA	NA	NA
Controls	99.25± 6.5	186.5-53=133.5				
Severe PE	82.5 ±6.4	140-41.5= 98.5				
Mild PE	82.25±28.8	151.5-35.25 =116.25				

PIGF Statistical Values:Significant with p value ≤ 0.05 , IQ Range- Inter Quartile Range, AUC- Area under the receiver –operating characteristics, SD-Standard Deviation, SE-Standard Error, NA- Not Applicable

Figure 1: AUC (Area Under the receiver operating characteristics (ROC) curve) The estimated value of PIGF is 0.93 with 92% sensitivity and 77% specificity with p value 0.007(significant with ≤ 0.05)




Table 2. Statistical Outcome in Early and Late Preeclamptic Women (Cases) versus Early and Late Normotensive Pregnant Women (Controls)

The gestational weeks in cases and controls were analysed by the Mann-Whitney U test. In gestational age, we found a major difference when compared to late and early PE. In late PE the gestational weeks are 37 and whereas in early PE was 30 weeks and in controls, late gestation was extended to 39-40 weeks. So, this designates that due to PE there were more probabilities of early delivery which affects the development of the fetus. If we compare the serum PIGF levels in early and late PE, in early gestational weeks of PE the serum PIGF levels were low in maternal circulation compared to late gestational weeks. In late gestational weeks in controls and PE, the levels were highly significant than early gestational weeks of PE and in controls significant with p value ≤ 0.0001 ,

Content	Gestational Weeks	Median \pm SE	P value
PIGF levels	Early Gestational Weeks in PE cases	139 \pm 18	0.3572
	Early Gestational Weeks in controls	140.3 \pm 62	
	Late Gestational Weeks in PE cases	615 \pm 128	0.00001
	Late Gestational Weeks in controls	195 \pm 65.8	

Numbers of subjects involved in the study were as follows: Early Gestational Weeks in PE cases n=36; Early Gestational Weeks in controls n=10; Late Gestational Weeks in PE cases n=114; Late Gestational Weeks in controls n=140. n- Number of subjects, SE- Standard Error,

IV. Discussion

For healthy fetal growth there should be a proper uterine establishment with healthy placental angiogenesis. This plays a vital role in placental perfusion. Imbalance in angiogenesis and vasculogenesis factors affects the remodeling of uterine spiral arteries and angiogenesis which can be one of the reasons for the pathogenesis of PE.[8]In PIGF, 4 altered forms were found. PIGF may respond to hypoxia or ischemia, and other comorbidities like obesity, ethnicity.The PIGF can be a diagnostic marker in individuals especially in southern part of India. [1]PIGF expression is persistent in placental hypoxia due to underdeveloped uteroplacental circulation. However, studies have been shown that PIGF expression in PE placental tissue is either increased or unchanged. PIGF expression regulation is unknown, but several mechanisms have been investigated. PIGF expression may be influenced by inflammation, which is evident in sepsis.[2]In a mouse-transgene designed model by Chau et al.,study revealed that PIGF promotes angiogenesis for the fetoplacental circulation and supports trophoblast development but the decidual invasion is not influenced. Consequently, abnormal spiral artery remodeling in mice does not lead to placental insufficiency or abnormal blood pressure regulation.[2]In a biomarker study conducted by Giardini et al., suggested that PIGF is a biochemical marker not only for PE but also for placental dysfunction.[9] In a multi-centric study, PIGF levels range from 12 to 3000 pg/mL can be a preferable early clinical marker of adverse suspected PE pregnancy.[10]In another study by Lai et al., Myatt et al and Youssef et al., AUC range from 0.61 and 0.734. Poon and Youssef et al., in a cohort study evidenced that PIGF is a diagnostic marker in PE. PIGF serum levels were significantly lower in first trimester,who later developed PE. [11,12,13] In a cross-sectional study performed by El- Deen et al., with 74 pregnant women, and 50 PE Egyptian women examined for the maternal serum PIGF levels between severe and mild PE. They found there is no statistically significance between severe PE and mild PE groups. When compared with controls to PE women, the PIGF levels were significant which matches with the present study.[4]

The Indian studies on PIGF role or in combinations for prediction of PE availability has been very limited.[14] From India total of 3 studies (original studies) were performed, from north India (all from Delhi) 3 studies and from south India 1 study was done and present study is 2nd in southern part of India. From northern part of India, in Delhi by Agarwal et al., a study was performed on 17 PE cases, the levels of PIGF were compared between early and late PE but not statistically significant between early and late PE but when compared to PE and controls the levels were significant. The cut-off values in early PE were <30 pg/ml with 88.2% sensitivity and 71.4% and in late PE was <32 pg/ml with a sensitivity of 86.7% and specificity of 74.3%. [3]In a cohort study by Ghosh et al., from Delhi, measured maternal serum PIGF levels at 11–14 weeks and 22–24 weeks of gestation.So it proved that PIGF is an early second trimester marker as maternal serum PIGF levels are <144 pg/ml at 22–24 weeks of gestation were much higher than those of serum PIGF <228 pg/ml at 11–14 weeks of gestation.[15] In a case control study by Varughese et al., from Delhi with 80 study subjects measured the maternal sera PIGF levels. The mean PIGF levels were 236.77 pg/ml in PE where as in controls the mean PIGF levels are 744.98 pg/ml. [16]Another twin study by Agarwal et al., studied on PIGF and biophysical markers in late first trimester. They deliberate on 291 singleton pregnant women at 11–14 weeks of gestation who are primigravida of age group <40 years. PIGF alone had detection rate of 40% in PE.[14]The reduced levels of PIGF are evident from the first trimester, before 5 weeks the onset of PE. So, this is evident that PIGF levels will be lower from early gestational weeks itself especially in suspected PE. So, the maternal

serum PIGF levels, can be 90% and 50% accuracy are there in who developing early PE and late PE. There was a significant association between PIGF and the severity of PE also. [15] Present study results match with this study results. Chappell et al., in their study observed that PIGF has high sensitivity and predictive value for PE within 14 days in early onset suspected PE. So PIGF alone has a better predictive value. [18] Treatment of PE with PIGF is promising but with many uncertainties still remain.[2]Present study results also confirms that PIGF can be a diagnostic marker in late gestational weeks is significant and even in severe PE also.

V. Conclusion

In this present study PIGF is a PE diagnostic marker at late gestational weeks especially in south-eastern Kolar population. This is the 2nd study in southern part of India.