

A Study To Determine The Protein To Creatinine Ratio In Random Urine Sample To Predict The Protein Loss In 24 Hours Urine In Cases Of Chronic Kidney Disease In South Indian Population

Ms.Meghalatha.T.S^{1*}, Dr.N.Muninathan², Dr.Suma.H.R³ and Dr.R.Arunakumari⁴.

^{1,2}Department of Biochemistry, Meenakshi Medical College & Research Institute, Enathur, Kanchipuram, Tamilnadu, India.

³Department of Biochemistry, PES Institute of Medical Sciences & Research, Kuppam, Andhra Pradesh, India.

⁴Department of Biochemistry, Karpaga Vinayaga Institute of Medical Sciences, Chengalpattu, Tamilnadu, India.

Abstract: *Aim:* The aim of this study was to assess the ability of Protein to Creatinine ratio from random urine samples to rule in or rule out proteinuria in patients with Chronic Kidney Disease. Chronic renal failure is a clinical syndrome resulting in the progressive loss of renal function. The symptoms of chronic renal failure result not only from simple excretory failure but also from the onset of regulating failure, the kidney's failure to regulate certain substances, such as sodium & water. **Design/Methods:** The conducted study was a Correlation study. Chronic kidney disease patients were selected on the basis of serum creatinine levels. Serum creatinine >2.5mg/dl values were considered to be having chronic kidney disease. **Results:** The present study shows a very good correlation between 24hr urine protein loss and protein to creatinine ratio (PCR) in second void, third voids and early morning urines, and these correlations were very similar for each of the three urine aliquots. **Conclusion:** The tests are to be used for monitoring the level of proteinuria in established renal disease, they can then be used as surrogates for 24hr measurements

Key words: Creatinine, Proteinuria, Protein to Creatinine Ratio, Chronic kidney disease (CKD).

I. Introduction

Chronic renal failure is a clinical syndrome resulting in the progressive loss of renal function. The symptoms of chronic renal failure result not only from simple excretory failure but also from the onset of regulating failure, the kidney's failure to regulate certain substances, such as sodium & water.⁽¹⁾

According to the National Kidney Foundation –Kidney Dialysis Outcomes Quality Initiative (NKF-KDOQI) guidelines, chronic kidney disease is classified into 5 stages.⁽²⁾ The NKF-KDOQI guidelines stratify chronic kidney disease from stage 1 at the mild end of the spectrum to stage 5 which is severe.⁽³⁾ Stage 1&2 chronic kidney disease are usually not associated with any symptoms arising from the decrement in GFR. If the decline in GFR progresses to stages 3&4, clinical and laboratory complications of chronic kidney disease becomes more prominent. Virtually all organ systems are affected, but the most evident complications include anemia & associated easy fatigability. If the patient progresses to stage 5 chronic kidney disease, toxins accumulate such that patients usually experience a marked disturbance in their activities of daily living, wellbeing, nutritional status, and water & electrolyte homeostasis, culminating in the uremic syndrome. As discussed above, this state will end in death unless renal replacement therapy is instituted.

The dispiriting term end stage renal disease represents stage 5 of Chronic Kidney Disease where the accumulation of toxins, fluid and electrolytes results in the uremic syndrome.⁽²⁾ Early identification is essential so that adverse outcomes of Chronic Kidney Disease can be prevented or treated.⁽³⁾ The identification and quantification of Proteinuria is important in the initial diagnosis and subsequent follow-up and monitoring of renal disease. Proteinuria is also recognized as an independent risk factor for cardiovascular disease and as a predictor end organ damage. Urine protein or albumin measurements now form a central part of the classification and guidelines for Chronic Kidney Disease.

There is no 'gold standard' for identifying proteinuria and although it has been generally accepted that the best measure of protein loss is that based on a 24 h urine collection, the variability of results obtained by this method make this view questionable. Quoted reference ranges for 24 h protein loss vary, but most would regard a protein loss of more than 150 mg/24 h as abnormal with significant proteinuria being greater than 300 mg/ 24 h.

24hours urine collections, however, can be cumbersome to perform and prone to errors in collection. For this reason, the use of protein to creatinine ratios in random urine samples has been proposed. One

systematic review of studies using random urine Protein: Creatinine Ratio concluded that they provided evidence to rule out significant proteinuria as defined by a 24 hours measurement. ⁽⁴⁾

The aims of this study was to assess the ability of protein to creatinine ratio from random urine samples to rule in or rule out proteinuria in patients with Chronic Kidney Disease admitted in the Medicine department, PESIMSR, Kuppam. By collecting additional urine at different points during the day the most appropriate random urine sample, if any could be determined.

II. Materials And Methods

SUBJECTS

Chronic kidney disease patients were selected on the basis of serum creatinine levels. Serum creatinine >2.5mg/dl values were considered to be having chronic kidney disease. Both male and females patients with age ranging from 20-70yrs are taken in this study. The conducted study was a Correlation study. 60 samples were tested. Among that 36cases were males and 24 cases were females. Patients with cardiac disease, pregnant women and those who required urgent dialysis are excluded from this study. Patients with all stages of chronic kidney disease of any cause admitted in PESIMSR, Kuppam are included in this study.

III. Methodology and data collection

1. Collection of serum samples:-

- Venous blood was collected from the CKD patients admitted in Medicine department of PES Hospital, Kuppam.

2. Collection of Urine samples :-

- After discarding the first sample of the day 24hours urine sample of each patients is to be collected, in four separate containers 1) Second void, 2) Third void, 3) Remaining urine passed in that day &4) Early morning urine passed the following day.

Biochemical analysis

Serum creatinine, urinary creatininewas done manually by Jaffe’s method using calorimeter, and urinary protein measurement was done using a kit method which is basically calorimetric method.

The present study comprises of 60 clinically diagnosed cases of chronic kidney disease. The age group ranges from 20-70 years, out of these, 36 patients were males and 24 patients were females. Statistical data analysis was performed using “Excel”.

The main contributory causes for CKD in the 60 patients were diabetic nephropathy, & hypertensive damage. Causes in 17 patients were not certain or under investigation. There were 19 patients with diabetes mellitus(DM) and 24 patients with hypertension(HTN). Out of this 12 patients with DM and 15 patients with HTN were males, and 7patients with DM and 9 patients with HTN were females (Table.1).

Table.1 Causes of chronic kidney disease.

	N	Diabetes mellitus	Hypertension	Unknown
All patients	60	19	24	17
Males	36	12	15	09
females	24	07	09	08

57% were men and the median age for all patients was 51.5 ±14.2175. For all patients, median 24hr urinary protein loss was 2089.5 ± 580.192 mg/24hr and median urine volumes was 1.35± 0.43333 L/24hr (Table.2).

	n	Age (years)	Urine volume (L/24hr)	Urine protein (mg/24hr)	Creatinine clearance (ml/min)
All patients	60	51.5 ± 14.2175	1.35 ± 0.43333	2089.5 ± 580.192	10.4602 ± 0.53743
Male	36	52 ± 12.20064	1.34 ± 0.3975	2089.5 ± 485.7697	10.4775 ± 0.52218

Female	24	49.5 ± 12.6026	1.41 ± 0.4841	2077 ± 553.3312	10.387 ± 0.56858
--------	----	-------------------	------------------	-----------------	------------------

Protein: Creatinine Ratio Versus 24 H Urine Proteins

The present study shows a very good correlation between 24hr urine protein loss and protein to creatinine ratio (PCR) in second void, third voids and early morning urines, and these correlations were very similar for each of the three urine aliquots. PCR to be a good predictor of both abnormal urine protein loss and clinically significant urine protein loss.

Correlation between PCR in second void urine and 24hr urinary protein loss.

There is a good correlation between 24hr urinary protein loss and PCR in second void urine. The correlation coefficient between them is 0.50479. The graphical representation of this is shown in **diagram.1**




Diagram.1. Protein to creatinine ratio measured in second void urine compared with 24hr urine protein loss. Correlation coefficient were 0.50479.

Correlation between PCR in third void urine and 24hr urinary protein loss.

There is a good correlation between 24hr urinary protein loss and PCR in third void urine. The correlation coefficient between them is 0.5064. The graphical representation of this is shown in diagram.2




Diagram.2. Protein to creatinine ratio measured in third void urine compared with 24hr urine protein loss. Correlation coefficient were 0.5064.

Correlation between PCR in first void urine and 24hr urinary protein loss.

There is a good correlation between 24hr urinary protein loss and PCR in first void urine. The correlation coefficient between them is 0.69184. The graphical representation of this is shown in diagram.3.




Diagram.3. Protein to creatinine ratio measured in first void urine compared with 24hr urine protein loss. Correlation coefficient were 0.69184.

CRAETININE CLEARENCE

There is a negative correlation between 24hr urinary protein loss and creatinine clearance, it means that if the creatinine clearance decreases the protein loss increases. That is, the creatinine clearance and 24hr urinary protein are inversely proportional to each other. The correlation coefficient between is -0.2167. The graphical representation of this is shown in diagram.4.




Diagram.4. Creatinine clearance compared with 24hr urine protein loss. Correlation coefficient was-0.2167.

III. Discussion

Collection of 24hr urine samples is cumbersome and prone to errors, hence analyte measurement in random urine is often proposed with correction for variation in urine flow rate by expressing results as a ratio to creatinine concentration. It was not possible to formally assess completeness of collection of 24hr samples, particularly as there is little published data for CKD. Although samples were from a well-motivated group of patients who regularly made 24hr collections.

Several studies have assessed the benefits of using random urine PCR as a surrogate for 24hr protein measurement, but relatively few of these have involved patients with kidney disease. Most studies have shown good correlations between PCR and 24hr protein loss.^{(5), (6), (7), (8)}

The NKF K/DOQI guidelines suggests that untimed spot urine samples should be used to detect and monitor proteinuria in children and adults, it prefers a first-morning sample, but accepts a random sample if a first-morning specimen is not available. Few studies have previously assessed agreements rather than correlations between these tests and found wide limits. The limits of agreement were also wide, but similar across a wide range of protein excretion when data were log-transformed, the absolute difference between PCR and UP becomes very large as protein excretion increases. Urinary protein excretion is not constant and daily excretion varies by as much as 40% besides repeated 24-hour urine protein excretion varies by at least 15%. Rodby et al. repeated measurements on 33 patients at least three months apart and found discordant results were the PCR increased in some patients whereas the UP fell, and vice versa. Agrawal found a day to day variability in 24-hour urinary protein excretion of 10% and in protein-to-creatinine ratio of 2%. This variability is a likely reason for the poor agreement between the two methods of assessing proteinuria⁽⁹⁾. Several investigators studied the relationship between the protein:creatinineratio and 24-h protein excretion. Ginsberg et al. reporteda correlation coefficient of 0.972; these authors also studiedthe variation of this relationship during the course of 24 h by studying the ratio and absolute amount of protein excretedin urine samples from 46 patients collected over timed periodsthroughout the day. They found that the relationship variedby as much as 30% but that during normal daylight activity—whenmost random samples are likely to be collected—the variationwas minimal. The greatest differences were seen during the timeswhen the patients were most likely to be recumbent. These authorsconcluded on the basis of these data that the protein:creatinineratio of a spot urine could be used as a reliable indicatorof the 24-h protein excretion. Several investigators have makesimilar observations and drawn similar conclusions, whereasothers have stated a preference for the first sample collectedafter the first morning void.

However, some authorshave pointed out that regression analysis and the reportingof a correlation coefficient indicate the degree of linear associationbetween the two variables but do not enable a reliable decisionto be made to replace one with the other. Thus, the highdegree of association between the protein:

creatinine ratio and the 24-h protein excretion does not necessarily give reliable information on whether use of the ratio in a random sample will enable clinicians to reduce their dependence on the 24-h urine collection.⁽¹⁰⁾

For detection of proteinuria, the P/C ratio presented a high level of accuracy. Two previous studies used the P/C ratio cut-off values of 0.2 and 3.5 in patients with various nephropathies and stable renal function to establish the diagnosis of *pathologic proteinuria* ($P_{24} \geq 0.2$ g) and *nephrotic range proteinuria* ($P_{24} \geq 3.5$ g), respectively. In a study with pregnant women, Ramos *et al.* reported that the best cut-off to define normal protein excretion (<0.3 g) was a P/C ratio. Chitalia *et al.* described the P/C values of 0.26 and 3.2 as the best cut-off points to establish critical levels of proteinuria.⁽¹¹⁾

IV. Conclusion

This study has demonstrated good correlation between 24hr urine protein loss and PCR in second, third and early morning urines. But early morning urine PCR showed more correlation with 24hr urine protein loss.

Bibliography

- [1] Clinical chemistry (theory, analysis, correlation) by Lawrence A. Kaplan, Amadeo J. Pesce, Stanen C. Kajmierczak. 4th edition, page - 486.
- [2] Harrison's Internal Medicine Vol.2, 17th edition, page - 1762.
- [3] Tietz text book of clinical chemistry and Molecular diagnosis 4th edition page – 822, 1241-45, 1687-89, 1694.
- [4] Mark Guy, Joanna K Borzomato Ronald G Newall, Philip A Kalra and cheistopher P Price. Protein and albumin to creatinine ratios in random urines accurately predict 24 hours protein and albumin loss in patients with kidney disease. *Ann.ClinBiochem* 2009; 46:468/476.
- [5] Anne-Marie Côté, Mark A Brown, Elaine Lam, Peter von Dadelszen, Tabassum Firoz, Robert M Liston and Laura A Magee. Diagnostic accuracy of urinary spot protein: creatinine ratio for proteinuria in hypertensive pregnant women: systematic review. *BMJ*. 2008 May 3; 336(7651): 1003–1006.
- [6] Wahbeh AM, Ewais MH, Elsharif ME. Comparison of 24hour urinary protein and protein-to-creatinine ratio in the assessment of proteinuria. *Saudi J Kidney Dis Transpl*. 2009 May; 20 (3):443-7.
- [7] Morgenstern BZ, Butani L, Wollan P, Wilson DM, Larson TS. Validity of protein-osmolality versus protein-creatinine ratios in the estimation of quantitative proteinuria from random samples of urine in children. *Am J Kidney Dis*. 2003 Apr; 41(4):760-6.
- [8] Cathei Lane, Mark Brown, William Dunsmoir, John Kelly and George Mangos. Can spot urine protein/creatinine ratio replace 24 h urine protein in usual clinical nephrology? *Nephrology (Carlton)* 2006 Jun; 11(3):245-9.
- [9] Ayman M Wahbeh, Mohammad H Ewais, Mahamed E Elsharif. Comparison of 24-hour urinary protein and protein-to-creatinine ratio in the assessment of proteinuria. *Saudi J Kidney Dis Transpl Year* : 2009 | Volume : 20 | Issue : 3 | Page : 443-447
- [10] Christopher P. Price, Ronald G. Newall and James C. Boyd. Use of Protein:Creatinine Ratio Measurements on Random Urine Samples for Prediction of Significant Proteinuria: A Systematic Review *Clinical Chemistry* 51: 1577-1586, 2005.
- [11] Verônica Verleine Hörbe Antunes, Francisco José Veríssimo Veronese and José Vanildo Morales. Diagnostic accuracy of the protein/creatinine ratio in urine samples to estimate 24-h proteinuria in patients with primary glomerulopathies: a longitudinal study. *Nephrol. Dial. Transplant.* (2008) 23 (7): 2242-2246.