

# A New Geometric Average Technique to Solve Multi-Objective Linear Fractional Programming Problem and Comparison with New Arithmetic Average Technique

Samsun Nahar<sup>1</sup>, Md. Abdul Alim<sup>2</sup>

<sup>1</sup>Department of Basic Sciences and Humanities, University of Asia Pacific

<sup>2</sup>Department of Mathematics, Bangladesh University of Engineering and Technology  
Dhaka-1000, Bangladesh

---

**Abstract :** In this paper, we have suggested a new technique to solve a MOLFP by using new geometric averaging method. An algorithm is suggested for its solution. We have used some other techniques such as arithmetic averaging, geometric averaging, and new arithmetic averaging method to solve the same problem. New geometric averaging method gives better result than all those methods.

**Keywords:** Linear Fractional Programming Problem, Statistical average method, Arithmetic average, geometric average

---

## I. Introduction

Linear programming is comparatively a recent mathematical concept. A study of multi-objective linear programming problem (MOLPP) is introduced in [1] which suggests an approach to set up multi-objective function (MOF) under the limitation so that the optimum value of individual problem was greater than zero. Using mean and median the MOF was studied by solving multi-objective programming problem [2]. The multi-objective linear fractional programming problem (MOLFPP) was solved by the technique developed by Chandra Sen. The industrial production planning, financial and corporate planning, healthcare and hospital planning are important fields which use linear fraction maximum problems. So it has attracted considerable research and interest. There are several methods to solve these problems discussed in [3] where linear fractional programming is transformed to an equivalent linear program. Sing conducted a useful study about the optimality condition in fractional programming [4]. In [5], Sulaiman and Othman studied optimal transformation technique to solve MOLFP. In [6], Hamad Amin studied MOLPP using Arithmetic average. The study in [7] suggested a new technique to transform MOLPP to the single objective linear programming problem by using harmonic mean for values of functions.

In this paper I have defined a MOLFPP and suggested a new geometric average technique to optimize the objective function where a single objective function is developed from multi-objective functions. The result is compared with that of optimization using new arithmetic average technique. The results are also compared with those of optimization which are obtained by using arithmetic mean and geometric mean. This new geometric average technique gives better result than all those results.

## II. Problem Formulation

The main objective of this study is to solve multi-objective linear fractional programming problems. Before going to this problem, I would like to discuss some common definitions which will be used to understand the target of this paper.

1.1 Common Definitions:

1.1.1 Linear Programming:

Linear Programming deals with the optimization of a function of variables known as objective function, subject to set of linear equalities/inequalities known as constraints. The objective function may be profit, loss, cost, production capacity or any other measure of effectiveness which is to be obtained in the best possible or optimal manner. G.B. Dantzig in 1947 proposed the simplex algorithm as an efficient method to solve a linear programming problem [8].

$$\text{Max}(\text{min}) \quad c^T \bar{x}$$

A linear program is of the form:  $s/t \quad A\bar{x} = \bar{b}$

$$\bar{x} \geq 0$$



### III. MOLFPF Solution Techniques

We can solve MOLFPF using Chandra Sen's technique but it gives comparatively poor result of the objective function. But statistical average techniques (arithmetic and geometric) which are proposed in this paper for solving MOLFPF give better result of the objective function. These techniques are briefly described below.

#### 1.1 Solving MOLFPF by Chandra Sen's Technique:

We obtain a single value corresponding to each of the objective functions of MOLFPF of equation (1.1). They are being optimized individually subject to the constraints of equation (1.1) as follows:

$$\begin{aligned}
 &\text{Max } z_1 = \varphi_1 \\
 &\text{Max } z_2 = \varphi_2 \\
 &\dots \dots \dots \dots \dots \\
 &\text{Max } z_r = \varphi_r \\
 &\text{Min } z_{r+1} = \varphi_{r+1} \\
 &\dots \dots \dots \dots \dots \\
 &\text{Min } z_s = \varphi_s ;
 \end{aligned} \tag{1.2}$$

where  $\varphi_1, \varphi_2, \dots, \varphi_s$  are values of the objective functions. These values can be put in the equation (1.3), which is known as Chandra Sen's method, to find out a single objective function.

$$\max z = \sum_{i=1}^r \frac{z_i}{|\varphi_i|} - \sum_{i=r+1}^s \frac{z_i}{|\varphi_i|} \tag{1.3}$$

where,  $\varphi_i \neq 0, i = 1, 2, \dots, s$ . Subject to the constraints of equation (1.1) and the optimum value of the objective functions  $\varphi_i$  may be positive or negative.

#### 1.2 Proposed Technique

##### 1.2.1 Statistical averaging method:

$$\max z = \sum_{i=1}^r \frac{z_i}{A.M.(AA_i)} - \sum_{i=r+1}^s \frac{z_i}{A.M.(AL_i)} \tag{1.4}$$

$$\max z = \sum_{i=1}^r \frac{z_i}{G.M.(AA_i)} - \sum_{i=r+1}^s \frac{z_i}{G.M.(AL_i)} \tag{1.5}$$

Where,  $AA_i = |\varphi_i|, i = 1 \dots r$  and  $AL_i = |\varphi_i|, i = 1 + r \dots s$

##### 1.2.2 Solving MOLFPF by using the new arithmetic average technique:

Let  $m_1 = \min \langle AA_i \rangle$ , where  $AL_i = |\varphi_i|, \varphi_i$  is maximum value of  $z_i, i = 1 \dots r$

$m_2 = \min \langle AL_i \rangle$ , where  $AA_i = |\varphi_i|, \varphi_i$  is minimum value of  $z_i, i = r + 1 \dots s$

$A.Av = \frac{m_1 + m_2}{2}$  so

$$\max z = \left( \sum_{i=1}^r z_i - \sum_{i=r+1}^s z_i \right) / A.Av \tag{1.6}$$

##### 1.2.3 Solving MOLFPF by using the new geometric average technique:

Using  $m_1$  and  $m_2$  obtained in section 1.2.2, we can find the geometric average as follows:

$$G.Av = \sqrt{m_1 m_2}$$

$$\max z = \left( \sum_{i=1}^r z_i - \sum_{i=r+1}^s z_i \right) / G.Av \tag{1.7}$$

1.3 Algorithm for new Arithmetic and Geometric average technique:

- Step 1: Find the value of each of individual objective functions which is to be maximized or minimized.
- Step 2: Solve the first objective problem by simplex method.
- Step 3: Check the feasibility of the solution in step 2. If it is feasible then go to step 4. Otherwise, use dual simplex method to remove infeasibility.
- Step 4: Assign a name to the optimum value of the first objective function  $z_1$  say  $\phi_1$ .
- Step 5: Repeat the step 2,  $i=1, 2, \dots, s$
- Step 6: Select  $m_1 = \min\langle AA_i \rangle$ ,  $m_2 = \min\langle AL_i \rangle$ ,  $i = 1 \dots s$

$$A.Av = \frac{m_1 + m_2}{2} \quad \text{and} \quad G.Av = \sqrt{m_1 m_2}$$

Step 7: Optimize the combined objective function with the same constraints

$$\max \quad z = \left( \sum_{i=1}^r z_i - \sum_{i=r+1}^s z_i \right) / A.Av \quad \text{and} \quad \max \quad z = \left( \sum_{i=1}^r z_i - \sum_{i=r+1}^s z_i \right) / G.Av$$

1.4 Program solution for new Arithmetic and Geometric average technique:

The following program can be used to solve MOLFP by proposed method.  
For this, let

- $\phi A_i$  = value of objective functions which is to be maximized.
- $\phi L_i$  = value of objective functions which is to be minimized

So

$$AA_i = |\phi A_i|; \quad \forall i = 1 \dots r; \quad AL_i = |\phi L_i|; \quad \forall i = 1 + r \dots s$$

$$SM = \sum_{i=1}^r z_i; \quad SN = \sum_{i=r+1}^s z_i$$

$$m_1 = \min\langle AA_i \rangle; \quad m_2 = \min\langle AL_i \rangle$$

$$\max \quad z = (SM - SN) / A.Av \text{ and}$$

$$\max \quad z = (SM - SN) / G.Av$$

1.5 Flow Chart for new Arithmetic and Geometric average technique:

The flow chart, shown in Fig. 1.1, describes how the objective functions are optimized.

### IV. Test Calculations

Example: Consider the following multi-objective linear fractional programming problems. But as these functions are fractional, we have to use modified simplex method first to obtain the values of these objective functions. Then we can use these values to develop a single objective function using Chandra Sen's technique. Multi-objective fractional functions:

$$\begin{aligned}
 \max \quad z_1 &= \frac{3x_1 - 2x_2}{x_1 + x_2 + 1} \\
 \max \quad z_2 &= \frac{9x_1 + 3x_2}{x_1 + x_2 + 1} \\
 \max \quad z_3 &= \frac{3x_1 - 5x_2}{2x_1 + 2x_2 + 2} \\
 \min \quad z_4 &= \frac{-6x_1 + 2x_2}{2x_1 + 2x_2 + 2} \\
 \min \quad z_5 &= \frac{-3x_1 - x_2}{x_1 + x_2 + 1}
 \end{aligned}
 \quad
 \begin{aligned}
 &x_1 + x_2 \leq 2 \\
 \text{s/t} \quad &9x_1 + x_2 \leq 9 \\
 &x_1, x_2 \geq 0
 \end{aligned}
 \tag{1.8}$$




Fig. 1.1 Flow chart.

We have to solve by modified simplex method (equation (1.9)) [11] which implies

$$\begin{aligned} \max z = \frac{cx + \alpha}{dx + \beta} \quad & Ax(\leq, =, \geq)b \\ \text{s/t } x \geq 0 \quad & b \geq 0 \end{aligned} \tag{1.9}$$

The objective function  $H(y) = Iy + j$

Where,  $I = \frac{c\beta - d\alpha}{\beta}$ ,  $y = \frac{x}{dx + \beta}$ ,  $j = \frac{\alpha}{\beta}$ ;  $Ky \leq L$ ,  $K = A\beta + bd$ ,  $L = b$

Now from equation (1.8)

$$\begin{aligned} \max z_1 = \frac{3x_1 - 2x_2}{x_1 + x_2 + 1} \quad & x_1 + x_2 \leq 2 \\ \text{s/t } 9x_1 + x_2 \leq 9 \quad & x_1, x_2 \geq 0 \end{aligned}$$

$c = (3, -2)$ ,  $d = (1, 1)$ ,  $\alpha = 0$ ,  $\beta = -1$ ,  $A_1 = (1, 1)$ ,  $A_2 = (9, 1)$ ,  $b_1 = 2$ ,  $b_2 = 9$

$$\max H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] = (3, -2) \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = 3y_1 - 2y_2$$

For first constraint,

$$K_1 = (1, 1).1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3), Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1).1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10), Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$$

$$\begin{aligned} \max, H(y) = 3y_1 - 2y_2 \quad \max H = 3y_1 - 2y_2 \\ \text{Thus } \begin{aligned} s/t \ 3y_1 + 3y_2 \leq 2 &\Leftrightarrow s/t \ 3y_1 + 3y_2 + s_1 = 2 \\ 18y_1 + 10y_2 \leq 9 &\Leftrightarrow 18y_1 + 10y_2 + s_2 = 9 \\ y_1, y_2 \geq 0 &\quad \text{where } y_1, y_2, s_1, s_2 \geq 0 \end{aligned} \end{aligned}$$

Table I

c <sub>B</sub>	Basis	c <sub>j</sub>					
		3	-2	0	0		
		Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>	3	3	1	0	2	0.67
0	S <sub>2</sub>	18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>	3	-2	0	0	0	
		↑					
0	S <sub>1</sub>	0	4/3	1	-1/6	1/2	
3	Y <sub>1</sub>	1	5/9	0	1/18	1/2	
	C <sub>j</sub> -E <sub>j</sub>	0	-11/3	0	-1/6	3/2	

Thus  $y_1 = 1/2, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/2, 0).1}{1 - (1)(1/2, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus  $\max Z_1 = 3/2$  with  $x_1=1, x_2=0$

Second objective function in equation (1.8),

$$\begin{aligned} \max z_2 = \frac{9x_1 + 3x_2}{x_1 + x_2 + 1} \quad x_1 + x_2 \leq 2 \\ \text{s/t } 9x_1 + x_2 \leq 9 \\ x_1, x_2 \geq 0 \end{aligned}$$

So,  $c = (9, 3), d = (1, 1), \alpha = 0, \beta = 1, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$

$$\max H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (9, 3) \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = 9y_1 + 3y_2$$

For first constraint,

$$K_1 = (1, 1).1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3), Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1).1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10), Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$$

$$\begin{aligned} \max, H(y) &= 9y_1 + 3y_2 & \max H &= 9y_1 + 3y_2 \\ s/t \ 3y_1 + 3y_2 &\leq 2 & s/t \ 3y_1 + 3y_2 + s_1 &= 2 \\ 18y_1 + 10y_2 &\leq 9 & 18y_1 + 10y_2 + s_2 &= 9 \\ y_1, y_2 &\geq 0 & \text{where } y_1, y_2, s_1, s_2 &\geq 0 \end{aligned}$$

Table II

c <sub>B</sub>	Basis	c <sub>j</sub>					
		9	3	0	0		
		Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>	3	3	1	0	2	0.667
0	S <sub>2</sub>	18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>	9	3	0	0	0	
		↑					
0	S <sub>1</sub>	0	4/3	1	-1/6	1/2	
9	Y <sub>1</sub>	1	5/9	0	1/18	1/2	
	C <sub>i</sub> -E <sub>i</sub>	0	-2	0	-1/2	9/2	

Thus  $y_1 = 1/2, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/2, 0) \cdot 1}{1 - (1, 1)(1/2, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus  $\max Z_2 = 9/2$  with  $x_1=1, x_2=0$

For third objective function from equation (1.8)

$$\begin{aligned} \max z_3 &= \frac{3x_1 - 5x_2}{2x_1 + 2x_2 + 2} & x_1 + x_2 &\leq 2 \\ & & s/t \ 9x_1 + x_2 &\leq 9 \\ & & x_1, x_2 &\geq 0 \end{aligned}$$

So,  $c = (3, -5), d = (2, 2), \alpha = 0, \beta = 2, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$

$$\max H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (3, -5) \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = 3y_1 - 5y_2$$

For first constraint,

$$K_1 = (1, 1) \cdot 2 + 2(2, 2) = (2, 2) + (4, 4) = (6, 6),$$

$$Ky \leq L \Rightarrow (6, 6) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 6y_1 + 6y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1) \cdot 2 + 9(2, 2) = (18, 2) + (18, 18) = (36, 20)$$

$$Ky \leq L \Rightarrow (36, 20) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 36y_1 + 20y_2 \leq 9$$

$$\begin{aligned} \max, H(y) &= 3y_1 - 5y_2 & \max H &= 3y_1 - 5y_2 \\ s/t \ 6y_1 + 6y_2 &\leq 2 & s/t \ 6y_1 + 6y_2 + s_1 &= 2 \\ 36y_1 + 20y_2 &\leq 9 & 36y_1 + 20y_2 + s_2 &= 9 \\ y_1, y_2 &\geq 0 & \text{where } y_1, y_2, s_1, s_2 &\geq 0 \end{aligned}$$

Table III

c <sub>B</sub>	c <sub>j</sub> Basis	3	-5	0	0		
		Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>	6	6	1	0	2	0.33
0	S <sub>2</sub>	36	20	0	1	9	0.25
	C <sub>j</sub> -E <sub>j</sub>	3	-5	0	0	0	
		↑					
0	S <sub>1</sub>	0	8/3	1	-1/6	1/2	
3	Y <sub>1</sub>	1	5/9	0	1/36	1/4	
	C <sub>j</sub> -E <sub>j</sub>	0	-20/3	0	-1/2	3/4	

Thus  $y_1 = 1/4, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/4, 0) \cdot 2}{1 - (2, 2)(1/4, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus max  $Z_3 = 3/4$  with  $x_1=1, x_2=0$

For fourth objective function from equation (1.8)

$$\min z_4 = \frac{-6x_1 + 2x_2}{2x_1 + 2x_2 + 2} \quad \begin{matrix} x_1 + x_2 \leq 2 \\ \text{s/t } 9x_1 + x_2 \leq 9 \\ x_1, x_2 \geq 0 \end{matrix}$$

So,  $c = (-6, 2), d = (2, 2), \alpha = 0, \beta = 2, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$

$$\min H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (-6, 2) \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = -6y_1 + 2y_2$$

For first constraint,

$$K_1 = (1, 1) \cdot 2 + 2(2, 2) = (2, 2) + (4, 4) = (6, 6), Ky \leq L \Rightarrow (6, 6) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 6y_1 + 6y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1) \cdot 2 + 9(2, 2) = (18, 2) + (18, 18) = (36, 20)$$

$$Ky \leq L \Rightarrow (36, 20) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 36y_1 + 20y_2 \leq 9$$

$$\max H(y) = 6y_1 - 2y_2 \quad \max H = 6y_1 - 2y_2$$

$$\text{Thus } \begin{matrix} \text{s/t } 6y_1 + 6y_2 \leq 2 \\ 36y_1 + 20y_2 \leq 9 \\ y_1, y_2 \geq 0 \end{matrix} \Leftrightarrow \begin{matrix} \text{s/t } 6y_1 + 6y_2 + s_1 = 2 \\ 36y_1 + 20y_2 + s_2 = 9 \\ \text{where } y_1, y_2, s_1, s_2 \geq 0 \end{matrix}$$

Table IV

c <sub>B</sub>	c <sub>j</sub> Basis	6	-2	0	0		
		Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>	6	6	1	0	2	0.33
0	S <sub>2</sub>	36	20	0	1	9	0.25
	C <sub>j</sub> -E <sub>j</sub>	6	-2	0	0	0	
		↑					
0	S <sub>1</sub>	0	8/3	1	-1/6	1/2	
3	Y <sub>1</sub>	1	5/9	0	1/36	1/4	
	C <sub>j</sub> -E <sub>j</sub>	0	-16/3	0	-1/6	3/2	



Thus  $y_1 = 1/4, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1-d(y_1, y_2)} = \frac{(1/4, 0) \cdot 2}{1-(2, 2)(1/4, 0)} = \frac{(1/2, 0)}{1-1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus  $\max Z_4 = -3/2$  with  $x_1=1, x_2=0$

For fifth objective function from equation (1.8)

$$\min z_5 = \frac{-3x_1 - x_2}{x_1 + x_2 + 1} \quad \begin{matrix} x_1 + x_2 \leq 2 \\ \text{s/t } 9x_1 + x_2 \leq 9 \\ x_1, x_2 \geq 0 \end{matrix}$$

So,  $c = (-3, -1), d = (1, 1), \alpha = 0, \beta = 1, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$

$$\min H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (-3, -1) \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = -3y_1 - y_2$$

For first constraint,

$$K_1 = (1, 1) \cdot 1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3), Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1) \cdot 1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10), Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$$

$$\max, H(y) = 3y_1 + y_2 \quad \max H = 3y_1 + y_2$$

$$\text{Thus } \begin{matrix} \text{s/t } 3y_1 + 3y_2 \leq 2 \\ 18y_1 + 10y_2 \leq 9 \\ y_1, y_2 \geq 0 \end{matrix} \Leftrightarrow \begin{matrix} \text{s/t } 3y_1 + 3y_2 + s_1 = 2 \\ 18y_1 + 10y_2 + s_2 = 9 \\ \text{where } y_1, y_2, s_1, s_2 \geq 0 \end{matrix}$$

Table V

c <sub>B</sub>	Basis	c <sub>j</sub>	3	1	0	0		
			Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>		3	3	1	0	2	0.67
0	S <sub>2</sub>		18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>		3	1	0	0	0	
0	S <sub>1</sub>		0	4/3	1	-1/6	1/2	
3	Y <sub>1</sub>		1	5/9	0	1/18	1/2	
	C <sub>j</sub> -E <sub>j</sub>		0	-2/3	0	-1/6	3/2	

Thus  $y_1 = 1/2, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1-d(y_1, y_2)} = \frac{(1/2, 0) \cdot 1}{1-(1, 1)(1/2, 0)} = \frac{(1/2, 0)}{1-1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus  $\max Z_5 = -3/2$  with  $x_1=1, x_2=0$

The values obtained from the objective functions are summarized in Table VI.

Table VI

I	$\varphi_i$	$x_i$	$AA_i =  \varphi_i $	$AL_i =  \varphi_i $
1	3/2	(1, 0)	3/2	
2	9/2	(1, 0)	9/2	
3	3/4	(1, 0)	3/4	3/2
4	-3/2	(1, 0)		3/2
5	-3/2	(1, 0)		

Now, by Chandra Sen's approach:

$$\text{Max } Z = \sum_{k=1}^r \frac{z_k}{|\varphi_k|} - \sum_{k=r+1}^s \frac{z_k}{|\varphi_k|}$$

$$= \frac{2}{3} \frac{3x_1 - 2x_2}{x_1 + x_2 + 1} + \frac{2}{9} \frac{9x_1 + 3x_2}{x_1 + x_2 + 1} + \frac{4}{3} \frac{3x_1 - 5x_2}{2x_1 + 2x_2 + 2} - \left[ \frac{2}{3} \frac{-6x_1 + 2x_2}{2x_1 + 2x_2 + 2} + \frac{2}{3} \frac{-3x_1 - x_2}{x_1 + x_2 + 1} \right]$$

$$= \frac{1}{x_1 + x_2 + 1} [x_1(2 + 2 + 2) + x_2(2/3 - 10/3 - 4/3)] - \frac{1}{x_1 + x_2 + 1} [-4x_1] = \frac{10x_1 - 4x_2}{x_1 + x_2 + 1}$$

So the single objective function is

$$\max, z = \frac{10x_1 - 4x_2}{x_1 + x_2 + 1} \quad \begin{matrix} x_1 + x_2 \leq 2 \\ 9x_1 + x_2 \leq 9 \\ \text{s/t } x_1, x_2 \geq 0 \end{matrix} \tag{1.10}$$

Again the equation (1.10) is fractional, so we have to use modified simplex method to find its solution. So from modified simplex method

described in equation (1.9) we get

$$c = (10, 4), d = (1, 1), \alpha = 0, \beta = 1, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$$

$$\max H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (10, 4) \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = 10y_1 + 4y_2$$

For first constraint,

$$K_1 = (1, 1) \cdot 1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3)$$

$$Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1) \cdot 1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10)$$

$$Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$$

$$\max, H(y) = 10y_1 + 4y_2 \quad \max H = 10y_1 + 4y_2$$

$$\begin{matrix} \text{Thus} & \begin{matrix} \text{s/t } 3y_1 + 3y_2 \leq 2 \\ 18y_1 + 10y_2 \leq 9 \\ y_1, y_2 \geq 0 \end{matrix} & \Leftrightarrow & \begin{matrix} \text{s/t } 3y_1 + 3y_2 + s_1 = 2 \\ 18y_1 + 10y_2 + s_2 = 9 \\ \text{where } y_1, y_2, s_1, s_2 \geq 0 \end{matrix} \end{matrix}$$

Table VII

c <sub>B</sub>	Basis	c <sub>j</sub>					
		10	4	0	0		
0	S <sub>1</sub>	3	3	1	0	2	0.67
0	S <sub>2</sub>	18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>	10	4	0	0	0	
0	S <sub>1</sub>	0	4/3	1	-1/6	1/2	
10	Y <sub>1</sub>	1	5/9	0	1/18	1/2	
	C <sub>j</sub> -E <sub>j</sub>	0	-14/9	0	-5/9	5/2	

$$\text{Thus } y_1 = 1/2, \quad y_2 = 0$$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/2, 0) \cdot 1}{1 - (1, 1)(1/2, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus max Z = 5 with x<sub>1</sub>=1, x<sub>2</sub>=0

From Table VI, we get

$$A.M. (3/2, 9/2, 3/4) = 9/4, \quad A.M. (3/2, 3/2) = 3/2$$

So, by using equation (1.4) we have

$$\text{Max } Z = \sum_{i=1}^r \frac{z_i}{A.M(AA_i)} - \sum_{i=r+1}^s \frac{z_i}{A.M(AL_i)}$$

$$\begin{aligned} \text{Max } z &= \frac{4}{9} \left[ \frac{3x_1 - 2x_2}{x_1 + x_2 + 1} + \frac{9x_1 + 3x_2}{x_1 + x_2 + 1} + \frac{3x_1 - 5x_2}{2x_1 + 2x_2 + 2} \right] - \frac{2}{3} \left[ \frac{-6x_1 + 2x_2}{2x_1 + 2x_2 + 2} + \frac{-3x_1 - x_2}{x_1 + x_2 + 1} \right] \\ &= \frac{10x_1 - 2/3x_2}{x_1 + x_2 + 1} \end{aligned}$$

Thus  $\text{max}, z = \frac{10x_1 - 2/3x_2}{x_1 + x_2 + 1}$   $x_1 + x_2 \leq 2$   
 $9x_1 + x_2 \leq 9$   
 $s/t \ x_1, x_2 \geq 0$

$c = (10, -2/3), d = (1, 1), \alpha = 0, \beta = 1, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$

$\text{max } H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (10, -2/3) \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = 10y_1 - 2/3y_2$

For first constraint,

$K_1 = (1, 1).1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3), Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$

For second constraint,

$K_2 = (9, 1).1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10), Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$

$\text{max}, H(y) = 10y_1 - 2/3y_2$   $\text{max } H = 10y_1 - 2/3y_2$   
 Thus  $s/t \ 3y_1 + 3y_2 \leq 2$   $\Leftrightarrow \ s/t \ 3y_1 + 3y_2 + s_1 = 2$   
 $18y_1 + 10y_2 \leq 9$   $18y_1 + 10y_2 + s_2 = 9$   
 $y_1, y_2 \geq 0$   $where \ y_1, y_2, s_1, s_2 \geq 0$

Table VIII

c <sub>B</sub>	Basis	c <sub>j</sub>					
		10	-2/3	0	0		
		Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>	3	3	1	0	2	0.67
0	S <sub>2</sub>	18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>	10	-2/3	0	0	0	
0	S <sub>1</sub>	0	4/3	1	-1/6	1/2	
10	Y <sub>1</sub>	1	5/9	0	1/18	1/2	
	C <sub>j</sub> -E <sub>j</sub>	0	-56/9	0	-5/9	-5/9	

Thus  $y_1 = 1/2, \ y_2 = 0$

Now  $(x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/2, 0).1}{1 - (1, 1)(1/2, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$

Thus  $\text{max } Z = 5$  with  $x_1=1, x_2=0$

From Table VI, we get  $G.M. (3/2, 9/2, 3/4) = 1.7171; G.M. (3/2, 3/2) = 1.5$

So, by using equation (1.5) we have

$\text{Max } Z = \sum_{i=1}^r \frac{z_i}{G.M(AA_i)} - \sum_{i=r+1}^s \frac{z_i}{G.M(AL_i)}$

$$\begin{aligned} \text{Max } z &= \frac{1}{1.7171} \left[ \frac{3x_1 - 2x_2}{x_1 + x_2 + 1} + \frac{9x_1 + 3x_2}{x_1 + x_2 + 1} + \frac{3x_1 - 5x_2}{2x_1 + 2x_2 + 2} \right] - \frac{1}{1.5} \left[ \frac{-6x_1 + 2x_2}{2x_1 + 2x_2 + 2} + \frac{-3x_1 - x_2}{x_1 + x_2 + 1} \right] \\ &= \frac{11.862x_1 - 0.8736x_2}{x_1 + x_2 + 1} \end{aligned}$$

$$\text{Thus } \max Z = \frac{11.862x_1 - 0.8736x_2}{x_1 + x_2 + 1} \quad \begin{array}{l} x_1 + x_2 \leq 2 \\ 9x_1 + x_2 \leq 9 \\ \text{s/t } x_1, x_2 \geq 0 \end{array}$$

$$c = (11.862, -0.8736), d = (1, 1), \alpha = 0, \beta = 1, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$$

$$\max H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (11.862, -0.8736) \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = 11.862y_1 - 0.8736y_2$$

For first constraint,

$$K_1 = (1, 1) \cdot 1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3), Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$$

For second constraint

$$K_2 = (9, 1) \cdot 1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10), Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$$

$$\begin{array}{l} \max, H(y) = 11.862y_1 - 0.8736y_2 \quad \max H = 11.862y_1 - 0.8736y_2 \\ \text{s/t } 3y_1 + 3y_2 \leq 2 \quad \Leftrightarrow \quad \text{s/t } 3y_1 + 3y_2 + s_1 = 2 \\ 18y_1 + 10y_2 \leq 9 \quad \Leftrightarrow \quad 18y_1 + 10y_2 + s_2 = 9 \\ y_1, y_2 \geq 0 \quad \text{where } y_1, y_2, s_1, s_2 \geq 0 \end{array}$$

Table IX

c <sub>B</sub>	Basis	c <sub>j</sub>					
		11.862	-0.8736	0	0		
		Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>	3	3	1	0	2	0.67
0	S <sub>2</sub>	18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>	11.862	-0.8736	0	0	0	
		↑					
0	S <sub>1</sub>	0	4/3	1	-1/6	1/2	
11.862	Y <sub>1</sub>	1	5/9	0	1/18	1/2	
	C <sub>j</sub> -E <sub>j</sub>	0	-7.4636	0	-0.659	-5.931	

Thus  $y_1 = 1/2, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/2, 0) \cdot 1}{1 - (1, 1)(1/2, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus  $\max Z = 5.931$  with  $x_1=1, x_2=0$

**New Arithmetic average technique:**

$$\text{Let } m_1=0.75, m_2=1.5; \frac{m_1 + m_2}{2} = \frac{0.75 + 1.5}{2} = 1.125$$

$$\max z = \left( \sum_{i=1}^r z_i - \sum_{i=r+1}^s z_i \right) / A.Av = \frac{39x_1 - 3x_2}{2(1.125)(x_1 + x_2 + 1)} = \frac{39x_1 - 3x_2}{2.25 \cdot (x_1 + x_2 + 1)} = \frac{17.33x_1 - 1.33x_2}{x_1 + x_2 + 1}$$

$$\text{Thus } \max Z = \frac{17.33x_1 - 1.33x_2}{x_1 + x_2 + 1} \quad \begin{array}{l} x_1 + x_2 \leq 2 \\ 9x_1 + x_2 \leq 9 \\ \text{s/t } x_1, x_2 \geq 0 \end{array}$$

$$c = (17.33, -1.33), d = (1, 1), \alpha = 0, \beta = 1, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$$

$$\max H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (17.33, -1.33) \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = 17.33y_1 - 1.33y_2$$

For first constraint,

$$K_1 = (1, 1).1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3), Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1).1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10), Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$$

$$\begin{aligned} \max, H(y) &= 17.33y_1 - 1.33y_2 & \max H &= 17.33y_1 - 1.33y_2 \\ \text{Thus } s/t \quad & 3y_1 + 3y_2 \leq 2 & \Leftrightarrow & s/t \quad 3y_1 + 3y_2 + s_1 = 2 \\ & 18y_1 + 10y_2 \leq 9 & & 18y_1 + 10y_2 + s_2 = 9 \\ & y_1, y_2 \geq 0 & \text{where } & y_1, y_2, s_1, s_2 \geq 0 \end{aligned}$$

Table X

c <sub>B</sub>	Basis	c <sub>j</sub>		S <sub>1</sub>	S <sub>2</sub>		
		17.33	-1.33				
		Y <sub>1</sub>	Y <sub>2</sub>				
0	S <sub>1</sub>	3	3	1	0	2	0.67
0	S <sub>2</sub>	18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>	17.33	-1.33	0	0	0	
0	S <sub>1</sub>	0	4/3	1	-1/6	1/2	
17.33	Y <sub>1</sub>	1	5/9	0	1/18	1/2	
	C <sub>j</sub> -E <sub>i</sub>	0	-10.96	0	-0.963		

Thus  $y_1 = 1/2, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/2, 0).1}{1 - (1, 1)(1/2, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus max Z = 8.665 with  $x_1=1, x_2=0$

**New Geometric average technique:**

Let  $m_1=0.75, m_2=1.5$ ; So  $G.Av = \sqrt{0.75(1.5)} = 1.061$

$$\begin{aligned} \max z &= \left( \sum_{i=1}^r z_i - \sum_{i=r+1}^s z_i \right) / G.Av \\ &= \frac{39x_1 - 3x_2}{2(1.061)(x_1 + x_2 + 1)} = \frac{39x_1 - 3x_2}{2.122.(x_1 + x_2 + 1)} = \frac{18.379x_1 - 1.4137x_2}{x_1 + x_2 + 1} \end{aligned}$$

$$\begin{aligned} \text{Thus } \max z &= \frac{18.379x_1 - 1.4137x_2}{x_1 + x_2 + 1} \\ & \quad x_1 + x_2 \leq 2 \\ & \quad 9x_1 + x_2 \leq 9 \\ & \quad s/t \quad x_1, x_2 \geq 0 \end{aligned}$$

$c = (18.379, -1.4137), d = (1, 1), \alpha = 0, \beta = 1, A_1 = (1, 1), A_2 = (9, 1), b_1 = 2, b_2 = 9$

$$\max H(y) = [Iy] + j = \frac{c\beta - d\alpha}{\beta} [y] + 0 = (18.379, -1.4137) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = 18.379y_1 - 1.4137y_2$$

For first constraint,

$$K_1 = (1, 1).1 + 2(1, 1) = (1, 1) + (2, 2) = (3, 3), Ky \leq L \Rightarrow (3, 3) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 2 \Rightarrow 3y_1 + 3y_2 \leq 2$$

For second constraint,

$$K_2 = (9, 1).1 + 9(1, 1) = (9, 1) + (9, 9) = (18, 10), Ky \leq L \Rightarrow (18, 10) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \leq 9 \Rightarrow 18y_1 + 10y_2 \leq 9$$

$$\begin{aligned} \max, H(y) &= 18.379y_1 - 1.4137y_2 & \max H &= 18.379y_1 - 1.4137y_2 \\ \text{Thus } s/t \ 3y_1 + 3y_2 &\leq 2 & \Leftrightarrow \ s/t \ 3y_1 + 3y_2 + s_1 &= 2 \\ 18y_1 + 10y_2 &\leq 9 & 18y_1 + 10y_2 + s_2 &= 9 \\ y_1, y_2 &\geq 0 & \text{where } y_1, y_2, s_1, s_2 &\geq 0 \end{aligned}$$

Table XI

c <sub>B</sub>	c <sub>j</sub> Basis	18.379	-1.4137	0	0		
		Y <sub>1</sub>	Y <sub>2</sub>	S <sub>1</sub>	S <sub>2</sub>		
0	S <sub>1</sub>	3	3	1	0	2	0.67
0	S <sub>2</sub>	18	10	0	1	9	0.5
	C <sub>j</sub> -E <sub>j</sub>	18.379	-1.4137	0	0	0	
0	S <sub>1</sub>	0	4/3	1	-1/6	1/2	
18.379	Y <sub>1</sub>	1	5/9	0	1/18	1/2	
	C <sub>j</sub> -E <sub>1</sub>	0	-11.624	0	-1.021		

Thus  $y_1 = 1/2, y_2 = 0$

$$\text{Now } (x_1, x_2) = \frac{(y_1, y_2)\beta}{1 - d(y_1, y_2)} = \frac{(1/2, 0) \cdot 1}{1 - (1, 1)(1/2, 0)} = \frac{(1/2, 0)}{1 - 1/2} = \frac{(1/2, 0)}{1/2} = (1, 0)$$

Thus max Z = 9.1895 with  $x_1=1, x_2=0$

The solutions of the MOLFPF solved by different approaches are summarized in Table XII which shows that we get the improved value of the objective function using new statistical average method proposed in this paper. So the proposed technique is justified here.

Table XII

Chandra Sen's Approach	Statistical Average Method		New Statistical Average Method	
	Using A.M	Using G.M	New A. Av method	New G. Av method
Max Z=5 with $x_1=1, x_2=0$	Max Z=5 with $x_1=1, x_2=0$	Max Z=5.931 with $x_1=1, x_2=0$	Max Z=8.665 with $x_1=1, x_2=0$	Max Z=9.1895 with $x_1=1, x_2=0$

**V. Conclusion**

In this paper, a MOLFPF has been solved using different methods such as Chandra Sen's approach and proposed statistical average method, and the results are compared in Table XII. In statistical average method, we proposed geometric and arithmetic average approach. We also proposed a new statistical average approach to solve the problem. It is observed that statistical average method results better optimization than Chandra Sen's approach of the MOLFPF. The proposed new statistical average method optimizes the problem better than that of statistical average method. We also found that geometric average technique is suited for optimizing MOLFPF better than that of arithmetic average technique.

**References**

- [1]. Sen, Chandra., "A new approach for multi objective rural development planning", The Indian Economic Journal, Vol.30, pp.91-96.
- [2] Azapagic, A., R. Clift., (1999), "Life cycle assessment and multi objective optimization", Journal of cleaner production, Vol.7, pp.135-145, 1983
- [2]. Sulaiman, N.A. and Sadiq, G.W., "Solving the linear multi-objective programming problems; using mean and median value", Al-Rafiden Journal of Computer Sciences and Mathematics, University of Mosul, Vol.3, No. 1, pp.69-83, 2006
- [3]. A. Abdil-Kadir, M.S and Sulaiman, and N.A., "An approach for multi objective fractional programming problem", Journal of the College of Education, Univ. Salahaddin, Vol. 3. pp.1-5, 1993
- [4]. Sing, H.C., "Optimality condition in functional programming", Journal of Optimization Theory and Application, Vol. 33, pp.287-294, 1981
- [5]. Sulaiman, N.A. and Othman, A.Q., "Optimal transformation Technique to solve multi-objective linear programming problem", Journal of University of Kirkuk, Vol.2. No. 2, 2007
- [6]. Hamad-Amin A.O., "An adaptive Arithmetic Average Transformation Technique for solving MOLPP", M.Sc. Thesis, University of Koya, Koya/Iraq, 2008
- [7]. Nejmaddin A. Sulaiman, Rebaz B. Mustafa, " Using harmonic mean to solve multi-objective linear programming problems", American Journal of operations Research, 2016, 6, 25-30.
- [8]. Wikipedia, linear programming, [http://en.wikipedia.org/wiki/Linear\\_Programming](http://en.wikipedia.org/wiki/Linear_Programming)
- [9]. S. Sharma, Operation Research, Kedar Nath Ram Nath BCO. Meerut, India. (1988), 60, 559.
- [10]. Operation Research, [http://www.scribd.com/doc/100903589/Operation\\_Research](http://www.scribd.com/doc/100903589/Operation_Research), (21)
- [11]. Saha, S.K., Hossain, M.R., Uddin, M.K. and Mondal, R.N. 'A New Approach of Solving Linear Fractional Programming Problem (LFP) by Using Computer Algorithm' Open Journal of Optimization, 4, 74-86, 2015 <http://dx.doi.org/10.4236/ojop.2015.43010>