

# SDC Labeling For Cycle Of Join Zero Divisor Graphs

Dr. D. Bharathi<sup>1</sup>, M. Lakshmi<sup>2</sup>

Professor, Research Scholar, Department Of Mathematics, S. V. University, Tirupati- 517502, A. P, India.

## Abstract:

A sum divisor cordial labeling of a graph  $G$  with vertex set  $V(G)$  is a bijection  $f$  from  $V(G)$  to  $\{1, 2, 3, \dots, |V(G)|\}$  such that an edge  $uv$  is assigned the label 0 if 2 divides  $f(u)+f(v)$  and the label 1 otherwise, if it satisfies the condition

$|e_f(0) - e_f(1)| \leq 1$ . Then the graph with a sum divisor cordial labeling is called a sum divisor cordial graph. In this paper we introduce the concept of the sum divisor cordial labeling for cycle of join zero divisor graphs.

**Keywords:** Cycle, Zero divisor graphs, cordial graphs, cycle of graphs, Sum divisor cordial labeling, Sum divisor cordial graphs.

Date of Submission: 25-02-2025

Date of Acceptance: 05-03-2025

## I. Introduction

Let  $G = (V(G), E(G))$  be a simple, finite and undirected graph. Different types of graphs labeling techniques have been investigated and over to 2000 papers have been published in this area. A graph labeling is an assignment of labels traditionally represented by integers to edges or vertices or both. Beck [6] introduced Zero-divisor graph. Then Anderson and Livingston [7] modified it. Tamizh Chelvam et al [13] proposed graph labeling related to Zero-divisors in a commutative ring. obtained certain labeling for certain class of Zero-divisor graphs corresponding to finite rings. Let  $R$  be a commutative ring with non-zero identity,  $Z(R)$  it's set of all zero divisors in  $R$  and

$Z^*(R) = Z(R) \setminus \{0\}$ . the Zero-divisor graph of  $R$  is the simple undirected graph  $\Gamma(R)$  with vertex set  $Z^*(R)$  and two distinct vertices  $x$  and  $y$  are adjacent if  $xy=0$ . [11] S. Sajana and D. Bharathi studied the Intersection graph of zero-divisors of a finite commutative ring. [12] D. Eswara Rao and D. Bharathi studied the Total zero divisor graph of a commutative ring. All graphs consider in this paper are finite, simple and undirected. This graph product was introduced by Frucht and Harary [3]. After that A. Lourdusamy [10] introduced sum divisor cordial labeling of graphs. Here, we are interested in the sum divisor cordial (SDC) labeling for cycle of join zero divisor graphs. Based on these concepts we give several results on SDC labeling for Zero-divisor graphs. Here we study the cycle of join Zero-divisor graphs.

## II. Preliminaries:

### Definition 2.1

For a commutative ring  $Z_n$  with unity ( $1 \neq 0$ ), the zero-divisor graph of  $Z_n$ , denoted by  $\Gamma(Z_n)$ , is a simple graph with vertices as elements of  $Z_n$  and two distinct vertices are adjacent whenever the product of the vertices is zero.

### Definition 2.2

Let  $R$  be a finite ring. An element  $a \in R$  is called a zero divisor if there exists a non-zero element  $b \in R$  such that  $ab = 0$  or  $ba = 0$ .

### Definition 2.3

The join of two graphs  $G_1$  and  $G_2$  is denoted by  $G_1 + G_2$  and whose vertex set is  $V(G_1 + G_2) = V(G_1) \cup V(G_2)$  and edge set is  $E(G_1 + G_2) = E(G_1) \cup E(G_2) \cup \{uv : u \in V(G_1), v \in V(G_2)\}$ .

### Definition 2.4

A Graph  $G$  is said to be complete if every pair of its distinct vertices are adjacent.

### Definition 2.5

A bipartite graph is a graph whose vertex set  $V(G)$  can be partitioned into two subsets  $V_1$  and  $V_2$  such that every edge of  $G$  has one end in  $V_1$  and the other end in  $V_2$ .  $(V_1, V_2)$  is called a bipartition of  $G$ .

**Definition 2.6**

A complete bipartite graph is a bipartite graph with bipartition  $(V_1, V_2)$  such that every vertex of  $V_1$  is joined to all the vertices of  $V_2$ . It is denoted by  $K_{m,n}$ , where  $|V_1| = m$  and  $|V_2| = n$ .

A star graph is a complete bipartite graph  $K_{1,n}$ .

**Definition 2.7**

The complement of the graph  $G$  is the graph with vertex set  $V(G)$  and two vertices are adjacent in it and only if they are not adjacent in  $G$ .

**Definition 2.8**

For a cycle  $C_n$ , each vertex of  $C_n$  is replaced by connected graphs  $G_1, G_2, \dots, G_n$  and is known as cycle of graphs. We shall denote it by  $C(G_1, G_2, \dots, G_n)$ . If we replace each vertex by a graph  $G$ , i.e.  $G_1 = G = G_2 = \dots = G_n$ , such cycle of a graph  $G$  is denoted by  $C(n, G)$ .

**Definition 2.9**

A cycle graph is a graph that contains a single cycle, or a closed chain of vertices connected by edges. A cycle graph with  $n$  vertices is denoted by  $C_n$ .

**Definition 2.10**

A star graph is a Tree with a single internal node and multiple leaves.

**Definition 2.11**

Cordial labeling is a way to assign labels to vertices and edges using only the numbers 0 and 1. The labels are assigned so that the number of vertices and edges labeled 0 and 1 differ by at most 1.

**Definition 2.12**

Let  $G = (V(G), E(G))$  be a simple graph and  $f: V(G) \rightarrow \{1, 2, \dots, |V(G)|\}$  be the bijection. For each edge  $uv$ , assign the label 0, if  $2/f(u)+f(v)$  and the label 1 otherwise. The function  $f$  is a sum divisor cordial (SDC) labeling if it satisfies the condition  $|e_f(0) - e_f(1)| \leq 1$ , where  $e_f(0)$  is the number of edges labeled with 0 and  $e_f(1)$  is the number of edges labeled with 1. Then a graph which admits sum divisor cordial (SDC) labeling is called as a sum divisor cordial (SDC) Graph.

**III. Main Results**

In this section, we obtain cycle of join zero-divisor graphs which admits a SDC labeling.

**Theorem 3.1.** For any prime number  $p > 2$ , cycle of join zero-divisor graphs  $\Gamma(Z_{2p}) + \Gamma(Z_4)$  where  $n \geq 3$  is a SDC graph.

Proof: Let  $G = C(n, (\Gamma(Z_{2p}) + \Gamma(Z_4)))$ ,  $\forall n \geq 3$

Let the vertex set and edge set of the graph  $G$  are

$$V(G) = \{u_{1,1}, u_{1,2}, \dots, u_{1,p}, x_1, u_{2,1}, u_{2,2}, \dots, u_{2,p}, x_2, \dots, u_{n,1}, u_{n,2}, \dots, u_{n,p}, f\}, \forall n \geq 3$$

$$E(G) = \{u_{j,i}u_{j,p}, u_{j,i}x_j, u_{j,p}x_j, u_{1,p}u_{2,p}, u_{2,1}u_{3,1}, \dots, u_{n,p}u_{1,1}\}, \forall n \geq 3$$

Therefore  $|V(G)| = n(p+1)$  and

$$|E(G)| = n(2p-1) + n = 2np$$

Define  $f: V(G) \rightarrow \{1, 2, 3, \dots, n(p+1)\}$ ,  $\forall n \geq 3$  by

$$f(u_{1,p}) = 1, f(u_{2,p}) = p+2, f(u_{3,p}) = 2p+3, \dots, f(u_{n,p}) = (n-1)p + n, \forall n \geq 3$$

$$f(u_{1,i}) = i+1 \text{ for } ,$$

$$f(u_{2,i}) = p+2+i \text{ for}$$

.

.

.

$$f(u_{n,i}) = (n-1)p + n + i \text{ for}$$

$$f(x_1) = p+1, f(x_2) = 2p+2, \dots, f(x_n) = np + n.$$

Then the induced edge labeling is as follows  $f: E(G) \rightarrow \{0, 1\}$  is given by

$$f(u_{j,i}u_{j,p}) =$$

$$f(u_{j,i}x_j) =$$

$$f(u_{j,p}x_j) = 1, f(u_{j,p}u_{j+1,p}) = 0, f(u_{n,p}u_{1,p}) = 0.$$

we note that

$$(0) = n(p-1) + n = np \text{ and } (1) = n(p)$$

$$\text{then } |(0) - (1)|$$

Hence G is a SDC graph. □

**Example 3.2:**

A SDC labeling for  $G = C(4, (\Gamma(Z_{2p}) + \Gamma(Z_4)))$ , is given in figure 1 where  $n=4, p=3$

$$V(G) = \{u_{1,1}, u_{1,2}, u_{1,3}, x_1, u_{2,1}, u_{2,2}, u_{2,3}, x_2, u_{3,1}, u_{3,2}, u_{3,3}, x_3, u_{4,1}, u_{4,2}, u_{4,3}\}, \forall n \geq 3$$

$$E(G) = \{u_{j,i}u_{j,3}, u_{j,i}x_j, u_{j,3}x_j, u_{1,3}u_{2,3}, u_{2,p}u_{3,3}, \dots, u_{4,p}u_{1,3}\},$$

Therefore,  $|V(G)|=16$  and  $|E(G)|=24$

Here  $(0) = 12$  and  $(1) = 12$

Hence  $|(0) - (1)|$  then  $C(4, (\Gamma(Z_{2p}) + \Gamma(Z_4)))$ , is a SDC graph.




Figure 1.  $C(4, (\Gamma(Z_{2p}) + \Gamma(Z_4)))$

**Theorem 3.3.** For any prime number  $p > 2$ , cycle of join zero-divisor graphs  $\Gamma(Z_{2p}) + \Gamma(Z_6)$  where  $n \geq 3$  is a SDC graph.

Proof: Let  $G = C(5, (\Gamma(Z_{2p}) + \Gamma(Z_6)))$

Let the vertex set and edge set of the graph G are

$$V(G) = \{u_{1,1}, u_{1,2}, \dots, u_{1,p}, x_1, y_1, z_1, u_{2,1}, u_{2,2}, \dots, u_{2,p}, x_2, y_2, z_2, u_{n,1}, u_{n,2}, \dots, u_{n,p}, x_n, y_n, z_n\}$$

$$\text{and } E(G) = \{u_{j,i}u_{j,p}, u_{j,i}x_j, u_{j,p}x_j, u_{j,i}y_j, u_{j,p}y_j, u_{j,p}z_j, x_jy_j, y_jz_j, u_{1,p}u_{2,p}, u_{2,p}u_{3,p}, \dots, u_{n,p}u_{1,p}\}$$

Therefore  $|V(G)| = n(p+3)$  and  $|E(G)| = n(4p+1) + n = 4np + 2n = 2n(2p+1)$

Define the vertex labeling

$$f: V(G) \rightarrow \{1, 2, 3, \dots, n(p+3)\} \text{ by}$$

$$f(u_{i,i}) = i+1 \text{ for ,}$$

$$f(u_{2,i}) = p+4+i \text{ for}$$

$$f(u_{3,i}) = 2p+7+i \text{ for}$$

·  
·  
·

$$f(u_{n,i}) = (n-1)p + (3n-2) + i \text{ for } ,$$

$$f(u_{1,p}) = 1, f(u_{2,p}) = p+4, f(u_{3,p}) = 2p+7, \dots, f(u_{n,p}) = (n-1)p + (3n-2),$$

$$f(x_1) = p+1, f(x_2) = 2p+4, f(x_3) = 3p+7, \dots, f(x_n) = np+3n-2,$$

$$f(y_1) = p+3, f(y_2) = 2p+6, f(y_3) = 3p+9, \dots, f(y_n) = np+3n,$$

$$f(z_1) = p+2, f(z_2) = 2p+5, f(z_3) = 3p+8, \dots, f(z_n) = np+3n-1.$$

Then the induced edge labeling is as follows  $f^*: E(G) \rightarrow \{0, 1\}$  is given by

$$f^*(u_{j,i}u_{j,p}) =$$

$$f^*(u_{j,i}x_j) =$$

$$f^*(u_{j,i}y_j) =$$

$$f^*(u_{j,i}z_j) =$$

$$f^*(u_{j,p}x_j) = 1, f^*(u_{j,p}y_j) = 1, f^*(u_{j,p}z_j) = 0, f^*(x_jy_j) = 0, f^*(y_jz_j) = 1, f^*(u_{j,p}u_{j+1,p}) = 0,$$

$$f^*(u_{n-1,p}u_{1,p}) = 0$$

we observe that  $(0) = 2np+n = (2p+1)n$  and  $(1) = n(2p+1)$  then  $|(0) - (1)|$

Hence  $G$  is a SDC graph. □

**Example 3.4.**

A SDC labeling of the  $C(5, \Gamma(Z_6) + \Gamma(Z_6))$  is given in figure 2 where  $n=5, p=3$ .

Let  $G = C(5, \Gamma(Z_6) + \Gamma(Z_6))$

$V(G) = \{u_{1,1}, u_{1,2}, u_{1,3}, x_1, y_1, z_1, u_{2,1}, u_{2,2}, u_{2,3}, x_2, y_2, z_2, u_{3,1}, u_{3,2}, u_{3,3}, x_3, y_3, z_3, u_{4,1}, u_{4,2}, u_{4,3}, x_4, y_4, z_4, u_{5,1}, u_{5,2}, u_{5,3}, x_5, y_5, z_5\}$ , and

$E(G) = \{u_{j,i}u_{j,3}, u_{j,i}x_j, u_{j,3}x_j, u_{j,i}y_j, u_{j,3}y_j, u_{j,3}z_j, x_jy_j, y_jz_j, u_{1,3}u_{2,3}, u_{2,3}u_{3,3}, \dots, u_{5,3}u_{1,1}\}$

Therefore,  $|V(G)| = 30$  and  $|E(G)| = 70$

Here  $(0) = 35$  and  $(1) = 35$

Hence  $|(0) - (1)|$  then  $C(5, (\Gamma(Z_6) + \Gamma(Z_6)))$  is a SDC graph.




Figure 2.  $C(5, \Gamma(Z_6) + \Gamma(Z_6))$

**Theorem 3.5.** For any prime number  $p > 2$ , cycle of join zero-divisor graphs  $\Gamma(Z_{2p}) + \Gamma(Z_9)$

where  $n \geq 3$  does not admit SDC labeling.

□

**Example 3.6:**

A SDC labeling of the  $C(4, (\Gamma(Z_6) + \Gamma(Z_9)))$  is given in figure 3 where  $n=4, p=3$ .

Let  $G = C(4, (\Gamma(Z_6) + \Gamma(Z_9)))$


$V(G) = \{u_{1,1}, u_{1,2}, u_{1,3}, x_1, y_1, u_{2,1}, u_{2,2}, u_{2,3}, x_2, y_2, u_{3,1}, u_{3,2}, u_{3,3}, x_3, y_3, u_{4,1}, u_{4,2}, u_{4,3}, x_4, y_4\}$  and

$E(G) = \{u_{i,1}u_{i,3}, u_{i,1}x_j, u_{i,3}x_j, u_{i,1}y_i, u_{i,3}y_i, x_iy_j, u_{1,3}u_{2,3}, u_{2,3}u_{3,3}, \dots, u_{4,3}u_{11}\}$

Therefore,  $|V(G)|=20$  and  $|E(G)|=40$

Here  $|0|=16$  and  $|1|=24$

Hence  $|0| - |1|$  then  $C(4, (\Gamma(Z_6) + \Gamma(Z_9)))$  is not a SDC graph.



**Figure 3.**  $C(4, (\Gamma(Z_6) + \Gamma(Z_9)))$

**Theorem 3.7.** For any prime number  $p > 2$ , cycle of  $\Gamma()$  does not admit the SDC labeling.

□

**Example 3.8:**

A SDC labeling of the  $C(4, \Gamma())$  is given in figure 4 where  $n=4, p=5$ .

Let  $G = C(4, \Gamma())$ ,

$V(G) = \{u_{1,1}, u_{1,2}, u_{1,3}, u_{1,4}, u_{2,1}, u_{2,2}, u_{2,3}, u_{2,4}, u_{3,1}, u_{3,2}, u_{3,3}, u_{3,4}, u_{4,1}, u_{4,2}, u_{4,3}, u_{4,4}\}$  and

$E(G) = \{u_{j,1}u_{j,2}, u_{j,1}u_{j,3}, u_{j,4}u_{j,2}, u_{j,4}u_{j,3}, u_{j,2}u_{j,3}, u_{j,1}u_{j,4}\}$

Therefore,  $|V(G)|=16$  and  $|E(G)|=28$

Here  $|0|=12$  and  $|1|=16$

Hence  $|0| - |1|$  then  $C(4, (\Gamma()))$  is not a SDC graph.




Figure 4. C (4,Γ 0)

**Theorem 3.9.** For any prime number  $p > 2$ , cycle of join graph,  $C(3, \Gamma(Z_4))$  is a SDC graph.

□

**Theorem 3.10.** For any prime number  $p > 2$ , cycle of join graph,  $C(3, \Gamma(Z_6))$  is a SDC graph.

**Proof:** Let  $G = C(3, \Gamma(Z_6))$

Then vertex set of the graph  $G$  is

$$V(G) = \{u_{1,1}, u_{1,2}, \dots, u_{1,p-1}, x_1, y_1, z_1, u_{2,1}, u_{2,2}, \dots, u_{2,p-1}, x_2, y_2, z_2, u_{3,1}, u_{3,2}, \dots, u_{3,p-1}, x_3, y_3, z_3\}$$

And the edge set of the graph  $G$  is

$$E(G) = \{u_{i,x_j}, u_{i,y_j}, u_{i,z_j}, x_j y_j, y_j z_j, u_{1,1} u_{2,1}, u_{2,1} u_{3,1}, u_{3,1} u_{1,1} \text{ and } \}$$

Therefore  $|V(G)| = 3p+6$  and  $|E(G)| = 9p$

Define the vertex labeling  $f: V(G) \rightarrow \{1, 2, \dots, 3p+6\}$

By  $f(u_{1,i}) = i$  for

$f(u_{2,i}) = p+2+i$  for ,

$f(u_{3,i}) = 2p+4+i$  for ,

$f(x_i) = p, f(y_i) = p+2, f(z_i) = p+1$

$f(x_2) = 2p+2, f(y_2) = 2p+4, f(z_2) = 2p+3, f(x_3) = 3p+4, f(y_3) = 3p+6, f(z_3) = 3p+5.$

then the induced edge labeling is given by  $f^*: E(G) \rightarrow \{0, 1\}$  is given by

$$f^*(u_{j,i} x_j) =$$

$$f^*(u_{j,i} y_j) =$$

$$f^*(u_{j,i} z_j) =$$

$$f^*(x_j y_j) = 0, f^*(y_j z_j) = 1, f^*(u_{1,1} u_{2,1}) = 1, f^*(u_{2,1} u_{3,1}) = 1, f^*(u_{3,1} u_{1,1}) = 0.$$

we note that  $(0) \mid = 3(0)+1 =$  and  $(1) \mid = 3(0)+2 =$ ,

Thus  $\mid (0) - (1) \mid$

Hence  $G$  is a SDC graph.

□

**Example 3.11:**

A SDC labeling of the  $C(3, \Gamma(Z_6))$  is given in figure 5 where  $n=3, p=7$ .

Let  $G = C(3, \Gamma(Z_6))$

$$V(G) = \{u_{1,1}, u_{1,2}, \dots, u_{1,6}, x_1, y_1, z_1, u_{2,1}, u_{2,2}, \dots, u_{2,6}, x_2, y_2, z_2, u_{3,1}, u_{3,2}, \dots, u_{3,6}, x_3, y_3, z_3\}$$

$$E(G) = \{u_{j,i} x_j, u_{j,i} y_j, u_{j,i} z_j, x_j y_j, y_j z_j, u_{1,1} u_{2,1}, u_{2,1} u_{3,1}, u_{3,1} u_{1,1} \text{ and } \}$$

Therefore,  $|V(G)| = 27$  and  $|E(G)| = 63$

Here  $(0) \mid = 31$  and  $(1) \mid = 32$

Hence  $\mid (0) - (1) \mid$  then  $C(3, \Gamma(Z_6))$  is a SDC graph.




Figure 5'. C (3. (+ Γ(Z<sub>6</sub>)))

**Theorem 3.12.**

For any prime number  $p > 2$ , cycle of join graph  $+ \Gamma(Z_p)$  is a SDC graph when  $n \geq 3$ .

**Proof:**

Let  $G = C(n. (+ \Gamma(Z_p)))$

The vertex set of the graph G is

$$V(G) = \{u_{1,1}, u_{1,2}, \dots, u_{1,p-1}, x_1, y_1, u_{2,1}, u_{2,2}, \dots, u_{2,p-1}, x_2, y_2, \dots, u_{n,1}, u_{n,2}, \dots, u_{n,p-1}, x_n, y_n\}$$

$$E(G) = \{u_{j,i}x_j, u_{j,i}y_j, x_jy_j, u_{j,i}u_{j+1,i}\}$$

Therefore  $|V(G)| = n(p+1)$  and

$$|E(G)| = n(2p-1) + n = 2np$$

Define the vertex labeling.

$$f: V(G) \rightarrow \{1, 2, 3, \dots, n(p+1)\} \text{ by}$$

$$f(u_{1,i}) = i \text{ for ,}$$

$$f(u_{2,i}) = p+1+i \text{ for ,}$$

$$f(u_{3,i}) = 2p+2+i \text{ for ,}$$

.

.

.

$$f(u_{n,i}) = (n-1)p + (n-1) + i \text{ for}$$

$$f(x_1) = p, f(x_2) = 2p+1, f(x_3) = 3p+2, \dots, f(x_n) = np + (n-1),$$

$$f(y_1) = p+1, f(y_2) = 2p+2, f(y_3) = 3p+3, \dots, f(y_n) = np + n.$$

Then the induced edge labeling is given by

$$f(u_{j,i}x_j) =$$

$$f(u_{j,i}y_j) =$$

$$f(x_jy_j) = 1, f(u_{j,i}u_{j+1,i}) = 0, f(u_nu_{11}) = 0.$$

we observe that  $(0) = n(p-1) + n = np$  and  $(1) = np$ . Thus  $| (0) - (1) | = 0$ . Hence G is a SDC graph.

□

**Example 3.13:**

A SDC labeling of the  $C(4. (+ \Gamma(Z_5)))$  is given in figure 6 where  $n=4, p=5$ .

Let  $G = C(4. (+ \Gamma(Z_5)))$

$$V(G) = \{u_{1,1}, u_{1,2}, \dots, u_{1,4}, x_1, y_1, u_{2,1}, u_{2,2}, \dots, u_{2,4}, x_2, y_2, \dots, u_{4,1}, u_{4,2}, \dots, u_{4,4}, x_4, y_4\}$$

$$E(G) = \{u_{j,i}x_j, u_{j,i}y_j, x_jy_j, u_{j,i}u_{j+1,i}, u_{4,i}u_{1,i}\}$$

Therefore,  $|V(G)| = 24$  and  $|E(G)| = 40$

Here  $(0) = 20$  and  $(1) = 20$

Hence  $| (0) - (1) | = 0$  then  $C(4. (+ \Gamma(Z_5)))$  is a SDC graph.




Figure 6.  $C(4, (+\Gamma(Z_9)))$

Remark: In join graphs we used only paths in zero-divisor graphs,  $Z_4, Z_6, Z_9$ .

#### IV. Conclusion

The cycle of graphs are

- $C(n, (\Gamma(Z_{2p}) + \Gamma(Z_4)))$ ,  $n \geq 3$  and  $p > 2$  is a SDC graph.
- $C(n, (\Gamma(Z_{2p}) + \Gamma(Z_6)))$ ,  $n \geq 3$  and  $p > 2$  is a SDC graph.
- $C(n, (\Gamma(Z_{2p}) + \Gamma(Z_9)))$ ,  $n \geq 3$  and  $p > 2$  is not a SDC graph.
- $C(n, \Gamma())$ ,  $n \geq 3$  and  $p > 2$  is not a SDC graph.
- $C(n, (+\Gamma(Z_4)))$ ,  $n \geq 3$  and  $p > 2$  is a SDC graph.
- $C(n, (+\Gamma(Z_6)))$ ,  $n \geq 3$  and  $p > 2$  is a SDC graph.
- $C(n, (+\Gamma(Z_9)))$ ,  $n \geq 3$  and  $p > 2$  is a SDC graph.

#### References:

- [1] B.M. Stewart, Magic Graphs, Canadian J.Math, 18(1966), 1031-1059.
- [2] A. Rosa, On Certain Valuations Of The Vertices Of A Graph, Theory Of Graphs (Rome, July 1966), Gordon And Breach, N.Y. And Paris, 1967 349-355.
- [3] R. Frucht And F. Harary, On The Corona Of Two Graphs, Aequationes Math. 4(1970), 322-325.
- [4] Graph Theory With Applications To Engineering & Computer Science By Narsingh Deo, 1974.
- [5] Graph Theory With Applications By J.A Bondy And U.S.R. Murty, 1976
- [6] I. Beck, Zero Divisor Graph Of Commutative Rings. J Algebra. 1988; 116:208-226.
- [7] D.F Anderson And P. S Livingston, The Zero-Divisor Graph Of A Commutative Ring, J.Algebra, 217 (1999), 434-447.
- [8] S.P Redmond, On Zero-Divisor Graphs Of Small Finite Commutative Rings, Discrete Math., 307 (9-10) (2007), 1155-1166.
- [9] J. A. Gallian, A Dynamic Survey Of Graph Labeling, The Electronic J. Combin., 18 2015 #DS16.
- [10] A. Lourdasamy And F.Patrick, Sum Divisor Cordial Graphs, Proyecciones Journal Of Mathematics, 35(1), 2016 115-132.
- [11] S. Sajana, D. Bharathi, Intersection Graph Of Zero- Divisors Of A Finite Commutative Ring, International Journal Of Pure And Applied Mathematics 109 (2016) ,51-58.
- [12] D. Eswara Rao, D. Bharathi, Total Zero Divisor Graph Of A Commutative Ring, International Journal Of Scientific Research 2 (2013),28-29
- [13] T.T Chelvam And C. Subramanian, Signed Product Cordial Labeling Of Zero-Divisor Graphs, International Journal Of Mathematical Combinatorics, Special 1(2018), 164-169.
- [14] Kaneria V.J., Makadia H.M. And Jariya M. M., Graceful Labeling For Cycle Of Graphs, Int.J.Of Maths.Res.,6(2) (2014),173-178.