

## On Fuzzy Neutrosophic Pre $\Sigma$ -Baire Spaces.

E. Poongothai, E. Kalaivani

Department Of Mathematics, Shanmuga Industries Arts And Science College (Co-Ed.), Tiruvannamalai-606603, Tamil Nadu, India

---

### Abstract

In this paper, the concepts of fuzzy neutrosophic pre  $\sigma$ -nowhere dense set, fuzzy neutrosophic pre  $\sigma$ -first category set and fuzzy neutrosophic pre  $\sigma$ -second category set in fuzzy neutrosophic topological spaces are introduced and studied. By means of fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets, the concept of fuzzy neutrosophic pre  $\sigma$ -Baire space is defined and several characterizations of fuzzy neutrosophic pre  $\sigma$ -Baire spaces are studied. Several examples are given to illustrate the concepts introduced in this paper.

**Keywords:** Fuzzy neutrosophic pre-open(closed) set, fuzzy neutrosophic pre  $F\sigma$ -set, fuzzy neutrosophic pre  $G\delta$ -set, fuzzy neutrosophic pre dense, fuzzy neutrosophic pre residual, fuzzy neutrosophic nowhere dense set, fuzzy neutrosophic pre  $\sigma$ -first and second category sets, fuzzy neutrosophic pre  $\sigma$ -Baire space.

Date of Submission: 13-07-2025

Date of Acceptance: 23-07-2025

---

### I. Introduction

The fuzzy idea was invaded all branches of science as far back as the presentation of fuzzy sets by L. A. Zadeh [21]. The important concept of fuzzy topological space was offered by C.L. Chang [3]. The idea of fuzzy  $\sigma$ -Baire Spaces was introduced by G. Thangaraj and E. Poongothai [13]. The concept of neutrosophic sets was defined with membership, non-membership and indeterminacy degrees. In 2017, Veereswari [20] introduced fuzzy neutrosophic topological spaces. The idea of fuzzy neutrosophic Baire spaces was introduced by E. Poongothai and E. Padmavathi [10]. In this paper, the concepts of fuzzy neutrosophic pre  $\sigma$ -nowhere dense set, fuzzy neutrosophic pre  $\sigma$ -first category set and fuzzy neutrosophic pre  $\sigma$ -second category set in fuzzy neutrosophic topological spaces are introduced and studied. By means of fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets, the concept of fuzzy neutrosophic pre  $\sigma$ -Baire space is defined and several characterizations of fuzzy neutrosophic pre  $\sigma$ -Baire spaces are studied. Several examples are given to illustrate the concepts introduced in this paper.

## II. Preliminaries

**Definition 2.1** [2] A fuzzy neutrosophic set  $A$  on the universe of discourse  $X$  is defined as  $A = (x, T_A(x), I_A(x), F_A(x)), x \in X$  where  $T, I, F : X \rightarrow [0, 1]$  and  $0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3$ .

**Definition 2.2** [2] A fuzzy neutrosophic set  $A$  is a subset of a fuzzy neutrosophic set  $B$  (i.e.,)  $A \subseteq B$  for all  $x$  if  $T_A(x) \leq T_B(x), I_A(x) \leq I_B(x), F_A(x) \geq F_B(x)$ .

**Definition 2.3** [2] Let  $X$  be a non-empty set, and  $A = (x, T_A(x), I_A(x), F_A(x)), B = (x, T_B(x), I_B(x), F_B(x))$  be two fuzzy neutrosophic sets. Then

$$\begin{aligned} A \cup B &= (x, \max(T_A(x), T_B(x)), \max(I_A(x), I_B(x)), \min(F_A(x), F_B(x))) \\ A \cap B &= (x, \min(T_A(x), T_B(x)), \min(I_A(x), I_B(x)), \max(F_A(x), F_B(x))) \end{aligned}$$

**Definition 2.4** [2] The difference between two fuzzy neutrosophic sets  $A$  and  $B$  is defined as  $A \setminus B(x) = (x, \min(T_A(x), F_B(x)), \min(I_A(x), 1 - I_B(x)), \max(F_A(x), T_B(x)))$ .

**Definition 2.5** [2] A fuzzy neutrosophic set  $A$  over the universe  $X$  is said to be null or empty fuzzy neutrosophic set if  $T_A(x) = 0, I_A(x) = 0, F_A(x) = 1$  for all  $x \in X$ . It is denoted by  $0_N$ .

**Definition 2.6** [2] A fuzzy neutrosophic set  $A$  over the universe  $X$  is said to be absolute (universe) fuzzy neutrosophic set if  $T_A(x) = 1, I_A(x) = 1, F_A(x) = 0$  for all  $x \in X$ . It is denoted by  $1_N$ .

**Definition 2.7** [2] The complement of a fuzzy neutrosophic set  $A$  is denoted by  $A^c$  and is defined as  $A^c = (x, T_{A^c}(x), I_{A^c}(x), F_{A^c}(x))$  where  $T_{A^c}(x) = F_A(x), I_{A^c}(x) = 1 - I_A(x), F_{A^c}(x) = T_A(x)$ . The complement of fuzzy neutrosophic set  $A$  can also be defined as  $A^c = 1_N - A$ .

**Definition 2.8** [1] A fuzzy neutrosophic topology on a non-empty set  $X$  is a  $\tau$  of fuzzy neutrosophic sets in  $X$  satisfying the following axioms.

- (i)  $0_N, 1_N \in \tau$
- (ii)  $A_1 \cup A_2 \in \tau$  for any  $A_1, A_2 \in \tau$
- (iii)  $\bigcup_{i \in I} A_i \in \tau$  for any arbitrary family  $\{A_i : i \in I\} \subseteq \tau$

In this case the pair  $(X, \tau)$  is called fuzzy neutrosophic topological space and any fuzzy neutrosophic set in  $\tau$  is known as fuzzy neutrosophic open set in  $X$ .

**Definition 2.9** [1] The complement  $A^c$  of a fuzzy neutrosophic set  $A$  in a fuzzy neutrosophic topological space  $(X, \tau)$  is called fuzzy neutrosophic closed set in  $X$ .

**Definition 2.10** [1] Let  $(X, \tau)$  be a fuzzy neutrosophic topological space and  $A = (x, T_A(x), I_A(x), F_A(x))$  be a fuzzy neutrosophic set in  $X$ . Then the closure and interior of  $A$  are defined by

$$\begin{aligned} \text{int}(A) &= \bigcup \{G : G \text{ is a fuzzy neutrosophic open set in } X \text{ and } G \subseteq A\} \\ \text{cl}(A) &= \bigcap \{G : G \text{ is a fuzzy neutrosophic closed set in } X \text{ and } A \subseteq G\} \end{aligned}$$

**Definition 2.11** [1] Let  $(X, \tau)$  be a fuzzy neutrosophic topological space over  $X$ . Then the following properties hold, (i)  $\text{cl}(A^c) = (\text{int } A)^c$ , (ii)  $\text{int}(A^c) = (\text{cl } A)^c$ .

**Definition 2.12** [10] A fy. neutrosophic set  $A_N$  in a fy. neutrosophic top. space  $(P, \tau_N)$  is called a fy. neutrosophic  $F_\sigma$ -set if  $A_N = \bigcup_{i=1}^{\infty} A_{N_i}$  where  $A_{N_i} \in \tau_N$  for  $i \in I$ .

**Definition 2.13** [10] A fy. neutrosophic set  $A_N$  in a fy. neutrosophic top. space  $(P, \tau_N)$  is called a fy. neutrosophic  $G_\delta$ -set in  $(P, \tau_N)$  if  $A_N = \bigcap_{i=1}^{\infty} A_{N_i}$  where  $A_{N_i} \in \tau_N$  for  $i \in I$ .

**Definition 2.14** [10] A fy. neutrosophic set  $A_N$  in a fy. neutrosophic top. space  $(P, \tau_N)$  is called a fy. neutrosophic dense if there exist no  $fnCS B_N$  in  $(P, \tau_N)$  s.t.  $A_N \checkmark B_N \checkmark 1_X$ . That is  $fn(A_N)' = 1_N$ .

**Definition 2.15** [10] A fy. neutrosophic set  $A_N$  in a fy. neutrosophic top. space  $(P, \tau_N)$  is called a fy. neutrosophic nowh. dense set if there exist no non zero  $fnOS B_N$  in  $(P, \tau_N)$  s.t.  $B_N \checkmark fn(A_N)'$ . That is,  $fn(((A_N)')) = 0_N$ .

**Definition 2.16** [10] Let  $(P, \tau_N)$  be a fy. neutrosophic top. space. A fy. neutrosophic set  $A_N$  in  $(P, \tau_N)$  is called fy. neutrosophic one category set if  $A_N = \bigcup_{i=1}^{\infty} A_{N_i}$ , where  $A_{N_i}$ 's are fy. neutrosophic nowh. dense sets in  $(P, \tau_N)$ .

Any other fy. neutrosophic set in  $(P, \tau_N)$  is said to be of fy. neutrosophic two category.

**Definition 2.17** [10] A fy. neutrosophic top. space  $(P, \tau_N)$  is called fy. neutrosophic one category space if the fy. neutrosophic set  $1_N$  is a fy. neutrosophic one category set in  $(P, \tau_N)$ . That is  $1_N = \bigcup_{i=1}^{\infty} A_{N_i}$ , where  $A_{N_i}$ 's are fy. neutrosophic nowh. dense sets in  $(P, \tau_N)$ . Otherwise  $(P, \tau_N)$  will be called a fy. neutrosophic two category space.

**Definition 2.18** [10] Let  $A_N$  be a fy. neutrosophic one category set in  $(P, \tau_N)$ . Then  $A_N^c$  is called fy. neutrosophic re. set in  $(P, \tau_N)$ .

**Definition 2.19** [10] A fy. neutrosophic top. space  $(P, \tau_N)$  is called fy. neutrosophic Baire space if  $fn(\bigcup_{i=1}^{\infty} (A_{N_i})^c) = 0_N$ , where  $(A_{N_i})$ 's are fy. neutrosophic nowh. dense sets in  $(P, \tau_N)$ .

**Theorem 2.1** [10] Let  $(P, \tau_N)$  be a fy. neutrosophic top. space. Then the following are equivalent

- (1)  $(P, \tau_N)$  is a fy. neutrosophic Baire space.
- (2)  $fn(A_N)^c = 0_N$ , for every fy. neutrosophic one category set  $A_N$  in  $(P, \tau_N)$ .
- (3)  $fn(B_N)^c = 1_N$ , for every fy. neutrosophic re. set  $B_N$  in  $(P, \tau_N)$ .

**Definition 2.20** [5] Let  $(X_N, \tau_N)$  be a fuzzy neutrosophic topological space. A fuzzy neutrosophic Set  $\lambda_N$  in  $(X_N, \tau_N)$  is called a fuzzy neutrosophic  $\sigma$ -nowhere dense set if  $\lambda_N$  is a fuzzy neutrosophic  $F_\sigma$ -set in  $(X_N, \tau_N)$  such that  $int(\lambda_N) = 0_N$ .

**Definition 2.21** [5] Let  $(X_N, \tau_N)$  be a fuzzy neutrosophic topological space. A fuzzy neutrosophic set  $\lambda_N$  in  $(X_N, \tau_N)$  is called fuzzy neutrosophic  $\sigma$ -first category set if  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_{N_i})$ , where  $(\lambda_{N_i})$ 's are fuzzy neutrosophic  $\sigma$ -nowhere dense sets in  $(X_N, \tau_N)$ . Any other fuzzy neutrosophic set in  $(X_N, \tau_N)$  is said to be fuzzy neutrosophic  $\sigma$ -second category sets in  $(X_N, \tau_N)$ .

**Definition 2.22** [5] Let  $\lambda_N$  be a fuzzy neutrosophic  $\sigma$ -first category set in  $(X_N, \tau_N)$ . Then  $1_N - \lambda_N$  is called a fuzzy neutrosophic  $\sigma$ -residual set in  $(X_N, \tau_N)$ .

**Definition 2.23** [5] A fuzzy neutrosophic topological space  $(X_N, T_N)$  is called fuzzy neutrosophic  $\sigma$ -first category space if the fuzzy neutrosophic set  $1_{X_N}$  is a fuzzy neutrosophic  $\sigma$ -first category set in  $(X_N, T_N)$ . That is  $1_{X_N} = \bigcup_{i=1}^{\infty} (A_N)_i$ , where  $(A_N)_i$ 's are fuzzy neutrosophic  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Otherwise  $(X_N, T_N)$  will be called a fuzzy neutrosophic  $\sigma$ -second category space.

**Definition 2.24** [5] Let  $(X_N, T_N)$  be a fuzzy neutrosophic topological space. Then  $(X_N, T_N)$  is called a fuzzy neutrosophic  $\sigma$ -Baire space if  $\text{int}(\bigcup_{i=1}^{\infty} (A_N)_i) = 0_N$ , where  $(A_N)_i$ 's are fuzzy neutrosophic  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ .

**Theorem 2.2** [5] Let  $(X_N, T_N)$  be a fuzzy neutrosophic topological space. Then the following are equivalent

- (1)  $(X_N, T_N)$  is a fuzzy neutrosophic  $\sigma$ -Baire space.
- (2)  $\text{int}(A_N) = 0_N$ , for every fuzzy neutrosophic  $\sigma$ -first category set  $A_N$  in  $(X_N, T_N)$ .
- (3)  $\text{cl}(\mu_N) = 1_N$ , for every fuzzy neutrosophic  $\sigma$ -residual set  $\mu_N$  in  $(X_N, T_N)$ .

**Definition 2.25** [12] A fuzzy neutrosophic topological space  $(X, \tau_N)$  is called a fuzzy neutrosophic submaximal space if for each fuzzy neutrosophic set  $A_N$  in  $(X, \tau_N)$  such that  $(A_N)^- = 1$  then  $A_N \in \tau_N$  in  $(X, \tau_N)$ . That is  $(X, \tau_N)$  is a fuzzy neutrosophic submaximal space if each fuzzy neutrosophic dense set in  $(X, \tau_N)$  is a fuzzy neutrosophic open set in  $(X, \tau_N)$ .

**Definition 2.26** [12] A fuzzy neutrosophic topological space  $(X, \tau_N)$  is called a fuzzy neutrosophic resolvable space if there exist a fuzzy neutrosophic dense set  $A_N$  in  $(X, \tau_N)$  such that  $(1 - A_N)^- = 1$ . Otherwise,  $(X, \tau_N)$  is called a fuzzy neutrosophic irresolvable space.

**Definition 2.27** [12] A fuzzy neutrosophic topological space  $(X, \tau_N)$  is called a fuzzy neutrosophic hyperconnected space if every non-null fuzzy neutrosophic open subset of  $(X, \tau_N)$  is fuzzy neutrosophic dense in  $(X, \tau_N)$ .

**Definition 2.28** [12] A fuzzy neutrosophic topological space  $(X, \tau_N)$  is called a fuzzy neutrosophic  $P$ -space if each fuzzy neutrosophic  $G_\delta$ -set in  $(X, \tau_N)$  is fuzzy neutrosophic open set in  $(X, \tau_N)$ .

**Definition 2.29** [12] A fuzzy neutrosophic topological space  $(X, \tau_N)$  is called a fuzzy neutrosophic almost resolvable space if  $\bigcup_{i=1}^{\infty} (A_N)_i = 1$ , where  $(A_N)_i$ 's in  $(X, \tau_N)$  are such that  $(A_N)_i^- = 0$ , otherwise,  $(X, \tau_N)$  is called a fuzzy neutrosophic almost irresolvable space.

**Definition 2.30** [4] FNS  $\lambda_N$  in FNTS  $(X, \tau)$  is called Fuzzy neutrosophic regular - open set (Briefly, FNR-open) if  $\lambda_N = FNInt(FNcl(\lambda_N))$ .

**Definition 2.31** [4] FNS  $\lambda_N$  in FNTS  $(X, \tau)$  is called Fuzzy neutrosophic regular - closed set (Briefly, FNR-closed) if  $\lambda_N = FNcl(FNInt(\lambda_N))$ .

**Definition 2.32** [4] Fuzzy neutrosophic pre-open set (Briefly, FNP-open) if  $\lambda_N \check{D} FNInt(FNcl(\lambda_N))$ .

**Definition 2.33** [4] Fuzzy neutrosophic pre-closed set (Briefly, FNP-closed) if  $FNcl(FNInt(\lambda_N)) \check{D} (\lambda_N)$ .

### 3 Fuzzy Neutrosophic Pre $\sigma$ —Nowhere Dense sets

**Definition 3.1** A fuzzy neutrosophic set  $\lambda_N$  in a fuzzy neutrosophic topological space  $(X_N, T_N)$  is called a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$  if  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre-closed sets in  $(X_N, T_N)$ .

**Definition 3.2** A fuzzy neutrosophic set  $\lambda_N$  in a fuzzy neutrosophic topological space  $(X_N, T_N)$  is called a fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$  if  $\lambda_N = \bigcap_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre-open sets in  $(X_N, T_N)$ .

**Definition 3.3** A fuzzy neutrosophic set  $\lambda_N$  in a fuzzy neutrosophic topological space  $(X_N, T_N)$  is called a fuzzy neutrosophic pre dense if there exist no fuzzy neutrosophic pre-closed set where  $\mu_N$  in  $(X_N, T_N)$  such that  $\lambda_N \check{a} \mu_N \check{a} 1_N$ . That is,  $pcl(\lambda_N) = 1_N$  in  $(X_N, T_N)$ .

**Definition 3.4** A fuzzy neutrosophic set  $\lambda_N$  in a fuzzy neutrosophic topological space  $(X_N, T_N)$  is called a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set if  $\lambda_N$  is a non-zero fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$  such that  $pint(\lambda_N) = 0_N$ .

**Example 3.1** Let  $X_N = \{a, b, c\}$ . The fuzzy neutrosophic sets  $\lambda_N, \mu_N$  and  $\gamma_N$  are defined on  $X_N$  as follows:

$\lambda_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$\lambda_N = \{(a, (0.7, 0.6, 0.7)), (b, (0.5, 0.6, 0.5)), (c, (0.7, 0.7, 0.6))\}$$

$\mu_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$\mu_N = \{(a, (0.5, 0.4, 0.8)), (b, (0.7, 0.6, 0.5)), (c, (0.8, 0.6, 0.6))\}$$

$\gamma_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$\gamma_N = \{(a, (0.6, 0.7, 0.5)), (b, (0.5, 0.6, 0.6)), (c, (0.7, 0.6, 0.6))\}$$

Then,  $T_N = \{0_N, \lambda_N, \mu_N, \gamma_N, \lambda_N \vee \mu_N, \mu_N \vee \gamma_N, \lambda_N \vee \gamma_N, \lambda_N \times \mu_N, \mu_N \times \gamma_N, \lambda_N \times \gamma_N, 1_N\}$  is a fuzzy neutrosophic topology on  $X_N$ . The fuzzy neutrosophic set  $\delta_N = [(1_N - \lambda_N) \vee (1_N - \mu_N) \vee (1_N - (\lambda_N \vee \mu_N))]$  in  $(X_N, T_N)$ . Then  $\delta_N$  is a fuzzy neutrosophic pre  $F$ -set in  $(X_N, T_N)$  and  $\text{pint}(\delta_N) = 0_N$  and hence  $\delta_N$  is a fuzzy neutrosophic pre  $\alpha$ -nowhere dense set in  $(X_N, T_N)$ .

**Example 3.2** Let  $X_N = \{a, b\}$ . The fuzzy neutrosophic sets  $\alpha_N$  and  $\delta_N$  are defined on  $X_N$  as follows:

$$\alpha_N : X_N \rightarrow [0_N, 1_N] \text{ is defined as } \alpha_N = \{(a, (0.4, 0.3, 0.4)), (b, (0.5, 0.4, 0.3))\}$$

$$\delta_N : X_N \rightarrow [0_N, 1_N] \text{ is defined as } \delta_N = \{(a, (0.5, 0.3, 0.3)), (b, (0.4, 0.3, 0.5))\}$$

Then,  $T_N = \{0_N, \alpha_N, \delta_N, \alpha_N \vee \delta_N, \alpha_N \times \delta_N, 1_N\}$  is a fuzzy neutrosophic topology on  $X_N$ . Now, consider

$$\eta_N = [(1_N - (\alpha_N \vee \delta_N)) \vee (1_N - (\alpha_N \times \delta_N))]$$

$$\eta_N = 1_N - (\alpha_N \times \delta_N)$$

Therefore  $\eta_N$  is a fuzzy neutrosophic pre  $F$ -set in  $(X_N, T_N)$ .  $\text{pint}(\eta_N) \neq 0_N$ .

Therefore  $\eta_N$  is not a fuzzy neutrosophic pre  $\alpha$ -nowhere dense set in  $(X_N, T_N)$ .

**Remark 3.1** If  $\lambda_N$  and  $\mu_N$  are fuzzy neutrosophic pre  $\alpha$ -nowhere dense sets in a fuzzy neutrosophic topological space  $(X_N, T_N)$ , then  $\lambda_N \vee \mu_N$  be a fuzzy neutrosophic pre  $\alpha$ -nowhere dense set in  $(X_N, T_N)$ . For, consider the following example:

**Example 3.3** Let  $X_N = \{a, b, c\}$ . The fuzzy neutrosophic sets  $A_N, B_N$  and  $C_N$  are defined  $X_N$  as follows:

$A_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$A_N = \{(a, (0.7, 0.6, 0.6)), (b, (0.7, 0.5, 0.6)), (c, (0.5, 0.7, 0.6))\}$$

$B_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$B_N = \{(a, (0.5, 0.4, 0.5)), (b, (0.4, 0.7, 0.6)), (c, (0.8, 0.5, 0.6))\}$$

$C_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$C_N = \{(a, (0.7, 0.6, 0.6)), (b, (0.6, 0.5, 0.5)), (c, (0.5, 0.4, 0.5))\}$$

Then,  $T_N = \{0_N, A_N, B_N, C_N, A_N \vee B_N, B_N \vee C_N, A_N \vee C_N, A_N \times B_N, B_N \times C_N, A_N \times C_N, A_N \vee B_N \vee C_N, A_N \times B_N \times C_N, 1_N\}$  is clearly a fuzzy neutrosophic topology on  $X_N$ .

Now, consider,  $a_N = [(1_N - A_N) \vee (1_N - B_N) \vee (1_N - C_N)] = [1_N - (A_N \times B_N)]$

Therefore  $a_N$  is a fuzzy neutrosophic pre  $F_2$ -set in  $(X_N, T_N)$ .

$$\delta_N = [(1_N - (\delta_N \vee C_N)) \vee (1_N - (A_N \vee C_N)) \vee (1_N - (A_N \times B_N))] = [1_N - (A_N \times B_N)]$$

$\text{pint}(a_N) = 0_N$ , which implies that  $a_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$

$(1_N - C_N) \vee (1_N - (A_N \vee B_N \vee C_N)) = \gamma_N$  is a fuzzy neutrosophic pre  $F_2$ -set in  $(X_N, T_N)$ .

$\text{pint}(\gamma_N) = 0_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

$\text{pint}(a_N \vee \gamma_N) = 0_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

**Proposition 3.1** A fuzzy neutrosophic set  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in a fuzzy neutrosophic topological space  $(X_N, T_N)$  if and only if  $(1_N - \lambda_N)$  is a fuzzy neutrosophic pre dense and fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$ .



**Proof.** Let  $\lambda_N$  be a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set  $(X_N, T_N)$ . Then  $\lambda_N = \bigcup_{i=1}^{\infty} \lambda_N^i$ , where  $\lambda_N^i$ 's are fuzzy neutrosophic pre-closed sets in  $(X_N, T_N)$  and  $\text{pint}(\lambda_N) = 0_N$ . Then  $1_N - \text{pint}(\lambda_N) = 1_N - 0_N = 1_N$  and hence  $\text{pcl}(1_N - \lambda_N) = 1_N$ . Also  $(1_N - \lambda_N) = 1_N - \bigcup_{i=1}^{\infty} \lambda_N^i = \bigcap_{i=1}^{\infty} (1_N - \lambda_N^i)$ , where  $(1_N - \lambda_N^i)$ 's are fuzzy neutrosophic pre-open sets in  $(X_N, T_N)$ , implies that  $1_N - \lambda_N$  is a fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$ . Hence  $(1_N - \lambda_N)$  is a fuzzy neutrosophic pre dense and fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$ . Conversely, Let  $\lambda_N$  be a fuzzy neutrosophic pre dense and fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$ . Then  $\lambda_N = \bigcup_{i=1}^{\infty} \lambda_N^i$ , where  $\lambda_N^i$ 's are fuzzy pre-open sets in  $(X_N, T_N)$ . Now,  $1_N - \lambda_N = 1_N - \bigcup_{i=1}^{\infty} \lambda_N^i = \bigcap_{i=1}^{\infty} (1_N - \lambda_N^i)$ , where  $(1_N - \lambda_N^i)$ 's are fuzzy neutrosophic pre-closed sets in  $(X_N, T_N)$ . Hence  $(1_N - \lambda_N)$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$  and  $\text{pint}(1_N - \lambda_N) = 1_N - \text{pcl}(\lambda_N) = 1_N - 1_N = 0_N$ . [Since  $\lambda_N$  is a fuzzy neutrosophic pre dense in  $(X_N, T_N)$ ]. Therefore  $(1_N - \lambda_N)$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

**Proposition 3.2** If  $\lambda_N$  is a fuzzy neutrosophic pre dense set in a fuzzy neutrosophic topological space  $(X_N, T_N)$  such that  $\mu_N \subseteq (1_N - \lambda_N)$ , where  $\mu_N$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$ , then  $\mu_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

**Proof.** Let  $\lambda_N$  be a fuzzy neutrosophic pre dense set in  $(X_N, T_N)$  such that  $\mu_N \subseteq (1_N - \lambda_N)$ . Now,  $\mu_N \subseteq (1_N - \lambda_N)$ , implies that  $\text{pint}(\mu_N) \subseteq \text{pint}(1_N - \lambda_N)$ . Then  $\text{pint}(\mu_N) \subseteq 1_N - \text{pcl}(\lambda_N) = 1_N - 1_N = 0_N$  and hence  $\text{pint}(\mu_N) = 0_N$ . Therefore,  $\mu_N$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$  such that  $\text{pint}(\mu_N) = 0_N$  and hence  $\mu_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

**Definition 3.5** Let  $(X_N, T_N)$  be a fuzzy neutrosophic topological space. A fuzzy

neutrosophic set  $\lambda_N$  in  $(X_N, T_N)$  is called fuzzy neutrosophic pre  $\sigma$ -first category set if  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Any other fuzzy neutrosophic set in  $(X_N, T_N)$  is said to be of fuzzy neutrosophic pre  $\sigma$ -second category set in  $(X_N, T_N)$ .

**Definition 3.6** Let  $\lambda_N$  be a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ . Then  $(1_N - \lambda_N)$  is called a fuzzy neutrosophic pre  $\sigma$ -residual set in  $(X_N, T_N)$ .

**Definition 3.7** A fuzzy neutrosophic topological space  $(X_N, T_N)$  is called fuzzy neutrosophic pre  $\sigma$ -first category space is the fuzzy neutrosophic set  $1_{X_N}$  if a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ . That is,  $1_{X_N} = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Otherwise,  $(X_N, T_N)$  will be called a fuzzy neutrosophic pre  $\sigma$ -second category space.

**Proposition 3.3** If  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -first category set in a fuzzy neutrosophic topological space  $(X_N, T_N)$ , then there is a fuzzy neutrosophic pre  $F_\sigma$ -set  $\delta_N$  in  $(X_N, T_N)$  such that  $\lambda_N \subseteq \delta_N$ .

**Proof.** Let  $\lambda_N$  be a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ . Then  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Now,  $[1_N - pcl(\lambda_N)_i]$ 's ( $i = 1$  to  $\infty$ ) are fuzzy neutrosophic pre-open sets in  $(X_N, T_N)$ . Then  $\mu_N = \bigcup_{i=1}^{\infty} [1_N - pcl(\lambda_N)_i]$  is a fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$  and  $1_N - \mu_N = 1_N - \bigcup_{i=1}^{\infty} [1_N - pcl(\lambda_N)_i] = \bigcap_{i=1}^{\infty} [pcl(\lambda_N)_i]$ . Now,  $(\lambda_N)_i \subseteq pcl(\lambda_N)_i$ , implies that  $\bigcup_{i=1}^{\infty} (\lambda_N)_i \subseteq \bigcup_{i=1}^{\infty} [pcl(\lambda_N)_i]$ . Hence  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i \subseteq \bigcup_{i=1}^{\infty} [pcl(\lambda_N)_i] = [1_N - \mu_N]$ . That is,  $\lambda_N \subseteq [1_N - \mu_N]$  and  $[1_N - \mu_N]$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$ . Let  $\delta_N = [1_N - \mu_N]$ . Hence, if  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ , then there is a fuzzy neutrosophic pre  $F_\sigma$ -set  $\delta_N$  in  $(X_N, T_N)$  such that  $\lambda_N \subseteq \delta_N$ .

**Proposition 3.4** *If  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -first category set in a fuzzy neutrosophic topological space  $(X_N, T_N)$ , then there is a fuzzy neutrosophic pre  $F_\sigma$ -set  $\delta_N$  in  $(X_N, T_N)$  such that  $\lambda_N \leq \delta_N \leq cl(\lambda_N)$ , where  $\delta_N$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$ .*

**Proof.** Let  $\lambda_N$  be a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ . Then  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Now,  $[1_N - pcl(\lambda_N)]_i$ 's ( $i = 1$  to  $\infty$ ) are fuzzy neutrosophic pre-open sets in  $(X_N, T_N)$ . Then,  $\mu_N = \bigcup_{i=1}^{\infty} (1_N - pcl(\lambda_N))_i$  is a fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$  and  $1_N - \mu_N = 1_N - [\bigcup_{i=1}^{\infty} (1_N - pcl(\lambda_N))_i] = [\bigcap_{i=1}^{\infty} pcl(\lambda_N)_i]$ . Now,  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i \leq [\bigcap_{i=1}^{\infty} pcl(\lambda_N)_i] \leq \bigcup_{i=1}^{\infty} (cl(\lambda_N))_i \leq cl(\bigcup_{i=1}^{\infty} (\lambda_N)_i)$ . That is,  $\lambda_N \leq [1_N - \mu_N] \leq cl(\lambda_N)$  and  $[1_N - \mu_N]$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$ . Let  $\delta_N = [1_N - \mu_N]$ .

Hence, if  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$  such that then there is a fuzzy neutrosophic pre  $F_\sigma$ -set  $\delta_N$  in  $(X_N, T_N)$   $\lambda_N \leq \delta_N \leq cl(\lambda_N)$ , where  $\delta_N$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$ .

**Proposition 3.5** *If  $\lambda_N$  is a fuzzy neutrosophic pre-closed set in a fuzzy neutrosophic topological space  $(X_N, T_N)$  and if  $pint(\lambda_N) = 0_N$ , then  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .*

**Proof.** Let  $\lambda_N$  be a fuzzy neutrosophic pre-closed set in  $(X_N, T_N)$ . Then we have  $pcl(\lambda_N) = \lambda_N$ . Now,  $pint[pcl(\lambda_N)] = pint(\lambda_N)$  and  $pint(\lambda_N) = 0_N$ , implies that  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

**Proposition 3.6** *If  $\lambda_N$  is a fuzzy neutrosophic closed and fuzzy neutrosophic  $\sigma$ -nowhere dense set in a fuzzy neutrosophic topological space  $(X_N, T_N)$ , then  $pint(\lambda_N) = 0_N$  in  $(X_N, T_N)$ .*

**Proof.** Let  $\lambda_N$  be a fuzzy neutrosophic  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ . Then  $\lambda_N$  is a fuzzy neutrosophic  $F_\sigma$ -set such that  $\text{int}(\lambda_N) = 0_N$ . We have,  $\text{pint}(\lambda_N) \subseteq \lambda_N \times \text{intcl}(\lambda_N)$ . Then,  $\text{pint}(\lambda_N) \subseteq \lambda_N \times \text{int}(\lambda_N)$ . [Since  $\lambda_N$  is a fuzzy neutrosophic closed set,  $\lambda_N = \text{cl}(\lambda_N)$ ] and hence  $\text{pint}(\lambda_N) \subseteq \lambda_N \times 0_N$ . That is,  $\text{pint}(\lambda_N) = 0_N$  in  $(X_N, T_N)$ .

**Proposition 3.7** If each fuzzy neutrosophic  $\sigma$ -nowhere dense set  $\lambda_N$  is a fuzzy neutrosophic closed set in a fuzzy neutrosophic topological space  $(X_N, T_N)$ , then  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

**Proof.** Let  $\lambda_N$  be a fuzzy neutrosophic  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ . Then  $\lambda_N$  is a fuzzy neutrosophic  $F_\sigma$ -set in  $(X_N, T_N)$  such that  $\text{int}(\lambda_N) = 0_N$ . We have,  $\text{pint}(\lambda_N) \subseteq \lambda_N \times \text{intcl}(\lambda_N)$ . Since  $\lambda_N$  is a fuzzy neutrosophic closed set in  $(X_N, T_N)$ ,  $\text{cl}(\lambda_N) = \lambda_N$ . Then  $\text{pint}(\lambda_N) \subseteq \lambda_N \times \text{int}(\lambda_N)$ . That is,  $\text{pint}(\lambda_N) \subseteq \lambda_N \times 0_N = 0_N$ . Hence,  $\text{pint}(\lambda_N) = 0_N$  and therefore  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

## 4 Fuzzy Neutrosophic Pre $\sigma$ -Baire Spaces

**Definition 4.1** Let  $(X_N, T_N)$  be a fuzzy neutrosophic topological space. Then  $(X_N, T_N)$  is called a fuzzy neutrosophic pre  $\sigma$ -Baire Space if  $\text{pint}(\bigcup_{i=1}^{\infty} \lambda_{N_i}) \neq 0_N$ , where  $(\lambda_{N_i})$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ .

**Example 4.1** Let  $X_N = \{a, b, c\}$ . The fuzzy neutrosophic sets  $A_N, B_N$  and  $C_N$  are defined on  $X_N$  as follows:

$A_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$A_N = \{(a, (0.7, 0.6, 0.5)), (b, (0.5, 0.6, 0.8)), (c, (0.7, 0.5, 0.6))\}$$

$B_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$B_N = \{(a, (0.6, 0.5, 0.7)), (b, (0.6, 0.6, 0.7)), (c, (0.5, 0.7, 0.5))\}$$

$C_N : X_N \rightarrow [0_N, 1_N]$  is defined as,

$$C_N = \{(a, (0.5, 0.5, 0.7)), (b, (0.7, 0.5, 0.5)), (c, (0.6, 0.5, 0.7))\}$$

Then,  $T_N = \{0_N, A_N, B_N, C_N, A_N \vee B_N, A_N \vee C_N, B_N \vee C_N, A_N \times B_N, A_N \times C_N, B_N \times C_N, A_N \times B_N \times C_N, 1_N\}$  is a fuzzy neutrosophic topology on  $X_N$ . Now,  $\alpha_N = [(1_N - B_N) \vee (1_N - C_N) \vee (1_N - (A_N \vee B_N))] = [1_N - (B_N \times C_N)]$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$ .

$\text{pint}(\alpha_N) = 0_N$ ,  $\alpha_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set in  $(X_N, T_N)$ .

$\delta_N = [(1_N - (A_N \times B_N)) \vee (1_N - A_N \times C_N) \vee (1_N - (B_N \times C_N))]$  is a fuzzy neutrosophic pre  $F_\sigma$ -set in  $(X_N, T_N)$ .

$\text{pint}(\delta_N) = 0_N$ ,  $\delta_N$  is a fuzzy neutrosophic pre  $\sigma$ -nowhere dense set.  $\text{pint}(\alpha_N \vee \delta_N) = 0_N$ , then  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -Baire space.

**Proposition 4.1** Let  $(X_N, T_N)$  be a fuzzy neutrosophic topological space. Then the following are equivalent:

- (1)  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -Baire Space.
- (2)  $\text{pint}(\lambda_N) = 0_N$ , for each fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ .
- (3)  $\text{pcl}(\mu_N) = 1_N$ , for each fuzzy neutrosophic pre  $\sigma$ -residual set  $\mu_N$  in  $(X_N, T_N)$ .

**Proof.** (1)  $\Rightarrow$  (2)

Let  $\lambda_N$  be a fuzzy neutrosophic  $\sigma$ -first category set in  $(X_N, T_N)$ . Then  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Then,  $\text{pint}(\lambda_N) = \text{pint}(\bigcup_{i=1}^{\infty} (\lambda_N)_i)$ . Since  $(X_N, T_N)$  is a fuzzy neutrosophic  $\sigma$ -Baire space,  $\text{pint}(\bigcup_{i=1}^{\infty} (\lambda_N)_i) = 0_N$ . Hence,  $\text{pint}(\lambda_N) = 0_N$  for a fuzzy neutrosophic pre  $\sigma$ -first category set  $\lambda_N$  in  $(X_N, T_N)$ .

(2)  $\Rightarrow$  (3)

Let  $\mu_N$  be a fuzzy neutrosophic pre  $\sigma$ -residual set  $\mu_N$  in  $(X_N, T_N)$ . Then  $(1_N - \mu_N)$  is a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ . By hypothesis,  $\text{pint}(1_N - \lambda_N) = 0_N$ . Then,  $1_N - \text{pcl}(\mu_N) = 0_N$ . Hence,  $\text{pcl}(\mu_N) = 1_N$ , for a fuzzy neutrosophic pre  $\sigma$ -residual set  $\mu_N$  in  $(X_N, T_N)$ .

(3)  $\Rightarrow$  (1)

Let  $\lambda_N$  be a fuzzy neutrosophic  $\sigma$ -first category set in  $(X_N, T_N)$ . Then  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Now,  $\lambda_N$  is a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ , implies that  $(1_N - \lambda_N)$  is a fuzzy neutrosophic pre  $\sigma$ -residual set in  $(X_N, T_N)$ . By hypothesis,  $\text{pcl}(1_N - \lambda_N) = 1_N$ . Then,  $1_N - \text{pint}(\lambda_N) = 1_N$ . Hence  $\text{pint}(\lambda_N) = 0_N$ . That is,  $\text{pint}(\bigcup_{i=1}^{\infty} (\lambda_N)_i) = 0_N$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Hence  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -Baire space.

**Proposition 4.2** If the fuzzy neutrosophic topological space  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -Baire space, then  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -second category space.

**Proof.** Let  $(X_N, T_N)$  be a fuzzy neutrosophic pre  $\sigma$ -Baire space. Then,  $\text{pint}(\bigcup_{i=1}^{\infty} (\lambda_N)_i) = 0_N$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Then  $\bigcup_{i=1}^{\infty} (\lambda_N)_i \neq 1_N$  [Otherwise,  $\bigcup_{i=1}^{\infty} (\lambda_N)_i = 1_N$ , implies that  $\text{pint}(\bigcup_{i=1}^{\infty} (\lambda_N)_i) = \text{pint}(1_N) = 1_N$ , which in turn implies that  $0_N = 1_N$ , a contradiction]. Hence  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -second category space.

**Proposition 4.3** Let  $(X_N, T_N)$  be a fuzzy neutrosophic topological space. If  $\bigcup_{i=1}^{\infty} (\lambda_N)_i \neq 0_N$ , where  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre dense and fuzzy neutrosophic pre G $\delta$ -sets in  $(X_N, T_N)$ , then  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -second category space.

**Proof.** Given that  $\bigcup_{i=1}^{\infty} (\lambda_N)_i \neq 0_N$ , implies that  $1_N - \bigcup_{i=1}^{\infty} (\lambda_N)_i \neq 1_N$ . Then  $\bigcap_{i=1}^{\infty} (1_N - (\lambda_N)_i) \neq 1_N$ . Since  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre dense and fuzzy neutrosophic pre  $G_\delta$ -set in  $(X_N, T_N)$ , by proposition 3.1.,  $(1_N - (\lambda_N)_i)$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Hence,  $\bigcap_{i=1}^{\infty} (1_N - (\lambda_N)_i) \neq 1_N$ , where  $(1_N - (\lambda_N)_i)$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Hence  $(X_N, T_N)$  is not a fuzzy neutrosophic pre  $\sigma$ -first category space. Therefore  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -second category space.

**Proposition 4.4** *If a fuzzy neutrosophic topological space  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -Baire space, then no non-zero fuzzy neutrosophic pre-open set is a fuzzy neutrosophic  $\sigma$ -first category set in  $(X_N, T_N)$ .*

**Proof.** Let  $\lambda_N$  be a non-zero fuzzy neutrosophic pre-open set in a fuzzy pre  $\sigma$ -Baire space  $(X_N, T_N)$ . Suppose that  $\lambda_N = \bigcup_{i=1}^{\infty} (\lambda_N)_i$ , where the fuzzy neutrosophic sets  $(\lambda_N)_i$ 's are fuzzy neutrosophic pre  $\sigma$ -nowhere dense sets in  $(X_N, T_N)$ . Then  $\text{pint}(\lambda_N) = \text{pint}(\bigcup_{i=1}^{\infty} (\lambda_N)_i)$ . Since  $(X_N, T_N)$  is a fuzzy neutrosophic pre  $\sigma$ -Baire space,  $\text{pint}(\bigcup_{i=1}^{\infty} (\lambda_N)_i) = 0_N$ . This implies that,  $\text{pint}(\lambda_N) = 0_N$ . Then we will have  $\lambda_N = \text{pint}(\lambda_N) = 0_N$ , a contradiction. Since  $\lambda_N$  is a non-zero fuzzy neutrosophic set in  $(X_N, T_N)$ . Hence no non-zero fuzzy neutrosophic pre-open set is a fuzzy neutrosophic pre  $\sigma$ -first category set in  $(X_N, T_N)$ .

### III. Conclusion

In this study, we have introduced and analyzed the concept of fuzzy neutrosophic pre  $\sigma$ -Baire Spaces, extending classical and fuzzy Baire space theories into the neutrosophic framework. The work lays a foundation for further exploration of fuzzy neutrosophic spaces in advanced topology, particularly in applications involving decision-making, artificial intelligence and Information systems, where vagueness and indeterminacy play a critical role.

### References

- [1] Arockiarani I.; J. Martina Jency. More On Fuzzy Neutrosophic Sets And Fuzzy Neutrosophic Topological Spaces., International Journal Of Innovative Research & Studies, Volume 3(5), 2014, 643-652.
- [2] Arockiarani I.; I. R. Sumathi.; J.Martina Jency. Fuzzy Neutrosophic Soft Topological Spaces. International Journal Of Mathematical Archive, Volume 4(10), 2013., 225-238.
- [3] Chang C.L, Fuzzy Topological Spaces, Journal Of Mathematical Analysis And Applications. Volume 24(1), 1968., 182-190.
- [4] Fatimah M. Mohammed I X And Anas A. Hijab I And Shaymaa F. Matar I. Fuzzy Neutrosophic Weakly-Generalized Closed Sets In Fuzzy Neutrosophic Topological Spaces, J.University Of Anbar For Pure Science : Vol.12(2), 2018, 63-73.
- [5] Poongothai,E.; Kalaivani, E. Neutrosophic  $\Sigma$ -Baire Spaces On Fuzzy Setting. Journal Of Xedian University, 2024, Volume 18(1.) Pp. 253-264.
- [6] Poongothai,E.; Kalaivani, E. A Characterization Of Neutrosophic  $\Sigma$ -Baire Space On Fuzzy Setting. Neutrosophic Sets And Systems, 2025, Volume 83, Pp. 87-95.
- [7] Poongothai,E.; Kalaivani, E. Some Remarks On Fuzzy Neutrosophic  $\Sigma$ -Baire Spaces And Other Fuzzy Neutrosophic Spaces, Communicated To The Journal Of Neutrosophic Sets And Systems.
- [8] Poongothai,E.; Kalaivani, E. Fuzzy Neutrosophic Regular  $\Sigma$ -Baire Space. Communicated To The Journal Of Neutrosophic Sets And Systems.
- [9] Ganesan Thangaraj.; Eswaran Poongothai. A Note On Fuzzy  $\Sigma$ -Baire Spaces, Annals Of Fuzzy Mathematics And Informatics, 2015, Volume 9(2), Pp. 197-203.
- [10] Poongothai,E.; Padmavathi, E. Baire Spaces On Fuzzy Neutrosophic Topological Spaces. Neutrosophic Sets And Systems, 2022, Volume 51(1), Pp. 708-723.
- [11] Poongothai,E.; Padmavathi, E. On Fuzzy Neutrosophic Directed Baire Spaces And Characterizations Of Other Fuzzy Neutrosophic Spaces, Mukta Shabd Journal, 2023, Volume 12(12), Pp. 570-586.
- [12] Poongothai,E.; Padmavathi, E. C-Almost P-Spaces And Other Spaces On Fuzzy Neutrosophic Topological Spaces. International Journal Of Advanced Research In Science, Engineering And Technology, Nov.2023, Volume 10(11).
- [13] Thangaraj, G.; Poongothai, E. On Fuzzy  $\Sigma$ -Baire Spaces, International Journal Of Fuzzy Math. And Sys., 2013, Volume 3(4), Pp.

- 275-283.
- [14] Thangaraj, G.; Poongothai, E., On Fuzzy Regular  $\Sigma$ -Baire Spaces, *Advances In Fuzzy Mathematics*, 2014, Volume 9(1), Pp. 95-104.
  - [15] Thangaraj, G.; Poongothai, E., On Fuzzy Pre- $\Sigma$ -Baire Spaces, *Advances In Fuzzy Mathematics*, 2016, Volume 11(1), Pp. 39-49.
  - [16] Turksen I., Interval-Valued Fuzzy Sets Based On Normal Form, *Fuzzy Sets And Systems*, Vol 20, 1986., 191-210.
  - [17] Salama A.A., And S.A.Alblowi, Neutrosophic Sets And Neutrosophic Topological Spaces, *IOSR Journal Of Mathematics*, Volume 3(4), 2012.
  - [18] Salama A.A., Generalized, Neutrosophic Set And Generalized Neutrosophic Topological Spaces, *Computer Science And Engineering*, Volume 2(7),LI 2012., 129-132.
  - [19] Smarandache Fuzzy, Neutrosophic Set, A Generalization Of The Intuitionistic Fuzzy Sets, *Inter.J.Pure Appl.Math.*, Volume 24, 2005., 287-297.
  - [20] Veereswari V., An Introduction To Fuzzy Neutrosophic Topological Spaces, *IJMA*, Volume 8(3), 2017., 144-149.
  - [21] Zadeh L.A., Fuzzy Sets, *Information And Control*, Volume 8, 1965, 338-353.