

Effect of Olive Oil Massage on Postoperative Cesarean Pain and Sleep Quality: a Randomized Controlled Trial

Dr. Rania Mahmoud Abdel Ghani¹, Dr. Amel Shaaban Abd Elmonem²

¹Assistant Professor of Maternal and Newborn Health Nursing, Faculty of Nursing, Cairo University-EGYPT

²Lecturer of Maternal and Newborn Health Nursing, Faculty of Nursing, Cairo University-EGYPT

Corresponding author: Dr. Rania Mahmoud Abdel Ghani

ABSTRACT: The aim of the study is to assess the effect of olive oil massage on post-operative cesarean pain and sleep quality. Design: randomized controlled trial. Sample: simple random sample of two hundred primiparous women was recruited in the study. The inclusion criteria was women who can read and write, their age ranged between 15-38 years, undergoing elective cesarean with spinal anesthesia, term pregnancy, have intact skin in the massage areas (i.e. scalp, neck, upper shoulder, hands and feet), no history of olive oil sensitivity, full consciousness after the surgery, willingness to participate, free from any medical disorders and post-cesarean section complications. Procedure: massage took place 8 hours post-cesarean section operation. Olive oil (5 ml) was applied for massaging each part of the woman's body (i.e. scalp, neck, upper shoulder, hands, and feet). Results: there were statistical significant differences between the study and the control groups related to levels of the post-operative cesarean section pain and fatigue after olive oil massage ($p \leq 0.05$). Further, there were statistical significance differences between both groups related to mean of total sleep disturbance ($t=-12.07$, $p \leq 0.0001$) and total sleep effectiveness ($t=15.77$, $p \leq 0.0001$). Conclusion: Olive oil massage may act as positive intervention in managing the post-operative cesarean pain and sleep disturbance.

Key Words: Olive Oil- Massage- Postoperative Cesarean Pain- Sleep Quality- RCT.

Date of Submission: 20-04-2018

Date of acceptance: 05-05-2018

I. Introduction

Cesarean section pain delays mothers' contact with their newborn, besides interferes to take proper breastfeeding position, self-care, newborn care and to carry out the essential daily activities (Abbaspoor, Akbari, and Najari, 2014). Further, after cesarean section, women usually experience worsening in their total sleep hours. Wound pain, uterine contraction pain and frequent breast feeding are leading factors of sleep complaints (Manjurai & Latheef, 2016). In order to resume the energy and engage with the routine daily activities, the main requirements are free from pain and adequate sleeping. Sleep adequacy improves tissue restoration and immune system functions (Sook, et al. 2013). The new modalities in medical intervention today are integrating the medical therapy with the alternative and complementary one. One of the important cost effective and applicable alternative therapy is massage. Massage affects different body systems and their functions positively. Recently, massage is defined as a type of aero-hydrotherapy. Massage helps to improve local and general circulation, promote immunological function, enhance natural healing ability and maintain homeostasis (Nelson, 2015). Many researches addressed the effect of massage alone, and with the essential oils for intervening pain and enhancing quality of sleep. But up till now, there are few scattered researches evaluated the effect of massage with olive oil. Olive oil is one of herbal products of its known history in the traditional medicine. Olive oil acts as pain killer, anti-inflammatory agent, besides its tranquilizing properties (Chauhan, Rani and Bansal, 2016). Nurse has an important and integral role in early post cesarean section intervention. The main role is helping post cesarean section women to restore their energy in order to adapt to new motherhood responsibilities.

AIM

To assess the effect of olive oil massage on post-operative cesarean pain and sleep quality

HYPOTHESES

1. Olive oil massages will minimize post-operative cesarean pain intensity of primiparous women.
2. Olive oil massages will improve sleep quality of post-cesarean primiparous women.

POSTOPERATIVE CAESAREAN PAIN

It is an acute pain as a result of surgical procedure, present on the day of the caesarean section and persistent for 18-24 hours. It is triggered by changes position of the supine to a sitting upright (Peter, etal. 2013).

II. MATERIALS AND METHODS

DESIGN

Randomized controlled trial

SETTING

The present study was conducted at El Kasr Aini, Cairo University maternity hospitals, at the postnatal unit.

SAMPLE


Simple random sample of two hundred primiparous women was recruited in the study. The inclusion criteria was women who can read and write, their age ranged between 15-38 years, undergoing elective cesarean with spinal anesthesia, term pregnancy, have intact skin in the massage areas (i.e. scalp, neck, upper shoulder, hands and feet), no history of olive oil sensitivity, full consciousness after the surgery, willingness to participate, free from any medical disorders and post- cesarean section complications. The sample size has been determined utilizing sample equation based on the daily numbers of cesarean section operations. Sample size was determined with a risk of type I error of 5% and a power of 80% .

Recruitment of the participants

In order to conduct correct randomization, randomization was done by the following three steps; *first*, identifying the total number of women who admitted in the operation room in one day from the operation's time table notes. *Second*, listing the mothers who met the inclusion criteria. *Third*, determining and ordering the hospital's numbers wrote on the women admission paper. Woman who had an odd number on her admission paper was included in the study. Further, sealed envelopes with numbers from one to two hundred were utilized for random assignment. Allocation concealment was approached through a trained professional researcher; figure (1) demonstrates women flow across the study.

ETHICAL CONSIDERATIONS

Written permission was obtained from the administrative authorities of maternity teaching hospital. The aim and the study procedure were explained. Consent was obtained. Confidentiality of data was considered, and participants were identified in the computerized database by numbers.



III. TOOLS AND MEASUREMENTS

1. **A structured interview schedule** includes socio-demographic and obstetrical data.
2. **Sleep Scale (Verran and Synder-Halpern, 1987):** it was designed to assess the sleep quality of the last 24 hours of hospitalized persons; those without pre-existing difficulties of sleep. The VSH assesses two main branches of sleep experience; 1) disturbance and 2) effectiveness for a total of 8 questions. Each question rated from zero to 10 (a total range of 0–80). High score indicates worst sleep for the sleep disturbance and best sleep for the sleep effectiveness subscale. The reliability coefficient of the scale was 0.82. Factors analysis and correlation showed validity.
3. **Visual Analogue Scale for pain (Huskisson, 1974):** it is a line of 10 cms marked from zero (no pain) to 10cms (worst pain). The respondent was requested to mark among the two extremes that best described the degree of pain intensity being experienced at that moment. The VAS possesses both concurrent validity and discriminate validity (Gift, 1989).
4. **The Fatigue Severity Scale (FSS):** it is likert scale of 7 points to assess the fatigue effects on daily activities. The scale consists of nine statements; each statement should be rated from one to seven and women were asked to check on the suitable number. A value of one indicates strong disagreement with the statement, while a value of seven indicates strong agreement. A total score of less than thirty six suggests no effect of fatigue on women activities. Moreover a score of thirty six or more indicates that further medical intervention is needed (Lauren, et al. 1989).
5. **Olive oil:** it is one of herbal products; it is composed of 98% triglycerides, which includes predominantly mono-unsaturated oleic acid which has been established to be vital for skin maintenance (Kapellakis, et al. 2008). In the present study, it was used without dilution or any additions. The researchers get the original olive oil from the pharmacists.

IV. INTERVENTIONS

Data collection procedure took place between September 2017 to February 2018. The present study was based on the gate control theory of Melzack and Wall (1965). They propose that pain impulses can be manipulated and inhibited through closing the gate. It hypothesized that massage might be compelling in shutting the gate. Massage interferes with the transmission of noxious stimuli and modifies the perception of pain (Adams, White and Beckett, 2010). Massage took place 8 hours post- cesarean section operation. In the study group, woman was helped to get out from bed. Woman was supported by a pillow behind the back and two stairs for legs (i.e. in order to sit in a proper, comfort position). Olive oil (5 ml) was applied for massaging each part of the woman's body (i.e. scalp, neck, upper shoulder, hands, and feet). Massage was applied by rotational friction movements, grasping each part of hands (i.e. from the wrist to fingers) and feet without focusing or pressuring on a certain point (20 minutes, five minutes for each part) using the fingers' palm. Friction was applied for neck and shoulder (for 10 minutes, five minutes for each). Additional five minutes scalp massage had been provided with gentle circular, upward and downward directions. The total amount of oil consumed in the massage session was approximately 35 mls. while the total duration of session was 35 minutes. The control group received the routine hospital care. The study and the control groups were assessed for pain intensity sleep quality and fatigue level. Pain intensity was assessed before massage and immediately after massage. Also, two times assessment of fatigue were done before massage and next day intervention. While sleep pattern was assessed once, next day intervention. Phone calls have been utilized for fatigue and sleep quality assessment for women who discharged before conducting the second interview.

V. STATISTICAL ANALYSIS

Statistical package for the social science (SPSS), version 20 was used for data analysis. An intention to treat analysis was utilized without any drop out.

Characteristics of mothers among the study and the control groups.

Regarding characteristics of sample, data denoted that, there were no statistical significant differences between both groups; the study and the control groups related to mean of age, mean of gestational age, educational levels and occupation. The age ranged between 15-38 years old. Further, most of the mothers were housewives and could read and write $p \geq 0.05$ (table 1).

Comparisons between both groups related to pain intensity and fatigue's level before and after massage

Table 2 revealed that there were positive statistical significant differences between both groups; the study and the control groups related to levels of the post-operative cesarean section pain and fatigue after olive oil massage

($p \leq 0.05$). Although, the fatigue levels before and after massage for both groups were considered accepted and far from any medical intervention.

Table 1. Characteristics of mothers among the study and the control groups.

Items	Study group (n=100)		Control group (n=100)		t	P
	Mean	SD	Mean	SD		
Age	31.18	3.04	31.06	2.94	0.28	0.77
Gestational age (weeks)	38.45	0.95	38.55	0.98	-0.76	0.44
	No.	%	No.	%	χ^2	P
Education level						
Read & Write	35	35.0	32	32.0	0.88	1.00
Primary School	29	29.0	27	27.0	0.96	1.00
Preparatory School	11	11.0	13	13.0	0.55	0.66
Secondary School	11	11.0	10	10.0	0.93	1.00
University	14	14.0	18	18.0	0.33	0.43
Occupation						
Housewife	65	65.0	56	56.0	0.43	0.46
Employee	35	35.0	44	44.0	0.09	0.10

Level of significance at $p \leq 0.05$.

Table 2. Comparisons between both groups related to pain intensity and fatigue's level before and after massage

Items	Study group (n=100)		Control group (n=100)		t	P
	Mean	SD	Mean	SD		
Before massage						
Pain intensity	4.38	0.48	4.27	0.44	1.66	0.09
The fatigue severity score	29.37	0.48	29.36	0.48	0.14	0.88
After massage						
Pain intensity	3.85	0.38	4.27	0.44	-7.11	0.0001
The fatigue severity score	29.04	0.71	29.28	0.45	-2.83	0.005

Level of significance at $p \leq 0.05$

Comparisons between both groups related to sleep quality reports.

Regarding mothers' reports on their sleep disturbances and effectiveness, there were statistical significant differences between both groups related to mean of total sleep disturbance ($t=-12.07$, $p \leq 0.0001$) and total sleep effectiveness ($t=15.77$, $p \leq 0.0001$). Further, there were positive statistical significant differences between both groups of the sub-items related to each domain. Data reflects inadequate sleep quality of the control group mothers. ($p \leq 0.0001$), (table 3).

Table 3. Comparisons between both groups related to sleep quality reports.

Items	Study group (n=100)		Control group (n=100)		t	P
	Mean	SD	Mean	SD		
Sleep Disturbance	14.51	0.79	15.68	0.51	-12.07	0.0001
Mid-sleep awakening	3.90	0.62	4.75	0.43	-11.22	0.0001
*Soundness of sleep	5.21	0.40	5.40	0.49	-3.02	0.003
Quality of disturbance	4.10	0.30	4.72	0.44	-11.60	0.0001
*Sleep latency	3.68	0.46	4.63	0.48	-14.16	0.0001
Sleep Effectiveness	21.37	0.48	20.29	0.45	15.77	0.0001
Total sleep period	5.56	0.49	4.64	0.54	12.39	0.0001
Rest upon awakening	6.54	0.50	6.33	0.47	2.98	0.003
Subjective quality of sleep	5.39	0.49	5.16	0.36	3.70	0.0001
*Sleep sufficiency evaluation	5.49	0.50	5.16	0.36	5.24	0.0001

Level of significance at $p \leq 0.05$

*Sleep latency is the amount of time between settling down to sleep and actually falling asleep

*Sleep sufficiency evaluation represents a person's perception of having received sufficient quantity of sleep.

*Soundness of sleep is in good condition of sleep

Baseline and post-intervention pain intensity & fatigue severity score among the study group

Regarding the differences of the pain intensity and fatigue levels among the study group before and after massage, results revealed that there were positive statistical significant differences with $p \leq 0.0001$ (table 4).

Table 4. Baseline and post-intervention pain intensity & fatigue severity score among the study group

Items	Study group (n=100)				t	P
	Mean	SD	Mean	SD		
	Baseline		After massage			
Pain intensity	4.38	0.48	3.85	0.38	6.99	0.0001
The fatigue severity score	29.37	0.48	29.04	0.71	5.04	0.0001

Level of significance at $p \leq 0.05$

VI. DISCUSSION

Methodological considerations

It was difficult to conduct double or single blinding in the present study, because the nature of the intervention.

Results discussion

Caesarean section (CS) usually initiates moderate to severe pain for about 48 hours after operation. It requires optimal pain management; which not only enable the mother to ambulate early but also crucial in the post-cesarean section operative rehabilitation (Bonnet et al. 2010). Further, after cesarean section, women being at obvious risk of sleep disturbance (Hunter, Rychnovsky & Yount, 2009). This study looking at determining the effect of olive oil massages on post-operative cesarean pain and sleep quality. The present study results revealed that there were statistical significant differences between both groups; the study and the control groups related to the levels of the post-operative cesarean section pain after olive oil massage, although the levels of pain among both groups were moderate in its' intensity. These results may interpret as; massage increase para-sympathetic nerve activity resulting in an increase in the secretion of epinephrine and a decrease in the secretion of nor-epinephrine and cortisol. In addition, olive oil has analgesic and anti-inflammatory properties; olive oil as morphine induced marked analgesic effect (Riella, et al. 2012), further it inhibits releasing of cyclo-oxygenases enzymewhich is involved in prostaglandins as a result, inducing an anti-inflammatory effect (Babu, et al 2009).The present results goes on the same line with Kuriakose (2012) and Chauhan, Rani and Bansal (2016) who assessed the effectiveness of olive oil back massage on labor pain and concluded that olive oil massage may be able to modify the pain intensity in the active phase and improve patients' satisfaction. And with Degirmen, et al. (2010) who stated that, massaging of the hand and foot could be considered as an effective nursing management of post-operative cesarean pain.

In terms of women report of their sleep quality, results of the present study revealed that olive oil massage of scalp, neck, upper shoulder, hands and feet was effective in improving sleep disturbance in term of sleep latency, mid-sleep awakenings, and quality of disturbance and soundness of sleep. The neck muscles massage extends the range of cervical spine rotation (Gong, et al. 2012) and relaxes the tension of neck flexor muscles (Gong, 2013). In addition, scalp massage has its effect on inhibiting the stress hormones (epinephrine and nor-epinephrine) and as a result, it initiates the activation of para-sympathetic nerves such as decrease in heart rate, relaxation of blood vessels and muscle (Hong, Youngand Wan, 2016). Thus, massage has been proved to produce its positive effects on hormones (Morhenn, Beavin and Zak, 2012), blood pressure (Nelso, 2015), and heart rate (Supa'at, et al. 2013). Further, foot massage improved the quality of sleep and increase the nocturnal sleep duration (Ariamanesh, Malekshahi and Safari, 2015).While hand massage has its positive effect on fatigue and sleep quality (Park, Chun and Kwak, 2016). The present results go parallel with the results of Manjurai, et al. (2016) study, they reported that initiate foot and back massage early after cesarean section operation enhances women's sleep quality, and with Adams, White and Beckett (2010), who evaluated the effect of massage on pain in different occasions and specialties; in medical, surgical, and obstetrics units, they reported that, more than fifty percent of their sample had pain relief and improvement in their sleep pattern.

Moreover, postoperative cesarean section fatigue may be a result of many interrelated factors; pain, sleep disturbance, feeding the newborn and the limited women movement abilities first hours post-operation. The present study's result revealed that, there was statistical significant difference between both groups related to the post-operative cesarean section fatigue. This result may be based on the proposed effect of

massage on women sleep quality; as a result the level of fatigue was manipulated. Further, when woman gets adequate sleep, she can resume her energy to practice her motherhood role and activities. Many studies approved the effect of massage on relieving pain and reducing the symptoms of fatigue in different health conditions (Nunes, etal. 2016; Mori, etal. 2004; and Tanaka, etal. 2002). While recent study of Park (2016) correlated with massage, fatigue and sleep among hospice patients, and his results go on the same line with the findings of the present study.

CONCLUSIONS

Olive oil massage may act as positive intervention in managing the post-operative cesarean pain and sleep disturbance.

RECOMMENDATIONS

The long term effect of olive oil massage on postpartum woman sleep, fatigue and daily activities is needed to be examined in the future studies.

IMPLICATIONS FOR PRACTICE

Olive oil massage is a simple, inexpensive intervention that can be initiated post-caesarean section in a short time with no harm.

CONFLICT OF INTEREST

There is no conflict of interest.

REFERENCES

- [1]. Abbaspoor, Z.Akbari, M. and Najar, S. (2014). [Effect of Foot and Hand Massage in Post-Cesarean Section Pain Control: A Randomized Control Trial]. American Society for Pain Management Nursing. 15 (1):132-136.
- [2]. Adams, R. White, B. and Beckett. (2010).The Effects of Massage Therapy on Pain Management in the Acute Care Setting. Int J Ther. Massage Bodywork. 3(1): 4-11.
- [3]. Ariamanesh, F. Malekshahi, F. and Safari, M. (2015). The Effect of Foot Massage on Night Sleep Quality in Hemodialysis Patients. Yafte. 17(1):71-7.
- [4]. Babu, N. P. Pandikumar, P. Ignacimuthu, S. (2009). Anti-inflammatory Activity of Albizialebeck Benth, an Ethnomedicinal Plant, in Acute and Chronic Animal Models of Inflammation. Journal of Ethnopharmacology. 125 (2): 356-360
- [5]. Bonnet, M. P. Mignon, A. Mazoit, J. X. Ozier, Y. and Marret, E. (2010). [Analgesic Effect and Adverse Effects of Epidural Morphine Compared to Parenteral Opioids after Elective Caesarean Section: A Systemic Review]. European Journal of Pain". 14 (9): 894-899.
- [6]. Chauhan, K. Rani, S. Bansal, P. (2016). Effectiveness of Olive Oil Back Massage on Reduction of Labor Pain during First Stage of Labor. Int J Nurs Midwif Res. 3 (2&3): 32-36. ISSN: 2455-9318.
- [7]. Degirmen, N. Ozerdogan, N. Sayiner, D. Kosgeroglu, N. and Ayranci U. (2010). [Effectiveness of Foot and Hand Massage in Post cesarean Pain Control in a Group of Turkish Pregnant Women]. Appl Nurs Res. 23 (3):153-8.
- [8]. Gift, A. G. (1989). Visual Analogue Scales: Measurement of Subjective Phenomena. Nurs Res. 38: 286-288.
- [9]. Gong, W. (2013). Impact of Longus Colli Muscle Massage on the Strength and Endurance of the Deep Neck Flexor Muscle of Adults. J Phys Ther Sci. 25: 591-593.
- [10]. Gong, W.T. Park, G. D. and Lee, J. H. (2012). Effects of Longus Colli Muscle Massage on the Cervical Extension Range of Motion of Adults. J Phys Ther Sci. 24: 999-1001.
- [11]. Hong, I. K. Young, T. K. and Wan, K.Y. (2016).The Effect of a Scalp Massage on Stress Hormone, Blood Pressure, and Heart Rate of Healthy Female. J Phys Ther Sci. 28 (10): 2703-2707.
- [12]. Hunter, L. P. Rychnovsky, J. D. and Yount, S. M. (2009). A Selective Review of Maternal Sleep Characteristics in the Postpartum Period. J Obstet Gynecol Neonatal Nurs 38: 60-68.
- [13]. Huskisson, E. C. (1974). Measurement of Pain. Lancet. 9:1127-1131.
- [14]. Kapellakis, J. E. Tsagarakis, K. P. and Crowther, J.C. (2008). Olive Oil History, Production and by-Product Management. Rev Environ Sci Biotechnol. 7(1):1-26.
- [15]. Kuriakose, D. (2012).A Study to Evaluate The Effectiveness of Olive Oil Back Massage on Labor Pain and Progress in First Stage of Labor among Primipara Mothers. Indian Journal of Nursing and Midwifery Research. 88-90.
- [16]. Lauren, B. Krupp, M. D. Nicholas, G. L. Muir-Nash, J. and Alfred, D. S.(1989).The Fatigue Severities Scale Application to Patients with Multiple Sclerosis and Systemic Lupus Erythematosus. Arch Neurol. 46 (10):1121-1123.
- [17]. Manjurai, A. E. and Latheef, F. (2016). Effectiveness of Foot Reflexology and Back Massage in Improving the Quality of Sleep among Post Caesarean Mothers. Int. J. Res. Ayurveda. Pharm.7 (5): 103-108.
- [18]. Melzack, R. and Wall, P. (1965). Pain Mechanisms: a New Theory. Science. 150 (3699): 971-8.
- [19]. Morhenn, V. Beavin, L. E. and Zak, P. (2012). J. Massage Increases Oxytocin and Reduces Adrenocorticotropin Hormone in Humans. Altern Ther Health Med. 18: 11-18.
- [20]. Mori, H. Tanaka, T. H. Ohsawa, H. and Nishijo, K. (2004). [Effect of Massage on Blood Flow and Muscle Fatigue Following Isometric Lumbar Exercise]. International Medical Journal of Experimental and Clinical Research. 10(5): CR173-8.
- [21]. Nelson, N.L. (2015). Massage Therapy: Understanding the Mechanisms of Action on Blood Pressure. A Scoping Review. J Am Soc Hypertens. 9: 785-793.
- [23]. Nunes, G. S. Urio Bender, P. Menezes, F. S. Yamashitafuji, I. Vargas, V. Z. and Wageck, B. (2016). Massage Therapy Decreases Pain and Perceived Fatigue after Long-Distance Ironman Triathlon: a Randomized Trial. J Physiotherapy. 62 (2): 83-87.
- [24]. Park, H. Chun, Y. and Kwak, S. (2016). [The Effects of Aroma Hand Massage on Fatigue and Sleeping among Hospice Patients]. Open Journal of Nursing. 6: 515-523.
- [25]. Peter, H. Pan, M.S.E.E. Ashley, M. Tonidandel, M. D. and Carol, A.(2013). Predicting Acute Pain after Cesarean Delivery Using

- Three Simple Questions. *Anesthesiology*. 118: 1170-1179.
- [26]. Riella, K. R. Marinho, R. R. and Santos, J. S. et al. (2012). Anti-Inflammatory and Cicatrizing Activities of Thymol, a Monoterpene of the Essential Oil from *Lippia Gracilis*, in Rodents. *Journal of Ethnopharmacology*. 143(2):656–663.
- [27]. Verran, J. A. and Snyder-Halpern, R. (1987). Instrumentation to Describe Subjective Sleep Characteristics in Healthy Subjects. *Res. Nurs. Health*. 10:155–163.
- [28]. Sook, M. J. Lee, S. Bae, I. Hur, M. Seong, K and Lee, M., S. (2013). [Effects of Aroma Massage on Home Blood Pressure, Ambulatory Blood Pressure, and Sleep Quality in Middle-Aged Women with Hypertension]. *Evid Based Complement Alternat Med*. doi: 10.1155/2013/403251.
- [29]. Supa'at, I. Zakaria, Z. and Maskon, O. et al. (2013). Effects of Swedish Massage Therapy on Blood Pressure, Heart Rate, and Inflammatory Markers in Hypertensive Women. *Evid Based Complement Alternat Med*. 171852.
- [30]. Tanaka, T. H. Leisman, G. Mori, H. and Nishijo, K. (2002). [The Effect of Massage on Localized Lumbar Muscle Fatigue]. *BMC Complementary and Alternative Medicine*. 2(1):9. doi: 10.1186/1472-6882-2-9.

Dr. Rania Mahmoud Abdel Ghani "Effect of Olive Oil Massage on Postoperative Cesarean Pain and Sleep Quality: A Randomized Controlled Trial "IOSR Journal of Nursing and Health Science (IOSR-JNHS) , vol. 7, no.2 , 2018, pp. 92-98