

Prevalent and Multiple Antibiotic Resistance Patterns of Salmonella Serovars Isolated From Blood Samples of Hospitalized Patients in Kano, North-West, Nigeria.

¹A. Mas'ud, ²Musa Tijjani

¹Department of Medical Laboratory Science, School of Health Technology, Kano, Kano state, Nigeria

²Department of Microbiology, Ahmadu Bello University, Zaria, Kaduna State, Nigeria.

Abstract:

Aim: The aims of the study were to determine the prevalent serovars of Salmonella and the multiple antibiotic resistance patterns of Salmonella isolates recovered from the blood samples of hospitalized patients in Kano metropolis.

Study design: The study is a descriptive cross-sectional study.

Place and duration of study: One milliliter of venous blood was collected from each patient with some or all clinical features of salmonellosis that sign a consent form and transfer into EDTA bottles. If daily is unavoidable blood samples were stored at 4 °C. Samples were analyzed at the both laboratories of the authors. This work was carried out between May, 2012 and March, 2014.

Methodology: The blood specimens were cultured in thioglycollate broth and sub-cultured onto deoxycholate citrate agar (DCA), Salmonella-Shigella agar (SSA) and brilliant Green agar (BGA) followed by confirmation of presumptive colonies using different biochemical tests and analytical profile index 20E. Serologic identification of Salmonella was performed by slide agglutination test using polyvalent O and H Salmonella antisera. Antibiotic susceptibility studies were performed by the disc diffusion method using ampicillin, chloramphenicol, ciprofloxacin, nalidixic acid and Trimethoprim-sulfamethoxazole.

Results: Although, the relationship between different age groups was not significantly associated ($P < 0.05$), patients under age bracket of 21-30 years were found to be more susceptible to Salmonella infections with 34 representing 6.8% followed in that order by 11-20 years (28), 31-40 years (19), ≤ 10 years (15) and > 40 years (8) age groups, representing 5.6%, 3.6%, 3.0% and 1.6% respectively. The prevalent rate of Salmonella infections was significantly higher ($P > 0.05$) in males than the females patients with 68 (13.7%) and 36 (7.2%) respectively. The serovars of Salmonella Typhi was the most predominant with 39(7.8%) followed by Salmonella Typhimurium 35(7.0%), Salmonella Paratyphi C 14(2.8%), Salmonella Paratyphi B 11 (2.2%) and Salmonella Paratyphi A was the least prevalent serovar with 5 (1.0%). Of the 104 isolates 96 (92.3%) were Of the 104 isolates 96 (92.3%) were resistant to Ampicillin; 81 (77.9%) resisted to Nalidixic acid; 30 (30.8%) resisted to Chloramphenicol, 17(16.3%) resisted to Cotrimazole while none (1.0%) resisted to Ciprofloxacin. 37 isolates were found resistant to 1 of 3 first-line anti-Salmonella antibiotics (ampicillin, cotrimoxazole and chloramphenicol). Highest MARI was recorded from thirty seven (44.3%) isolates with MARI of 7.8 with two phenotypic patterns which differed in two different antibiotics (ampicillin and nalidixic acid).

Conclusion: The frequency of Salmonella infections was highest among 21-30 year age group lowest in ≥ 40 year age group. However the rates of infection among all the six (6) age groups were not significantly associated. The prevalent rate of Salmonella infections was significantly higher ($P > 0.05$) in males than the females' patients. However, Salmonella Typhi was predominant followed by Salmonella Typhimurium, Salmonella Paratyphi C, B and A. Most Salmonella serovars isolated from patients in Kano, Nigeria resisted to Ampicillin. They also resisted to Nalidixic acid, Chloramphenicol and Cotrimazole, in decreasing order. Ciprofloxacin remained effective against all the Salmonella isolates tested. 37 isolates were found resistant to 1 of 3 first-line anti-Salmonella antibiotics (ampicillin, cotrimoxazole and chloramphenicol).

Key words: Salmonella serovars; blood samples; multiple antibiotic resistance; Kano-Nigeria.

I. Introduction

Salmonella infection also known as salmonellosis is an infection caused by ingesting Salmonella in food that is contaminated by faeces of animals or humans directly or indirectly. Common sources of infection include poultry meat and meat products, eggs and egg products. Some of the symptoms of salmonellosis are diarrhoea, vomiting fever and abdominal pain these occur 12-36 hours after eating infected food, in acute infection, blood and mucous are present in faecal specimens [1,2]. Salmonellae can be isolated from blood, stool, urine, bone marrow, duodenal aspirates and rose spots. The infective dose of Salmonella is usually about 1000 colony forming units (CFU) but it depends on various factors including the serotype and the pH of the

stomach. If the pH of the stomach is high a low dose is sufficient to cause infection. A high stomach pH can also accelerate Salmonella invasion and allow the organism to spread into the circulatory system, liver and gall bladder causing further complications [3].

Salmonella, a primary inhabitant of the gastrointestinal tract, is recognized as one of the most common causes of food borne infection worldwide, resulting in millions of infections and significant human death annually [4]. Non-typhoidal salmonellosis is common in most parts of the world [5]. It is widely spread in Europe and North America [6,7], Latin America, the Middle East and Africa [8]; also in countries such as India [9], Japan [10] and the United States [11]. Several studies have documented isolation of non-typhoidal Salmonella from humans and poultry in different parts of Nigeria [12,13,14,15]. Outbreaks of salmonellosis caused by Salmonella serovars Gallinarum, Pullorum, Typhimurium, and Enteritidis have also been reported [13,15,16].

Laboratory diagnosis of typhoid fever is dependent upon either the isolation of Salmonella Typhi from clinical samples or detection of raised antibody titres as shown by agglutination tests. Subsequently, antibody levels > 1:160 for Salmonella Typhi and Salmonella Paratyphi are regarded as diagnostic of enteric fevers [17]. Salmonellae can be isolated from blood, stool, urine, bone marrow, duodenal aspirates and rose spots. The organisms can usually be detected in 75-90% of patients during the first ten days of infection and in about 30% of patients during the third week in the blood [18].

There are rampant indiscriminate prescriptions of antibiotics for the treatment of Salmonella infection and sales of expired, fakes and substandard antibiotics. This has been an important problem in the treatment and susceptibility of Salmonella serovars to antibiotics. Antibiotics use increases the spread of antibiotics resistant bacteria. This is called 'selective pressure'. As bacteria become resistant to increasing number of antibiotics, the remaining antibiotics are used more often; increasing the selective pressure of Salmonella and therefore, making them more resistant [19,20]. In Nigeria most typhoid endemic areas, culture facilities are often available but have not been used for the routine laboratory diagnosis of enteric fever and blood cultures may be negative in patients who have taken antibiotics prior to the test [21].

However, majority of antibiotic resistance studies on Salmonella in the study area are based on the use of isolates obtained from stool samples of patients with Salmonella infection admitted into hospitals. Recent studies indicate that antibiotic resistance genes are a growing problem indicating the need to pay closer attention on the isolation of Salmonella in blood and community acquisition of multiple antibiotic resistance in Salmonella [22,23].

II. Materials And Methods

2.1 Hospitals

The six most patronized hospitals were randomly selected including one Teaching Hospital (Aminu Kano Teaching Hospital), three specialist hospitals (Murtala Mohammed Specialist, Mohammed Abdullahi Wase Specialist and Sir Sunusi Specialist Hospital), one General Hospital (Sheik Waziru Gidado General Hospital) and one Private Hospital (Khadijat Memorial Private Hospital). All are situated within Kano metropolis. The selected hospitals are reference hospitals in the state where people from various parts of the state and neighboring states of various occupations attend. They gave more than 70% of health care delivery in the state at large.

2.2 Patients and Specimens

Patients (in and out) who patronized the six selected hospitals with some or all clinical symptoms of Salmonella infections (i.e. vomiting, diarrhoea, headache, abdominal pain, body ache, breathlessness, weight lost, constipation and anaemia) recruited to sign the consent form were used for the study.

Any patient (in and out) who brought his blood specimen to the laboratory reception of one of the six selected hospitals for widel test, malarial test and other related blood tests recruited to sign the consent form were used for the study.

Blood (1ml) collected from each patient diagnosed positive for salmonellosis was used as sample for the study.

2.3 Collection and Handling of Specimens

One milliliter of venous blood was obtained using sterile syringe from an antecubital vein of each patient recruited for the study and dispensed immediately into 10ml thioglycollate broth. Sterile bijoux bottle that contained blood and 10ml thioglycollate broth was then labeled with specimen number, type of medium and date of dispensing [18].

2.4 Isolation and identification of salmonellae

2.4.1; Presumptive isolation of Salmonella

One milliliter of venous blood specimen was dispensed into 10ml thioglycollate broth and sub-cultured onto SSA, BGA and DCA agar everyday and incubated aerobically at 37°C for 7 days [22]. The cultured plates, SSA, BGA and DCA agar were examined for the presence of typical colonies of Salmonella based on cultural and morphological characteristics, that is, transparent colonies with black centre on SSA and pink colonies surrounded by a red medium on BGA, and small red translucent and or dome-shaped colonies, which may have central black spot due to hydrogen sulphide production [18].

Bacterial isolates obtained were further sub-cultured by stabbing into nutrient agar slants and stored at 4°C after aerobic incubation 37°C for 24 hours for subsequent analysis.

2.4.2: Purification of isolates

Presumptive culture of Salmonella stored in nutrient agar slant was sub-cultured onto SSA aerobic incubation 37°C for 24 hours to observe for the colonial characteristics of Salmonella and isolation of pure culture for subsequent biochemical characterizations.

2.4.3: Biochemical characterization of Salmonella

Isolation and identification of organisms were carried out as described by ISO [24], Habtamu et al. [16], and OIE [25]. A 24 h pure culture of each isolate was used to determine their gram stain reaction. The following biochemical tests were carried out: Indole test, triple sugar iron test, citrate test, methyl-red test, Voges-Proskauer test, lysine decarboxylase test, ornithine decarboxylase test, urease test, sugar (trehalose, sucrose, inositol, glucose, dulcitol, maltose, mannitol, melibiose, salicin, rhamnose and arabinose) fermentation test and motility test. Isolates were further characterized using commercially available identification system-Analytical Profile Index (API) 20 E test kit (Biomérieux, France) following the manufacturer's guideline.

2.5 Sero-typing of the isolates

Serological identifications of presumptive Salmonella were performed by slide agglutination test. Presumptive isolates of Salmonella obtained from the series of biochemical tests were screened serologically with somatic O Salmonella Paratyphi A, B, C₁, Salmonella Typhimurium C₂ and Salmonella Typhi D.

An agglutination test was performed on a clean glass slide. The slide was divided into sections with a wax pencil and one small drop of physiological saline was placed in each test section on the slide. By using a sterile inoculating loop a portion of growth from the surface of TSI agar was removed and emulsified in each drop of physiological saline on the slide. It was then mixed thoroughly to create a moderately milky suspension. A bent inoculating loop was used to pick a small drop of antiserum and transferred to one of the suspensions; the second suspension served as the control (usually approximately equal volume of antiserum and growth suspension was mixed). The suspension and antiserum were mixed very well and then the slide was tilted back and forth to observe for auto-agglutination (agglutination is more visible if the slides is observed under a bright light and over a black background) [26].

If clumping appeared within 30 to 60 seconds the reaction is positive, the saline suspension (control) was examined carefully to ensure that it is even and does not show clumping resulting from auto agglutination. If auto-agglutination occurs, the culture is termed "rough" and cannot be serotype. When positive agglutination reaction was obtained in one of the antisera, the Salmonella serovars Paratyphi A, B, C₁, Typhimurium C₂ or Typhi D subgroup was confirmed, and no further testing with antisera needed to be conducted [26].

2.6 Test of antibiotic sensitivity of the Salmonella isolates.

In-vitro susceptibility of Salmonella isolates to various routine antibiotics was tested by the standard disc diffusion technique [1].

2.6.1 Standardization of inoculum

This was done as described by CLSI [27]. Pure culture of identified Salmonella isolate (s) from an 18-hour plate culture was selected. Sterile wire loop was used to pick 3 colonies of each Salmonella serotype and emulsified in 5 ml of sterile normal saline. The tube containing the bacterial suspension was inserted into a sensititre nephelometer (TREK Diagnostic systems, UK) after calibration. Adjustment was made with extra inoculum or diluents, if necessary, until 0.5 McFarland standards were obtained. Fifty microliter of the broth was further transferred into 5 ml of Mueller-Hinton broth (Oxoid, UK) in a tube [27].

2.6.2 Inoculation of test plates

Optimally, within 10 minutes after adjusting the turbidity of the inoculum suspension, a sterile cotton swab was dipped into the standardized suspension in Mueller-Hinton broth. The dried surface of a 20 ml

Mueller-Hinton agar plate in a 100 mm disposable plate (STERILIN, UK) was inoculated by streaking with the cotton swab over the entire sterile agar surface. The inoculated plates were air dried at 37°C to allow for any excess surface moisture to be absorbed before applying the antibiotic discs [27].

2.6.3 Application of discs to inoculated agar plates

All positive cultures of *Salmonella* serovars isolated from blood samples were tested in vitro for susceptibility to different antibiotic by agar diffusion technique as described by Kirby-Bauer [28] and WHO [29]. This was carried out according to WHO protocol [30]. The susceptibility testing of *Salmonella* isolates were carried out using Mueller Hinton agar and were tested in vitro for susceptibility to five different antibiotics; ampicillin (25µg), chloramphenicol (30µg), ciprofloxacin (25µg), nalidixic acid (30µg) and Trimethoprim-sulfamethoxazole (30µg) [31].

The inoculated plates were air dried under aseptic condition to eliminate the liquid on the surface of medium, sterile forceps was used to place the antibiotic discs on the inoculated plates. Within 30 minutes after applying the disc, the plate was inverted and incubated at 35°C for 18 hours. Meter ruler was then used on the underside of plate to measure the diameter of each zone of inhibition in millimeter. Zone diameter for ATCC 25922 was compared with NCCLS Published Limits; Interpretative chart was then used to interpret the zone sizes of Inhibition [31].

Results were recorded as susceptible, intermediate susceptible or resistant based on the zones size of each antibacterial disc used [30, 31].

2.9 Statistical analysis of results

Statistical Package for Social Science (SPSS) version 14 was used [32]. Descriptive statistics were used to categorical (frequency percentages) variables. Chi-square test analysis was used to determine association between the resistant rate of *Salmonella* isolates and antibiotics activities.

III. Results

3.1 Bacterial isolation.

Of the five hundred blood specimens sampled from six selected hospitals studied, total of 126 bacterial isolates and 104 *Salmonella* positive specimens were recorded: 110 were collected from Murtala Mohammed Specialist Hospital (35 bacterial isolates and 5.8% *Salmonella* positive specimens were obtained), 100 from Aminu Kano Teaching Hospital (25 bacterial isolates and 4.4% *Salmonella* positive specimens were obtained), 90 from Mohammed Abdullahi Wase Specialist Hospital (20 bacterial isolates and 3.4% *Salmonella* positive specimens were obtained), 80 from Sir Sunusi Specialist Hospital (17 bacterial isolates and 2.6% *Salmonella* positive specimens were obtained), 60 from Sheik Waziru Gidado General Hospital (15 bacterial isolates and 2.2% *Salmonella* positive specimens were obtained) and 60 from Khadijat Memorial Private Hospital (14 bacterial isolates and 2.4% *Salmonella* positive specimens were obtained).

3.2 Salmonella identification by biochemical characterization.

Out of one hundred and twenty six (25.2%) bacterial isolates obtained from six selected hospitals studied, One hundred and eighteen (23.6%) presumptive *Salmonella* isolates were obtained from various biochemical characterization and identification test.

3.3 Sero-typing of the Salmonella isolates.

One hundred and four (20.8%) *Salmonella* isolates were obtained after serologic identifications of presumptive *Salmonella* isolates were performed by slide agglutination test.

3.4 The distribution of Salmonella infections in relation to age and sex

Although, the relationship between different age groups was not significantly associated ($P < 0.05$), patients under age bracket of 21-30 years were found to be more susceptible to *Salmonella* infections with 34 representing 6.8% followed in that order by 11-20 years (28), 31-40 years (19), ≤ 10 years (15) and >40 years (8) age groups, representing 5.6%, 3.6%, 3.0% and 1.6% respectively (Table 1).

Table 1: The distribution of Salmonella infections in relation to age among hospitalized patients in Kano metropolis, Nigeria

Age group (Years)	No. examined	No. (%) infected
≤10	49	15(3.0)
11-20	132	28(5.6)
21-30	167	34(6.8)
31-40	93	19(3.8)
≥40	67	8(1.6)
Total	500	104(20.8)

KEY: ≥ = Greater than or equal to; ≤ = Less than or equal to; % = Percentage of total number of specimen tested (500).

When the 104 infected patients were assessed in relation to sex, the prevalence rate of Salmonella infections was significantly higher ($P > 0.05$) in males than the females patients with 68 (13.6%) and 36 (7.2%) respectively (Table 2).

Table 2: The distribution of Salmonella infections in relation to sex among hospitalized patients in Kano metropolis, Nigeria

Sex	No. infected (%)	No. not infected (%)	Total (%)
Males	68(13.6)	201(40.2)	269(53.8)
Females	36(7.2)	195(39.0)	231(46.2)
Total	104(20.8)	396(79.2)	500(100)

Fisher’s exact test two sides P value = 0.0081

KEY: NO. = Number; % = Percentage of total number of specimens tested (500).

3.5: The prevalence serovars of Salmonella

From Figure 1 out of 104 presumptive Salmonella isolates obtained in this study, Salmonella Typhi was the predominant serovar with 39(7.8%) followed by Salmonella Typhimurium 35(7.0%), Salmonella Paratyphi C 14(2.8%), Salmonella Paratyphi B 11(2.2%) and Salmonella Paratyphi A was the least prevalent serovar with 5(1.0%) prevalence rate. However, the relationship between each Salmonella serovar obtained was also statistically significant ($P > 0.05$).




Figure 1: The prevalence serovars of Salmonella from blood specimens of hospitalized patients in Kano metropolis, Nigeria.

t= 3.059 P value = 0.0377

KEY: S. = Salmonella; No. = Number; % = Percentage of total number of Salmonella isolated (104).

3.6 Antibiotic sensitivity of the Salmonella isolates.

Out of 104 Salmonella serovars isolated, 37 were triple, 51 double, and 14 single-antibiotic resistant phenotypes. Sixty five (62.5%) of the isolates displayed resistant to at least one or two antibiotics and thirty seven (35.6%) displayed resistant to three of antibiotics tested. However, the remaining two (1.9%) isolates did not display resistance to any of the five (5) antibiotics tested (Table 3).

Similarly, 37 isolates were found resistant to 1 of 3 first-line anti-Salmonella antibiotics (ampicillin, cotrimoxazole and chloramphenicol). In addition, 51 isolates were found resistant to 2 of 3 first-line anti-Salmonella antibiotics (ampicillin, cotrimoxazole and chloramphenicol), none of the isolates were resistant to all three first-line anti-Salmonella antibiotics (multi- antibiotics resistant salmonellae) (Table 3).

Table 3: Multiple antibiotic resistance patterns of Salmonella serovars isolated from blood specimens of hospitalized patients in Kano metropolis, Nigeria.

Single Resistance types	Number with type	Multiple antibiotic resistance patterns/phenotypes		
		Number of antibiotics	Number of isolates with pattern	Resistance Patterns/phenotypes
AMP.	13	2	39	AMP., NA.
CPX	1		7	AMP., COT.
			5	CH., NA.
		3	27	AMP., CH., NA.
			10	AMP., COT., NA.

KEY: AMP = Ampicillin; CH = Chloramphenicol; CPX = Ciprofloxacin; COT = Cotrimoxazole; NA= Nalidixic Acid

In this work, eighty eight (88) of the test Salmonella serovars exhibited multiple antibiotic resistances with five (5) different phenotypic resistant patterns. Thirty nine (44.3%) isolates have a MARI of 7.8 with two phenotypic patterns which differed in two different antibiotics (ampicillin and nalidixic acid). Twenty seven (30.7%) isolates have a MARI of 5.4 with three phenotypic patterns which also differed in three different antibiotics (ampicillin, chloramphenicol and nalidixic acid). Ten (11.4%) isolates have a MARI of 2.0 with three phenotypic patterns which differed in three different antibiotics (ampicillin, cotrimoxazole and nalidixic acid). Seven (8.0%) isolates have a MARI of 1.4 with two phenotypic patterns which also differed in two different antibiotics (ampicillin and cotrimoxazole). However, five (5.7%) has a MARI of 1.0 with also two phenotypic patterns which also differed in two different antibiotics (chloramphenicol and nalidixic acid) (Figure 4).




Figure 4: Multiple Antibiotic Resistances Indices (MARI) of Salmonella isolates from Kano metropolis, Nigeria.

The result of in vitro -antibiotic susceptibility testing demonstrated that of the five (5) antibiotics tested, most of the 104 isolates (92.3%) was significantly ($P > 0.05$) resistant to ampicillin. They were also resistant to nalidixic acid, chloramphenicol, and cotrimoxazole in decreasing order with 77.9%, 30.8%, and 16.3% resistance rate respectively. Ciprofloxacin remained effective against all the Salmonella isolates tested with exception of one (1) Salmonella isolate representing 1.0% resistance rate.

Table 5: Disc diffusion test on Salmonella serovars isolated from blood samples of hospitalized patients in Kano metropolis, Nigeria.

Antibiotic	Disc potency (μg)	No. (%) of Resistant Isolates n= 104
AMP	25	96(92.3)
CH	30	32(30.8)
COT	25	17(16.3)
CPX	30	1(1.0)
NA	30	81(77.9)

$X^2 = 11.03$ P value = 0.001

No. = Number; n= Number of isolates tested; S = Salmonella; % = Percent; μg = microgram; AMP = Ampicillin; CH = Chloramphenicol; CPX = Ciprofloxacin; COT = Cotrimoxazole; NA = Nalidixic Acid; % = Percentage of total number of each Salmonella serovar tested.

IV. Discussion

In the present study, the highest infections were recorded from 21-30 years age groups and least infection was recorded from patients with ≥ 40 year age group. However the rate of infection among all the six (6) age groups was not significantly associated. In this study, male were more infected than their female counter part in the same age groups. In addition, the rates of infections between males and females patients were statistically significant ($P > 0.05$). This work is in consonance with the findings of Adkins and Santiago [33]; Abdullahi [34] and Mashi [35].

The highest incidence in males of 21-30 years age group patients recorded in this study is probably because most of the males in the study area were more prone to contaminations than their female counter part of the same age group. Males usually eat and drink outdoors and they do not recognize the state of the food or drink they eat and the nature of the environment in which the food and drink are prepared [35]. It was further observed that the high prevalence of Salmonella infection in males of 21-30 years age group is connected with water exposure by these individuals in their community, who are supposed to be demographically and economically active and productive individuals in the communities, because they are more exposed to occupational hazard of farming, related water contact activities, contaminated environment, food and drink than children and elderly persons (Adkins and Santiago, 2006). Similarly, sharing of public toilet in school, market, bus stop by male would probably increase the level of infection among the male [34].

However, the low community awareness level of males of this age on the routes of Salmonella infection may increase the transmission potential of the disease since individuals would not care to protect them or take other cautions when exposure becomes necessary. It was further observed that even among informed males awareness level does not play important role in discouraging water activities, and practices have become necessary as they are the major means of subsistence; in term of irrigation, agriculture, gardening, outdoors business and other occupational hazards [36,37].

Females are less likely to be found eating, drinking and defaecating outdoors. This is because of culture and religious inclination. Moreover, females are less exposed to occupational hazard of farming, related water contact activities, contaminated environment, contaminated food and drink than their males' counterpart [36]. It was observed that males were responsible for 98% of activities involving contamination and water exposure [35].

On the other hand, adults (≥ 40 year age group) appreciate the quality of health they have, they are wise, matured and therefore, protect and maintain most of their valuable physical health. In addition, the adults do not usually predispose themselves to various contaminated areas than youth [35]. Furthermore, the presence of little or no infection in older individuals may be connected with the less water contact activities thus reducing the risk

of contracting infection. Another inference is the possibility of developed immunity by the older individuals who might have contracted the disease in their young age. Therefore, it can be concluded that such insusceptibility could reflect on age dependent acquired immunity [38].

Various studies have been conducted by many researchers in different parts of the world establishing the significance of Salmonella serovars in the causation of Salmonella infection. Five (5) serovars of Salmonella were encountered in the present study. In this study, Salmonella Typhi was predominant followed by Salmonella Typhimurium, Salmonella Paratyphi C, B and A. This finding is in consonant with studies by Asma et al. [39] and Akinyemi et al. [40]. This is contrary to the work conducted in Kano metropolis (the study area) by Abdullahi [34] who revealed that Salmonella Typhimurium was the most predominant serovar. The highest susceptibility to Salmonella Typhi in the study area could probably be because Salmonella infections occurs by ingesting organisms in contaminated food or water from contaminated hands (Salmonella Typhi is mainly water-borne, Salmonella Typhimurium and Salmonella Paratyphi are mainly food-borne). In the area where this research was conducted, residents experienced serious problem of inadequacy of water supply which resulted to high consumption of unhygienic water. However, sewages and contaminated water used for irrigation contained high level of Salmonella Typhi (mainly water-borne) that can be transferred to consumer through contaminated farm products [34].

In addition, in the study area many other reports indicated that multiple antibiotic resistances (MDRs) to Salmonella Typhi were among the most frequent salmonellae which are responsible for high infection among carriers and high opportunistic infections among AIDS patients. There is also diagnostic dilemma in the issue of chronic carriers, in the study area, the Schistosoma haematobium is common, and it was reported that a chronic urinary carrier state is resulted from localization of typhoid bacilli in schistosomiasis, the Schistosoma-Salmonella complex [41]. The Patient will become a variant that has chronic foci of infection but whom Salmonella Typhi is shed intravascularly rather than in stool or urine [41]. Similarly, as few as 10^3 mean infective dose of Salmonella Typhi organisms is required to produce clinical or subclinical infection in humans but perhaps 10^5 - 10^8 salmonellae is required for other salmonellae in African countries. Furthermore, waterborne disease outbreaks are often unreported or under reported in developing countries because of the lack of systematic studies. Presently and, indeed, in the future Kano may not be an exception in this regard [34].

On the other hand, less infection with Salmonella Typhimurium and Salmonella Paratyphi (mainly food-borne) could probably be because most food stuffs were washed properly or cooked before consumption and this help on the decrease or total destruction of Salmonella Typhimurium and Salmonella Paratyphi in food stuffs [42,43].

In this study, most of the Salmonella isolates from patients in Kano were significantly (X^2 11.03, P value = 0.001) resistant to ampicillin ($P < 0.05$) with 96 resistant strains representing 92.3% of the Salmonella isolates, followed by nalidixic acid with 81 resistant strains representing 77.9%. Ciprofloxacin remained effective against all the Salmonella isolates tested with exception of one (1) Salmonella isolate representing 1.0% resistance rate. However, the relationship between Salmonella serovars and antibiotics tested was not statistically significant ($p > 0.05$). Sixty five (62.5%) of the isolates displayed resistant to at least one or two antibiotics and thirty seven (35.6%) displayed resistant to three of antibiotics tested. Similarly, 37 isolates were found resistant to 1 of 3 first-line anti-Salmonella antibiotics (ampicillin, cotrimoxazole and chloramphenicol). Highest MARI was recorded from thirty seven (44.3%) isolates with MARI of 7.8 with two phenotypic patterns which differed in two different antibiotics (ampicillin and nalidixic acid). This work is in consonance with the findings of Asma et al. [39], Hemalatha et al. [44], Malla et al. [45], Gautum et al. [46] and Abdullahi et al. [36].

The highest significant resistance ($P < 0.05$) of Salmonella isolates to ampicillin and nalidixic acid could probably be due to the usage of antibiotics in the study area which is possibly the most important factor that promotes the emergence, selection and dissemination of antibiotic-resistant microorganisms in both veterinary and human medicine [44, 13]. However, the rate of development of resistance appears to have accelerated in the past decade and today multiple resistant Salmonella constitute a global problem [47]. It has been observed that antibiotic susceptibility of Salmonella isolates is not constant but dynamic and varies with time and environment. This therefore demands the need for periodic screening of common pathogens for their antibiotic susceptibility profiles in different communities. There is strong evidence that the use of antibiotics can lead to the emergence and dissemination of resistant salmonellae, which can then be passed onto people via food or through direct contact with animals. During recent years the wide spread use of antibiotics in the field of veterinary medicine have resulted in the development of increasing number of bacterial strains possessing resistance to many antibiotics. The property of multiple antibiotics resistance could be transferred through conjugation from resistant strains of salmonellae, to another by means of plasmid, which occur in cytoplasm of the donor bacterium and multiply independently of the chromosomal DNA. Thus a new bacterium with resistance factor emerges that is resistant to one or more antibiotics. In another instant the high resistance of Salmonella isolates to commonly used antibacterial drugs is probably due to some factors ranging from the use

of fake antibiotics, abuse and misuse of those antibiotics found commonly in circulation among the general populace and health resources centers [47, 46].

Abdullahi [36] reported that, acquired antibiotics resistance is a growing worldwide problem due to the increasing use of antibiotics in humans, animals, and agriculture. In developing countries the situation is particularly serious for the following reasons: In many countries, antibiotics can be obtained outside of recognized treatment centres, and taken without medical authorization or supervision. This leads to inappropriate use of antibiotics and their being taken at sub-optimal dosages and for an insufficient length of time. Often the high cost of an antibiotic, results in an incomplete course being purchased, sufficient only to alleviate symptoms. Patients are not sufficiently informed about antibiotics and their use [18,36]. Problems also arise when antibiotics sold in local markets are sub-standard or expired antibiotics. Guidelines regarding the selection of antibiotics, correct prescription, and information about antibiotic resistance and how to minimize its spread are not communicated to those purchasing the antimicrobials. Antibiotics are often prescribed when they are not needed or for self-limiting infections, e.g. diarrhoeal disease and viral respiratory infections [18, 36].

Other overlapping problems are worsening the situation regarding typhoid fever and other Salmonella infections within Africa: the failure to control the spread of the Salmonella species involved, due to unclean water, poor sanitation, malnutrition, the failure to control resistant organisms and resistance genes so that, when infections occur, they produce more adverse consequences. It is perhaps obvious, if unaddressed, that poor and displaced persons in Africa are least likely to be able to access potable water, safe sanitation, and other factors to prevent faecal-oral infection and that public health facilities need to be strengthened to protect the poor [48].

Broad spectrum antibiotics are frequently used prophylactically, e.g. ampicillin. Laboratory facilities for accurate diagnosis and isolation of pathogens are often not available, resulting in an overuse and inappropriate use of antibiotics [49]. Many countries do not have effective surveillance of important antibiotic-resistant bacteria. Training and facilities for performing standardized antibiotics sensitivity tests are often lacking. Developing countries are often unable to afford costly second-line antibiotics to treat infections due to resistant organisms. This results in prolonged illness with longer periods of infectivity and to the further spread of resistant strains [18].

V. Conclusion

In addition, the frequency of Salmonella infections was highest among 21-30 year age group lowest in ≥ 40 year age group. However the rates of infection among all the six (6) age groups were not significantly associated. The prevalent rate of Salmonella infections was significantly higher ($P > 0.05$) in males than the females patients with 68 (13.7%) and 36 (7.2%) respectively.

From the results of many studies conducted in different parts of the world, the Salmonella serovars that are frequently isolated in blood specimens of patients were also encountered in the present study. The study revealed that among the Salmonella serovars isolated, Salmonella Typhi was predominant followed by Salmonella Typhimurium, Salmonella Paratyphi C, B and A.

The result of in vitro – antibiotic susceptibility testing of Salmonella isolates using disc diffusion test demonstrated that most of the isolates (92.3%) was significantly ($P > 0.05$) resistant to ampicillin followed by nalidixic acid, chloramphenicol, and cotrimoxazole in decreasing order. However, ciprofloxacin remained effective against all the Salmonella isolates tested with exception of one Salmonella representing 1.0% resistance rate. However, the relationship between Salmonella serovars and antibiotics tested was not statistically significant ($P > 0.05$). In addition, 37 isolates were found resistant to 1 of 3 first-line anti-Salmonella antibiotics (ampicillin, cotrimoxazole and chloramphenicol). Highest MARI was recorded from thirty seven (44.3%) isolates with MARI of 7.8 with two phenotypic patterns which differed in two different antibiotics (ampicillin and nalidixic acid).

VI. Recommendation

Mass campaign, public enlightenment should be given more attention on the effective personnel hygiene, food hygiene and environmental sanitation. There is also need for reduced and appropriate consumption of antibiotics both for human and animals. Government should review existing antibiotic use policy to check abuse/misuse of antibiotics and ensure their correct prescription; this would reduce the rate of emergence of Salmonella resistant in particular and other bacteria in general. Research on development of new antibiotics and vaccines for treatment and prevention of Salmonella and other bacterial infections should be given more attention.

To reduce this antibiotic resistance, public health reference laboratory with a tool to produce standardized antibiotic susceptibility test results of antibiotic susceptibility tests are important for clinical treatment plans; adequate information must be provided to the health care providers. In addition, susceptibility testing help on providing guidance and monitor of treatment, narrower the spectrum of its antibiotic (the more proffered is it's use when one knows specifically the organism being treated), degree of susceptibility of

organism can assist in determining the length of therapy (but not the only factor) and choice of cheaper antibiotic agents with less side effects.

Consent

"All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images.

Ethical Approval

"All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki."

Acknowledgement

We thank all staff of the Management and Staff of Medical Laboratory Science Department, School of Health Technology, Kano, Department of Microbiology, Ahmadu Bello University, Zaria, Kaduna State, Biotechnology center Ahmadu Bello University Zaria, Kaduna State and Management and staff of all the six selected Hospitals.

Competing Interests

The authors declared that they have no competing interests exist.

References:

- [1.] Al –jurayyan, A. M., Al-Rasheedi, N. A. M., Mohammad, N., Alnasser, M. M., Mugeiren, A. T. and Abdullahi, S. A. Childhood bacterial diarrhoea in regional hospital in Saudi Arabia. *Journal of Tropical Medicine and Hygiene*. 2004; **97**: 87– 90.
- [2.] Agada, G. O. A., I. O. Abdullahi, M. Aminu, M. Odugbo, S. C. Chollom, P. R. Kumbish and E. J. Okwori. Prevalence and Antibiotic Resistance Profile of Salmonella Isolates from commercial Poultry and Poultry Farm-handlers in Jos, Plateau State, Nigeria. *British Microbiology Research Journal*. 2014; 4(4): 462-479.
- [3.] Poirel, L., Lagrutta, E., Taylor, P., Pham, J. and Nordmann, P. Emergence of metallo- β -lactamase NDM-1-producing multiple antibiotic-resistant *Escherichia coli* in Antimicrobiol Agents Chemotherapy, 2011; **4**: 4914-4916.
- [4.] Ellaine S, Robert M.H., Frederick J.A., Robert V.T., Marc-Alain W., Sharon L.R. *Emerging Infectious Disease*. 2011; 17(1).
- [5.] Wilson J. S., Hazel M. S., Williams J. N., Phiri A., French P. N., Hart C. A. Non-typhoid Salmonellae in United Kingdom Badgers: Prevalence and Spatial Distribution. *Applied Environmental Microbiology*. 2003; 69: 4312-4315.
- [6.] Ling J.M., Chan E.W., Cheng A.F. Molecular epidemiological analysis of Salmonella Enterica serotype Derby infections in Hong Kong. *Journal of Infection*. 2001; 42(2):145-153.
- [7.] Wright J.G., Tengelsen L.A., Smith K.E., Bender J.B., Frank P.K., Grendon J.H., Rice D.H., Thiessen A.M., Gilbertson C.J., Sivapalasingam S, Barret T.J., Besser T.E., Hancock D.D., Angulo F.J. Multidrug-resistance Salmonella Typhimurium in four animals' facilities. *Emerging Infectious Disease*. 2005; **11**(8): 1235-1241.
- [8.] Kariuki S., Revathi G., Kariuki N., Kiiru J., Mwituria J., Hart C.W. Characterization of community acquired nontyphoidal Salmonella from bacteraemia and diarrhoeal infection in children admitted to hospital in Nairobi, Kenya. *BMC Microbiol*. 2006; 6(101):1-10.
- [9.] Sahai S, Mahadevan S, Srinivasan S, Kanungo R. Childhood bacterial meningitis in Pondicherry, South India. *Indian Journal of Pediatrics*. 2001; 68:839-841.
- [10.] Aarii J., Tanabe Y., Miyake M. Acute encephalopathy associated with non-typhoid salmonellosis. *Journal Child Neurol*. 2001; 16: 539-540.
- [11.] Voetsch A. C., Van Gilder T. J., Angulo F. J., Farley M.M., Shallow S., Marcus R., Cieslak P.R., Deneen V.C., Tauxe R.V. Food Net estimate of the burden of illness caused by non-typhoid Salmonella infections in the United States. *CID*. 2004; 127–134.
- [12.] Muhammed M., Muhammed L.U., Ambali A.G., Mani A.U., Azard S., Barco L. Prevalence of Salmonella associated with chick mortality at hatching and their susceptibility to antimicrobial agents, *Veterinary Microbiology*. 2010; **140**: 131-135.
- [13.] Enabulele A.S., Amune P.O., Aborisade W.T. Antibigrams of Salmonella isolates from poultry farms in Ovia North East local government area, Edo State, Nigeria. *Agriculture and Biology Journal of North America*. 2010; **1**(6): 1287-1290.
- [14.] Agbaje M., Davies R., Oyekunle M.A., Ojo O.E., Fasina F.O., Akinduti P.A. Observation on the occurrence and transmission pattern of Salmonella Gallinarum in commercial poultry farms in Ogun State, South Western Nigeria. *African Journal Microbiological Research*. 2010; **4**(9): 796-800.
- [15.] Fasura A.K., Deji-Agboola A.M., Akinyemi K.O. Antimicrobial resistance patterns and Emerging fluoroquinolone resistant Salmonella isolates from poultry and asymptomatic poultry workers. *African Journal of Microbiology Research*. 2012; 6(11):2610-2615.
- [16.] Habtamu M.T., Rathore R., Dhama K., Rajesh K.A. Isolation, Identification and Polymerase Chain Reaction (PCR). Detection of Salmonella species from materials of Poultry origin. *International Journal of Microbiological Research*. 2011; **(2)**:135-142.
- [17.] Akinyemi, K. O., Smith, S. I., Oyefolu, A. O. and Coker, A. O. Multidrug resistance In Salmonella Typhi isolated from patients with typhoid fever complications in Lagos, Nigeria. *Journal of Public Health*, 2005; **119**: 321-327.
- [18.] Cheesbrough, M. *District laboratory practice in tropical countries*, ECBS edition Cambridge University Press, 2005; 2: 182-187.
- [19.] Arora, D. R. *Text book of Microbiology*. Second edition, New York, United Kingdom. 2006; Pp. 294-300.
- [20.] Threlfall, E. J., Day, M. and De-pinna, E. Antibiotic-resistant enteric fever in the United Kingdom; *Lancet*. 2006; **367**(9522): 1576.
- [21.] Parry, C. M., Hoa, N. T., Diep S., Wain, J., Chinh, N. T., Vinh, H., Hien, T. T., White, N. J. and Farrar, I. T. Value of a single-tube widal test in diagnosis of typhoid fever in Vietnam. *Journal of Clinical Microbiology*. 2006; **37**: 2882-2886.
- [22.] Okeke, N.I., Aboderin, O.A. and Byarugaba, D. K. *Journal of Emergent Infectious Diseases*, 2007; **13**(11): 1640-1646.
- [23.] Shekhar, P., Rajat, P., Deepak, J., Neelam, S.Amit, R., and Sandeep, N. The baseline widal titre among the healthy individuals of the hilly areas in the Garhwal region of Uttarakh and Indian Journal of Clinical and Diagnosis Research. 2013; **7**(3): 437–440.
- [24.] International Organization of Standardization (ISO) 6579. *Microbiology general guidelines on methods for the detection of Salmonella*. International organization of standardization, Geneva, Switzerland; 2002.

- [25.] Office International des Epizooties (OIE). Fowl typhoid and Pullorum disease. In: Terrestrial manual. Office International des Epizooties, Paris, France. 2012; 3-5.
- [26.] Kauffmann F. Serological diagnosis of Salmonella species, Kauffmann White Scheme Minkagarord, Copenhagen, Denmark; 2008; 1974.
- [27.] Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing; Twentieth Informational Supplement. CLSI document M100-S20. Wayne, PA: Clinical and Laboratory Standards Institute; 2010.
- [28.] Kirby – Bauer, O. S. Antimicrobial sensitivity testing by agar diffusion method. American Medical Journal of Clinical Pathology, 2008; **44**: 493-493.
- [29.] WHO. Manual for laboratory investigations of acute enteric infections. 2004; WHO. CDO/83.
- [30.] 30. World Health Organization (WHO). Traditional Medicine. In: Aliyu M. and Chedi B. A. Z. (editors). Effects of the ethanolic stem bark extract of *Pterocarpus Erinaceus* (Fabaceae) on some isolated smooth muscles. Bayero Journal of Pure and Applied Sciences, 2008; **3(1)**: 34-38.
- [31.] Heuer, O.E., Jensen, V.F., Hammerum, A.M. Antimicrobial antibiotic consumption in companion animals. Emerging Infectious Disease, 2005; **11(2)**: 344-345.
- [32.] Robert, H.C. and Jane, G.N.. Doing data analysis with Statistical Package for Social Science (SPSS) version 14. 2005; Pp 344.
- [33.] Adkins, H. J. and Santiago, L. T. Increased recovery of enteric pathogens by use of blood, stool and rectal swab specimen. Journal of Clinical Microbiology. 2006; **25**: 58 –59.
- [34.] Abdullahi, M. Incidence and antimicrobial susceptibility pattern of Salmonella species in children attending some hospitals in Kano metropolis, Kano state, Nigeria. Bayero Journal of Pure and Applied Sciences, 2010; **3(1)**: 10 –15.
- [35.] Mash, B. H. Typhoid fever: a case study of an endemic disorder in Katsina metropolis. Biological and Environmental Sciences Journal for the Tropics. 2013; **5(2)**: 27-30.
- [36.] Abdullahi, M., Olonitola, S. O. and Inabo, I. H. Isolation of Bacteria associated with diarrhoea among children attending some hospitals in Kano metropolis, Kano state, Nigeria. Bayero Journal of Pure and Applied Sciences. 2010; **3(1)**: 10 –15.
- [37.] Shamsuddeen, U., Bukar, A., Usman A. D., Kabir M. H. and Abdulmalik, S. A. Bacteriological quality of water used from ice making in some parts of Kano metropolis, Nigeria. Bayero Journal of Pure and Applied Sciences. 2010; **3(1)**: 199 – 201.
- [38.] Islam, A., Butler, T., Kabir, I. and Alam, N. H. Treatment of typhoid fever with ceftriaxone for 5 days of chloramphenicol for 14 days: a randomized clinical trial. Antibiotic Agents and Chemotherapy. 2008; **37**: 1572–1575.
- [39.] Asma, H., Abdul, H., Yasra, S., Aamir, A., Saira, B., Ayesha, T. and Mushkoor, M.. Identification of Drug resistance genes in clinical isolates of Salmonella Typhi for development of diagnostic multiplex polymerase chain reaction. Pakistan Journal of Medical Science, 2005; **21 (4)**: 402-407.
- [40.] Akinyemi, K. O., Coker, A. O., Olukoya, D. K., Oyefoly, A.O., Amorighoye, E. P. and Omonigbehim, E. O. Prevalence of multi-antibiotic resistant Salmonella Typhi Among clinically diagnosed typhoid fever patients in Lagos, Nigeria. Journal of Medical Laboratory Science. 2010; **13(2)**: 38-42.
- [41.] Hoshino, Y., Masuda, G., Negashi, M., Ajisawa, A., Imamura, N., Hachimori, K., Takayawa, N., Yamaguchi, T. and Kimura, T. Clinical bacteriology profiles of patients with typhoid fever treated in Tokyo metropolis. Journal of Medical Laboratory Science, 2010; **13(2)**: 38-42.
- [42.] Aarestrup, F. M., Lertworapreecha, M., Evans, M. C., Bangtrakulnonth, A., Chalermchaikit, T., Hendriksen, R. S. and Wagener, H. C. Antibiotic susceptibility and occurrence of resistance genes among Salmonella Enterica serovar Weltevreden from different countries. Journal of Antibiotic Chemotherapy. 2003; **52**: 715–718.
- [43.] Abdullahi, L. O. and Abdulkareem, S. Bacteriological quality of some ready to eat vegetables as retailed and consumed in Sabongari, Zaria, Nigeria. Bayero Journal of Pure and Applied Sciences. 2010; **3(1)**: 173-175.
- [44.] Hemalatha R., Vijayalakshmi P., Gyaneshwari, Rao M.V., Ramani A. Multidrug resistant Salmonella typhi in Hyderabad. Indian Journal of Medical Microbiology. 2008; **17**: 39–41.
- [45.] Malla, S., Kansaakar, P., Serichantalerg, S. Journal of Nepal Medical Association, 2008; **44**: 18-22.
- [46.] Gautam, V., Gupta, N. K., Choudhary, U., Arora, D. R. Sensitivity Pattern of Salmonella serotypes in Northern India. Brazilian Journal of Infectious Diseases, 2008; **6**: 281-287
- [47.] Pruden, A., Pei, R., Storteboom, H., et al. Environmental Science and Technology. 2006; **40**: 445-7450.
- [48.] Guha, S., Jalan, B. Y., Dey, S., Easow, J. M., Wilson, G. and Shivananda, P. G. Salmonella bacteraemia in Pokhara: emergence of antibiotic resistance. Nepal Medical College's Journal, 2005; **7**: 23–25.
- [49.] Kumurya, A. S. and Kawo, A. H. Quality assessment of some paediatric cotrimoxazole oral suspensions marketed in metropolitan Kano, Nigeria. Biological and Environmental Sciences Journal for the Tropics, 2010; **7(1)**: 26.