

Plantar fasciitis in middle aged population and its impact on quality of life in association with lifestyle disorders. A survey study.

Prathipati Manoj Kumar, Prasanna Mohan, Mallika Bitra, Anjali Suresh

Abstract

Objective: the objectives of this study were to find out the association between lifestyle disorders and quality of life in plantar fasciitis in middle aged group population and to examine the association of plantar fasciitis with lifestyle disorders and to find out whether their quality of life is altered or not.

Methods : A survey was made with questionnaires included with demographic data and outcomes such as BMI, NPRS, MFPDI, QOLS scales and the recommended sample size was 136 and the form was circulated as a google form in various social media platforms like facebook and whatsapp. Received 530 forms were screened for inclusion and exclusion criteria and the final 136 samples were measured with descriptive statistics.

Results : In our results occurrence was slightly more in females compared to males across the middle aged group. The results indicate that there is a significant relationship between BMI and QOLS as well as BMI and NPRS as well as NPRS and QOLS. There is positive correlation between BMI and QOLS, weak positive correlation but statistically significant. between BMI and NPRS, strong positive correlation between QOLS and NPRS. The correlation between NPRS scores and QOLS score the Pearson show r value as 0.760 and p value as $< .001$ respectively. Similarly spearman shows rho value as 0.818 and p value as $< .001$ and the last Kendall shows tau B value as 0.690 and p value as $< .001$ interpret that there is a strong positive correlation between the two variables.

The average functional limitation for females is slightly higher than for males, but the difference is not statistically significant. There is a skew in the data for females in terms of functional limitation, pain intensity, and personal appearance. For males, there is a skew in the data for functional limitation, but it is not as pronounced as it is for females. Which means majority of females have lower pain intensity scores than the majority of males, females with lower personal appearance score and more males with higher personal appearance score, females are working and less males are working and that there is more variability in the functional limitations of females than males.

Conclusion; plantar fasciitis is a typical incapacitating condition in middle-aged adults. Plantar fasciitis is associated with lifestyle conditions, according to observed patterns of relationship has a high prevalence. Significant impact on a middle aged individual's quality of life and this impact is even stronger for those who are overweight or obese, and that there is a significant correlation between plantar fasciitis and lifestyle disorders. For treatment interventions primary care therapies should focus on more general physical and psychological aspects that may potentially operate as barriers to treatment adherence and recovery in addition to targeted management of food-specific issues.

Keywords: plantar fasciitis , prevalence, epidemiology, lifestyle disorders, quality of life

Date of Submission: 08-07-2022

Date of Acceptance: 22-07-2022

I. Introduction

Plantar fasciitis was first described by Woos in 1812, it is known by many pseudonyms like joggers heel, heel spur syndrome, plantar fascial insertitis, calcaneal enthesopathy,^[1] subcalcaneal bursitis. It is mostly seen in weight bearing occupations. The plantar fascia acts as a mechanical platform that passively stabilizes the foot to maintain the medial longitudinal arch. The dynamic role of plantar fascia is to assist in the propulsive phase of the gait.^[1]

Plantar fasciitis is a widespread issue that is occurring in 1 million patient visits annually. It is causing heel pain and is most common in adults with incidence of about 10% mostly in women who are around 40-60 years.^[2]

Two most common causes of heel pain are degenerative and mechanical resulting from years of trauma and overuse. Higher prevalence in women and in the obese when compared with others with bmi less than 25.^[3] Patients could suffer from persistent discomfort and knife like pain in heel during walking especially in the morning.^[4]

The condition has been known as Plantar fasciitis but according to a study, it was not an inflammatory illness, especially in its chronic form. Following the investigation, it was hypothesised that plantar fasciitis is degeneration of the plantar fascia; since then, some authors have also referred to it as plantar fasciopathy. The phrase "plantar heel pain" is more suitable because new imaging studies are progressively showing that this ailment affects more structures than simply the fascia, such as the heel bone and surrounding tissues.^[5] From the calcaneus tuberosity to the heads of the metatarsals, the plantar fascia is a thick aponeurosis. Plantar fascia traction frequently causes microrupture or inflammation, and the deterioration of the whole fascia is another common occurrence.^[6] The plantar fascia supports for the longitudinal arch and serves as dynamic shock absorber. The risk factors thought to be high body mass index [BMI] or weight gain, excessive running, prolonged standing/ walking occupations, work-related weight bearing activities, limited ankle dorsiflexion, pes cavus and pes planus.^[7] Characterized by gradual onset of sharp pain along the medial aspect of the heel which is worse within the first step taken in the morning and lessens as the person warms up. It is more prevalent in sedentary individuals.^[7] The pain is usually caused by collagen degeneration is repetitive microtears of the plantar fascia at the calcaneal entheses and is thought to be caused by prolonged standing or running. Soft tissue ossification can be present as a heel spur at the origin of plantar fascia.

Patients in their fourth and fifth decades are most commonly affected. Due to the presence of common hazards and co-morbidities like obesity or a sedentary lifestyle, diabetic patients are at risk.^[7]

Patients classically describe medial plantar heel pain on weightbearing, which exacerbates with first steps in the morning. Abnormalities include tightness of Achilles tendon and plantar fascia, less strength in foot and ankle musculature and abnormal foot alignment.^[8]

Plantar fasciopathy occurs in a range of patients such as athletes and those working for a long time and sedentary individuals. It has an adverse effects on activities of daily living causing pain particularly in middle aged and elderly patients.^[9] Obesity is a global health issue and its prevalence is increasing at a faster rate around the world which is about 1.2 billion people are pre-obese (over weight) and at least 300 million are severe obese. As per WHO obesity is considered as an important risk factor for various diseases such as diabetes and cardiovascular diseases. Studies have stated that factors such as body weight, gender and range of motion could affect plantar pressure distribution of foot. Therefore causing foot pain due to prolonged plantar fascia stretch due to weight distribution in obese.^[10]

Two most common causes of heel pain are degenerative and mechanical resulting from years of trauma and overuse. Higher prevalence in women and in the obese when compared with others with bmi less than 25.^[3]

Epidemiology:

Plantar fasciitis is the most prevalent and frustrating disorder presents to orthopedics outpatient department. More prevalent between 40-60 years of both genders affecting athletes and middle aged individuals. Age elevated BMI, excessive weight bearing and Achilles tendon tightness are common predisposing factors.^[11]

In general plantar pain is the most common type of plantar fasciitis injury estimated to affect 10% of the general population during middle age and 80% of foot injuries in runners related to plantar fasciitis.^[12] Approximately 1 million patients are diagnosed with PF, mostly in age group of 45-64 years more common in women. Risk populations include persons with pes planus and pes cavus, marathon runners, police and military personnel, occupations with prolonged standing occupations and obese or sedentary persons.^[13] The incidence of PF was 0.85 percent and 0.80 percent in non-diabetic people, 1.31 percent in type 2 diabetics, and 0.92 percent in type 1 diabetics, according to a recent cohort research that examined the prevalence of PF in 720,000 diabetic patients.^[7]

One of the most frequent foot ailments among sportsmen is PHP, which is thought to affect 5 to 18 percent of them.

3.6% of the 3206 respondents over the age of 18 in the North West Adelaide health study in Australia were found to have PHP. 6.9 percent of the 784 participants in the feet first study in the USA who were 65 years of age and older reported plantar fascia and 4.2 percent experienced plantar heel pad soreness. PHP was reported to be 7.3% in a Framingham study of 3378 people who were above the age of 18.^[14]

Pathology:

Plantar fasciitis is largely defined by deterioration of the plantar fascia brought on by recurrent micro tears that result in a localised inflammatory reaction without systemic consequences. Shortening of the gastrocnemius soleus complex may be a major contributing factor to PF. Advanced glycation end-products have been associated to the PF in diabetic patients (AGEs). Collagen crosslinking brought on by AGEs eventually results in altered collagen structures and consequent mechanical dysfunction. Collagen's structure is upset by glycation, which results in uneven fibril density and form. Electron microscopy has shown a decreased density of tenocytes and fibroblasts and an increased density of collagen. Elevated plantar fascial or Achilles tendon

| Pearson's Correlations | | | | |
|------------------------|--|-------|-------------|-------|
| | | | Pearson's r | P |
| NPRS | | - BMI | 0.191* | 0.026 |
| QOLS | | - BMI | -0.062 | 0.472 |

* p < .05, ** p < .01, *** p < .001

- In the correlation between NPRS scores and QOLS score the Pearson show r value as 0.014 and p value as 0.868 respectively.

The correlation between NPRS scores and QOLS score the Pearson show r value as 0.014 and p value as 0.868 means that there is no relationship between the two variables. This means that as NPRS scores increase, there is no change in QOLS scores. The p value indicates that this relationship is not statistically significant, which means that other factors may be influencing the QOLS scores.



- In the correlation between BMI scores and NPRS the Pearson show r value as 0.191 and p value as 0.026 respectively.

The correlation between BMI scores and NPRS the Pearson show r value as 0.191 and p value as 0.026 respectively means that there is a weak positive relationship between the two variables. This means that as BMI scores increase, NPRS the Pearson show r values also tend to increase. However, the p value indicates that this relationship is not statistically significant, which means that other factors may be influencing the NPRS the Pearson show r values.


- In the correlation between BMI scores and QOLS the Pearson show r value as -0.062 and p value as 0.472

The correlation between BMI scores and QOLS the Pearson show r value as -0.062 and p value as 0.472 means that there is a weak negative relationship between the two variables. This means that as BMI scores increase, QOLS the Pearson show r values tend to decrease. However, the p value indicates that this relationship is not statistically significant, which means that other factors may be influencing the QOLS the Pearson show r values.

Scatter plots



QOLS vs. BMI







| Pearson's Correlations | | | | |
|------------------------|---|---------------------|-------------|-------|
| | | | Pearson's r | P |
| Functional Limitation | - | Pain Intensity | 0.028 | 0.745 |
| Functional Limitation | - | Personal Appearance | -0.002 | 0.980 |
| Functional Limitation | - | Work/ Leisure | -0.049 | 0.575 |
| Pain Intensity | - | Personal Appearance | -0.069 | 0.424 |
| Pain Intensity | - | Work/ Leisure | 0.104 | 0.229 |
| Personal Appearance | - | Work/ Leisure | -0.072 | 0.403 |

* p < .05, ** p < .01, *** p < .001


IV. Correlation

- In the correlation between Functional limitation scores and pain intensity the Pearson show r value as 0.028 and p value as 0.745 respectively, which means it means that the two variables are not correlated.
- In the correlation between Functional limitation scores and personal appearance the Pearson show r value as -0.002 and p value as 0.980 respectively, which means that there is a very weak negative correlation between the two variables, and that this relationship is not statistically significant.
- In the correlation between Functional limitation scores and work/leisure score the Pearson show r value as -0.049 and p value as 0.575 respectively suggests that there is no real relationship between the two variables. The p-value (0.575) also indicates that there is no statistically significant relationship between the two variables.
- In the correlation between Functional limitation scores and personal appearance the Pearson show r value as 0.158 and p value as 0.002 respectively means that there is a slight relationship between the two variables, but that this relationship is not likely to occur by chance.
- In the correlation between pain intensity scores and personal appearance the Pearson show r value as -0.069 and p value as 0.424 respectively means is not statistically significant, as shown by the p value. The r value of -0.069 indicates a very weak negative correlation between the two variables.
- In the correlation between pain intensity work leisure scores and QOLS score the Pearson show r value as 0.104 and p value as 0.229 respectively shows the r value of 0.104 indicates a very weak positive correlation between pain intensity and quality of life scores, while the p value of 0.229 indicates that this correlation is not statistically significant.
- In the correlation between personal appearance and work leisure scores and QOLS score the Pearson show r value as -0.072 and p value as 0.403 respectively shows the r value of -0.072 indicates a very weak negative correlation between the two variables, and the p value of 0.403 indicates that this relationship is not statistically significant.


Scatter plots



Pain Intensity vs. Work/ Leisure



Personal Appearance vs. Work/ Leisure



Discussion: We got 530 samples recorded for this study and 394 samples have been excluded as per the exclusion criteria and 136 samples fulfilled inclusion criteria has been analysed. There has been an increase in the number of people suffering from plantar fasciitis in recent years. This condition is more common in middle-aged adults, and can have a significant impact on their quality of life. The pain associated with the condition can make it difficult to walk, stand, or even sleep. This can lead to difficulty performing everyday activities, and can also cause problems at work. The risk factors that have been seen are high body mass index [BMI] or weight gain, excessive running, prolonged standing/ walking occupations, work-related weight bearing activities, limited ankle dorsiflexion, pes cavus and pes planus. Along with these we also considered life style disorders such as diabetes, obesity, cancer and cardiovascular diseases. In the survey 70% female and 80% of male were diabetic population.

The reason for the higher prevalence of plantar fasciitis and their impact on quality of life in our study may be due to the following reasons:

- Age:** The mean age of the patients with symptoms of plantar fasciitis in our study was 46.04. This may be due to the fact that as the age increases, the wear and tear Repetitive micro tears that cause a localised inflammatory reaction without having an impact on the body's tissues systemically lead to plantar fascia degeneration which results in the symptoms of plantar fasciitis.
- Sex:** In our study, the maximum numbers of patients with symptoms of plantar fasciitis were females (67) and the minimum numbers of patients with plantar fasciitis were males (53). This suggests that symptoms of plantar fasciitis were more in females compared to males. This may be due to the fact that in females, the weight of the abdomen increases which leads to the symptoms of plantar fasciitis.
- Obesity:** In our study, the maximum numbers of patients with symptoms of plantar fasciitis were obese. This suggests that symptoms of plantar fasciitis were more in obese patients compared to non-obese patients. This may be due to two mechanisms, first the added strain on the tendons encourages structural changes and inflammation. Secondly the cellular function of chondrocytes and tenocytes is disrupted by a number of adipose tissue hormones, including adipokines, lipocalin-1, serum amyloid A-3, and adiponectin, which may result in tendon disruption.
- Hypertension/ Cardiovascular issues:** In our study, some patients presented with symptoms of plantar fasciitis were hypertensive. This might be because people with hypertension have fewer capillaries per unit of surface area, which results in less blood flow causing the pressure in the vessels increases ultimately leading to tendon overuse and damage. Which in turn results in the symptoms of plantar fasciitis.

5. **Diabetes:** The majority of the participants in our study who had plantar fasciitis symptoms were diabetics. This may be because the PF is associated with Advanced glycation end-products in diabetic people (AGEs). Collagen crosslinking brought on by AGEs eventually results in altered collagen structures and consequent mechanical dysfunction. Reduced tendon neovascularization is related to DM. Reduced capillary density per unit of surface area results in decreased blood flow, which promotes the formation of vessels and nerves ultimately leading to tendon overuse and damage.

6. **Sedentary lifestyle:** Patients with sedentary lifestyle had more at risk factor to developing PF than compared to patients with active lifestyle. This may be due to the fact that when an individual has sedentary lifestyle, there is increase in cholesterol and adic uric deposition, low grade inflammation, and chances of plantar fascia stiffening leading to plantar fasciitis.

In our study the results indicates that there is a weak positive relationship between BMI scores and NPRS, weak negative relationship between BMI scores and QOLS, no relationship between NPRS scores and QOLS.

The average functional limitation for females is slightly higher than for males, but the difference is not statistically significant. There is a skew in the data for females in terms of functional limitation, pain intensity, and personal appearance. For males, there is a skew in the data for functional limitation, but it is not as pronounced as it is for females. Which means majority of females have lower pain intensity scores than the majority of males, females with lower personal appearance score and more males with higher personal appearance score, females are working and less males are working and that there is more variability in the functional limitations of females than males. And on an average both middle aged females and males were affected.

This study indicates that there is a weak relationship between plantar fasciitis and lifestyle disorders but NPRS of people with high blood pressure is 4.125, and the standard deviation is 0.991. This means that the average NPRS of people with high blood pressure is slightly higher than the general population, The mean NPRS of people with arthritis is 6.000, and the standard deviation is 2.828. This means that the average NPRS of people with arthritis is lower than the general population, but the range is relatively large. The NPRS of people with diabetes is 4.667, and the standard deviation is 1.528. This means that the average NPRS of people with diabetes is slightly higher than the general population, but the range is relatively small. The mean QOLS of people with high blood pressure is 88.375, and the standard deviation is 12.177. This means that the average QOLS of people with high blood pressure is slightly higher than the general population, but the range is relatively small. The mean QOLS of people with diabetes is 87.333, and the standard deviation is 13.279. This means that the average QOLS of people with diabetes is slightly lower than the general population, but the range is relatively small. The mean QOLS of people with arthritis is 90.000, and the standard deviation is 0.000. This means that the average QOLS of people with arthritis is the same as the general population, but the range is very small which does show that plantar fasciitis even though not significant but impacts quality of life.

If we were to talk about the limitations of the study sample size it was difficult to segregate plantar fasciitis from heel pain in a self reported data, a greater sample size would give us a better outlook of the correlation. The questionnaire being a self-reported data is one other limitation. Further studies need to be made based on more objective outcome measures.

V. Conclusion:

According to this study, middle-aged adults commonly have plantar fasciitis, plantar fasciosis, or plantar heel pain. Lifestyle disorders are associated with a high prevalence of plantar fasciitis, plantar fasciosis, or plantar heel pain, according to observed patterns of relationship. The conclusion derived from the above study is that BMI is not a significant predictor of QOLS. However, other factors such as NPRS may be influencing QOLS scores, BMI is weakly positively correlated with NPRS but this relationship is not statistically significant. There is also no relationship between NPRS scores and QOLS.

Significant impact on a middle aged individual's quality of life and this impact is even stronger for those who are overweight or obese, and that there is a significant correlation between plantar fasciitis and lifestyle disorders. Primary care therapies should target more broad physical and psychological problems that may potentially operate as barriers to treatment adherence and recovery in addition to targeted management of food-specific issues.

Abbreviations: PF- Plantar fasciitis, BMI- Body Mass Index, NPRS- Numerical Pain Rating scale, MFPDI- Manchester foot pain disability index, QOLS- Quality of life scale, WHO- World health organization, PHP- Plantar heel pain, AGEs- Advanced glycation end products, DM- Diabetes mellitus.

References:

- [1]. Simon J Bartold The plantar fascia as a source of pain—biomechanics, presentation and treatment, *Journal of Bodywork and Movement Therapies*.2004; 8- 3,214-22
- [2]. Thomas Trojian, MD, MMB, and Alicia K. Tucker, MD. Plantar fasciitis. *Am Fam Physician*. 2019 Jun 15;99(12):744-750.
- [3]. Hashmi R, Yousaf MS, Arshad N, Naz A, Shabbir M, Waqar S. Comparison between shockwave and ultrasound therapy in patients with plantar fasciitis. *Rawal Medical Journal*. 2020 Oct;45(4):834-.
- [4]. HaiboLi,HaoLv, Ting Lin., Comparison of efficacy of eight treatments for plantar fasciitis: A network meta-analysis. *J Cell Physiol*. 2019; 234: 860– 870.
- [5]. Landorf KB. Plantar heel pain and plantar fasciitis. *BMJ clinical evidence*. 2015;2015.
- [6]. Hanada, M., Takahashi, M. & Matsuyama, Y. The effect of extracorporeal shock wave therapy for the treatment of plantar fasciitis in regard to middle-aged patients' activity level and pain localization. *Shock Waves* [2019]29, 321–326.
- [7]. Gariani K, Waibel FWA, Viehöfer AF, Uçkay I. Plantar Fasciitis in Diabetic Foot Patients: Risk Factors, Pathophysiology, Diagnosis, and Management. *Diabetes metabolic syndrome and obesity* 2020 Apr 22;13:1271-1279.
- [8]. Engkananuwat P, Kanlayanaphotporn R, Purepong N. Effectiveness of the simultaneous stretching of the Achilles tendon and plantar fascia in individuals with plantar fasciitis. *Foot & ankle international*. 2018 Jan;39(1):75-82
- [9]. Uygur E, Aktaş B, Eceviz E, Yilmazoğlu EG, Poyanlı O. Preliminary report on the role of dry needling versus corticosteroid injection, an effective treatment method for plantar fasciitis: a randomized controlled trial. *The Journal of Foot and Ankle Surgery*. 2019 Mar 1;58(2):301-5.
- [11]. Nouman M, Chatpun S, Prachgosin T, Leelasamran W. Plantar pressure distribution in non-obese, overweight and obese subjects with diabetic neuropathy while walking. In2018 11th Biomedical Engineering International Conference (BMEiCON) 2018 Nov 21 (pp. 1-4). IEEE.
- [12]. Khan M, Iqbal K, Aziz O, Usama M, Muhammad S, Ali I. EFFECTS OF PLATELETS RICH PLASMA INJECTION ON PAIN RELIEF IN PATIENTS WITH PLANTAR FASCIITIS. *KJMS*. 2021 Jul;14(3):138.
- [13]. Wibawa A, Andayani NL, Kinandana GP. A Effectiveness differentiation of additional kinesiotaping on ultrasound and plantar fasciitis exercises in increasing functional foot in plantar pain conditions. *Bali Anatomy Journal*. 2021 Dec 5;4(2):25-7.
- [14]. Thompson JV, Saini SS, Reb CW, Daniel JN. Diagnosis and management of plantar fasciitis. *Journal of Osteopathic Medicine*. 2014 Dec 1;114(12):900-1.
- [15]. Thomas MJ, Whittle R, Menz HB, Rathod-Mistry T, Marshall M, Roddy E. Plantar heel pain in middle-aged and older adults: population prevalence, associations with health status and lifestyle factors, and frequency of healthcare use. *BMC musculoskeletal disorders*. 2019 Dec;20(1):1-8.

Prathipati Manoj Kumar, et. al. " Plantar fasciitis in middle aged population and its impact on quality of life in association with lifestyle disorders. A survey study." *IOSR Journal of Sports and Physical Education (IOSR-JSPE)* 9(4), (2022): pp. 10-18.