

## Evaluation of Cardiac Abnormalities by Echo Cardiographic Study In Non Haemorrhagic Cerebral Infarction

Dr. P. Thirumurugan, MD., D.M(Cardio)

Corresponding Author: Dr. A Mohamed Rafiq Babu, MD., D.M (Cardio)

---

### Abstract

**Background:** Strokes are a major cause of morbidity and mortality worldwide. Majority are due to ischemic cerebral infarction. Ischemia may result from occlusion of blood vessels due to a disease process intrinsic to carotid and intracranial vessels or may relate to coexisting heart diseases, predisposing to embolic phenomenon. **Objective:** Echocardiographic study in patients with cerebrovascular accident, confirmed to be of non hemorrhagic nature on CT scan/MRI, for evidence of potential cardiac abnormalities as a predisposing cause for the vascular event. **Material and Methods:** An observational study was conducted in department of medicine, Govt. Villupuram Medical College Hospital, Villupuram. 50 patients of ischemic stroke were taken and 2D transthoracic echocardiography was done to find potential cardioembolic abnormalities. **Results:** 76% of patients had echocardiographic abnormalities and 42% had potential cardioembolic abnormalities, out of which most common were ventricular wall hypokinesia, calcific aortic valve, rheumatic heart disease and dilated cardiomyopathy. The prevalence was almost similar in different age groups and both the sexes. **Conclusion:** Prevalence of potential cardioembolic abnormalities is high (42%) in ischemic stroke patients and 2D echocardiography is therefore recommended in the management and secondary prevention of cardioembolic stroke, which has a higher mortality and more chances of recurrence than atherothrombotic type of ischemic stroke.

**Key Words:** Stroke, echocardiography, potential cardioembolic abnormalities, cardioembolic stroke

---

Date Of Submission: 20-09-2018

Date of acceptance: 08-10-2018

---

### I. Introduction

Stroke is defined as an episode of acute neurological dysfunction presumed to be caused by ischemia or haemorrhage, persisting  $\geq 24$  hours or until death. CNS infarction is defined as brain, spinal cord, or retinal cell death attributable to ischemia, based on pathological, imaging, or other objective evidence of cerebral, spinal cord, or retinal focal ischemic injury in a defined vascular distribution; or clinical evidence of cerebral, spinal cord, or retinal focal ischemic injury based on symptoms persisting  $\geq 24$  hours or until death, and other etiologies excluded.[1] Strokes are a major cause of morbidity and mortality worldwide. [2] They may result from brain infarction or haemorrhage. Majority are due to ischemic cerebral infarction. [3] Ischemia may result from occlusion of blood vessels due to a disease process intrinsic to carotid and intracranial vessels or may relate to coexisting heart diseases, predisposing to embolic phenomenon. [4] About 20% of ischemic stroke and TIAs are probably due to embolism from the heart. The most common cause being non-rheumatic atria fibrillation.

Stroke is the leading cause of disability and the second most common cause of death worldwide. [6,7] Cardio embolic cerebral infarction accounts approximately for one quarter of all cerebral infarcts. [8] In most cases, recurrence of cardio embolism can be prevented by oral anticoagulants. Therefore, for a patient with a cerebral infarct, early confirmation of a diagnosis of cardio embolic cerebral infarction is extremely important in order to initiate anticoagulation therapy for an adequate secondary prevention. [9] The present study was conducted for finding the prevalence of cardioembolic abnormalities in patients of non hemorrhagic stroke.

### II. Material And Methods

The study was conducted in Govt. Villupuram medical college after taking approval from institutional thesis and ethical committee. 50 patients diagnosed as cerebrovascular accident, confirmed to be of non haemorrhagic nature on CT scan/MRI were taken. All patients were subjected to detailed history after taking consent and detailed systemic examination and investigations were performed. Patients diagnosed as cerebrovascular accident, confirmed to be of non haemorrhagic on CT scan/MRI were included in the study and those having intracranial hemorrhage and venous infarctions were excluded. This was a cross sectional single centre type of echocardiographic study of 50 patients with ischemic stroke undertaken to investigate the potential cardiac abnormalities as cause for ischemic stroke. A Trans Thoracic Echocardiography (TTE) examination was performed in all subjects using available portable Colour Doppler Echocardiography (CDE)

machine with an adult transducer of 2.5M Hz (Sonosite USA). Parasternal long-and short-axis, apical 4 chamber and two chamber views were obtained. All the measurements were done according to the American Society of Echocardiography guidelines and all the abnormalities were diagnosed according to the standard guidelines. All the significant cardiac findings on echocardiography were noted. Following are the potential cardioembolic abnormalities considered

- Rheumatic heart disease:
- Atrial fibrillation: Absence of 'A' wave on colour doppler.
- Infective/non infective endocarditis:
- Patent foramen ovale:
- Hypokinesia/akinesia of ventricular wall:
- Calcific aortic valve: An echo-dense lesion.
- LA/LV thrombus:
- Cardiac myxoma
- Prosthetic valve
- Atrial septal aneurysm
- Paradoxical embolism and congenital heart disease

The data was collected and analysed Chi square test was used.

### **III. Results**


This is a Cross sectional single centre type of echocardiographic study of 50 patients with non hemorrhagic cerebral infarction undertaken to investigate the potential cardiac abnormalities as cause for ischemic stroke. 50 patients diagnosed as cerebrovascular accident, confirmed to be of non haemorrhagic nature on CT scan/MRI were taken. Following observations were made the youngest patient was 32 years old and oldest 80 year old. The mean age of the patients was 56.42 with SD of 12.32. Maximum incidence was in 45-65 year age group. Both males and females were almost equally represented in the study.

In this study, an echocardiographic evaluation was done for all 50 patients included in the study. 12 (24%) patients had normal echocardiographic data, while 38 (76%) patients had some abnormal finding on echocardiography. Potential cardioembolic abnormality, identified by transthoracic echocardiography was found in 42% of the cases. Stroke incidence rates are generally relatively higher in males than females. In this study females constituted 56% of cases and males 44% of cases, the difference may be due to small sample size.


In our study ECG was found to be abnormal in 42 % cases. LVH was found to be the most common abnormality in 16% cases on ECG which is quite comparable to a study by Goldstein et al[11] in which LVH was found in 21.5 % cases. Atrial fibrillation was found in 13% of cases in that study compared to 4% in our study. Bundle branch blocks i.e. LBBB and RBBB were found to be in 6 % cases each as compared to 2 and 7 % in that same study In this study most common abnormalities found were left ventricular hypertrophy (34%), followed by diastolic dysfunction (32%), hypokinesia/akinesia of ventricles in 20% cases, valvular findings in 20% cases and systolic dysfunction in 16% of cases.

Potential cardioembolic abnormalities were present in 42% cases as compared to a study done by N. Uma et al.[12] at Lady Hardinge Medical College and Associated Smt. Sucheta Kriplani and Dr. Ram Manohar Lohia Hospitals, New Delhi where potential cardioembolic source was found in 27 (54%) patients. Most common abnormalities found were left ventricular wall hypokinesia in 13 (26%) followed by mitral regurgitation in 9(18%) and aortic valve calcification in (14%) patients. This difference may be due to small sample size (50 cases) in both the studies. Zenkers et al.[13] also reported a similar result and found a potential cardioembolic source in 50% patients evaluated by transthoracic echocardiography. Infarction secondary to cerebral embolism was diagnosed in 127 (23.5%) of 540 patients in the Michael Reese Stroke Registry.[14] Coronary artery disease, atrial fibrillation, valvular heart disease, mitral annulus calcification, and cardiomyopathy were the commonest etiologies. Echocardiography documented a potential embolic source in 7 patients without previously known heart disease and clarified the cardiac pathology in many of the patients with known heart disease. In a study by Sandercock P et al[10] 244 patients of cerebral infarction were studied.

Potential cardioembolic sources were found in 31 % of patients. Gagliardi et al[16] studied frequency of echocardiographic abnormalities in patients with ischemia of the carotid territory. Eighty-eight consecutive patients referred to a neurosurgical Department (63 men and 25 women) aged from 14 to 68 years, with cerebral ischemia in the carotid territory were taken. Only 5 (18%) of the 27 patients with abnormal angiograms had a potential cardiac source of emboli while 24 (39%) out of the remaining 61 patients had a potential cardiac source demonstrated at echocardiography.



**Fig.1 ECG findings**



**Fig.2 Echocardiographic findings**

In our study the Left ventricular ejection fraction was compared in patients with a potential cardioembolic abnormality and those without a potential cardio embolic abnormality. Ejection fraction was found to be significantly lower ( $p < 0.05$ ) in the group with potential cardio embolic abnormalities (56.06 %) than the group without potential cardio embolic abnormalities (41.14 %). Low EF was a risk factor for stroke in the multiethnic North Manhattan (NOMASS) population cohort, independently of age, sex and ethnicity; however, risk of stroke was not related to severity of EF reduction.[15]

#### IV. Conclusion

Strokes are a major cause of morbidity and mortality. More than 40% of ischemic stroke patients have potential cardioembolic abnormalities. Cardioembolic stroke has higher mortality and recurrence rates. 2D echocardiography is the main stay in diagnosing cardiac source of embolus. Many patients with no apparent clinical or ECG evidence of cardiac disease can be identified with echocardiography. 2D ECHO is recommended in every ischemic stroke patient to find potential cardioembolic abnormalities which will guide in the management of stroke, use of anticoagulants and secondary prevention of stroke

### References

- [1]. Sacco RL. An updated definition of stroke for the 21st century: statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2013 Jul;44(7):2064-89.
- [2]. Bamford J, Sandercock P, Dennis M, Burn J, Warlow C. Prospective study of acute cerebrovascular disease in the community: the Oxfordshire community stroke project 1981-86: Methodology, demography and incident cases of first ever stroke. *JNeurosurg Psych* 1998;51:1373-80.
- [3]. Kistler JP, Ropper AH, Martin JB. Cerebrovascular diseases. *HPIM* 1994;13(2):2233-56.
- [4]. Warlow CP. Cerebrovascular disease. *Oxford Text book of Medicine*. London: Oxford University press; 1988.p.155-61, 170.
- [5]. Ringlestein EB, Koschorke S, Holling A, Thron A, Lambertz H, Minale C. Computed tomographic patterns of proven embolic brain infarcts. *AnnNeurol* 1989;26:759-65.
- [6]. Weir NU. An update on cardioembolic stroke. *Postgrad Med J* 2008;84:133-42.
- [7]. Ferro JM. Brain embolism. Answers to practical questions. *JNeurol*. 2003;250:139-47.
- [8]. Sandercock PAG, Warlow CP, Jones LR, Starkey IR. Predisposing factors for cerebral infarction: the Oxfordshire community stroke project. *BMJ* 1989;298:80-2.
- [9]. Goldstein DS. The Electrocardiogram in Stroke: Relationship to Pathophysiological Type and
- [10]. N Uma. Echocardiography in patients with cerebral infarction. *JAPI* 1999;47:291-3.
- [11]. Zenker. Transesophageal echocardiography in young patients with cerebral ischaemic events. *Stroke* 1988;9:345-8.
- [12]. Registry. *Stroke* 2010;41(10):2223-8.
- [13]. Wang D, Liu M, Hao Z, Tao W, Lin S, Zhang S, et al. Features of acute ischemic stroke with rheumatic heart disease in a hospitalized Chinese population. *Stroke* 2012;43(11):2853-7.
- [14]. Gagliardi. Frequency of echocardiographic abnormalities in patients with ischaemia of the carotid territory – a preliminary report. *Stroke* 1985;16(1):118-21.

Dr. P. Thirumurugan. " Evaluation of Cardiac Abnormalities by Echo Cardiographic Study In Non Haemorrhagic Cerebral Infarction. " *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)*, vol. 17, no. 10, 2018, pp 21-24.